

Manfredo P. do Carmo

Differential Forms and Applications

微分形式及其应用



Springer

世界图书出版公司
www.wpcbj.com.cn

Manfredo P. do Carmo

Differential Forms and Applications

With 18 Figures

Springer-Verlag
Berlin Heidelberg New York
London Paris Tokyo
Hong Kong Barcelona
Budapest

图书在版编目 (CIP) 数据

微分形式及其应用: 英文 / (巴西) 杜卡莫著. —影印本. —北京: 世界图书出版公司北京公司, 2010. 2

书名原文: Differential Forms and Applications

ISBN 978-7-5100-0475-9

I. ①微… II. ①杜… III. ①微分几何—英文

IV. ①0186.1

中国版本图书馆 CIP 数据核字 (2010) 第 010580 号

书 名: Differential Forms and Applications

作 者: Manfredo P. do Carmo

中 译 名: 微分形式及其应用

责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河国英印务有限公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64021602, 010-64015659

电子信箱: kjb@wpcbj.com.cn

开 本: 24 开

印 张: 5.5

版 次: 2010 年 01 月

版权登记: 图字: 01-2009-1074

书 号: 978-7-5100-0475-9/0 · 690

定 价: 19.00 元

Manfredo P. do Carmo
Instituto de Matematica Pura e Aplicada (IMPA)
Estrada Dona Castorina, 110
22460-320 Rio de Janeiro
Brazil

This is a translation of the Portuguese book "Formas Diferenciais e Aplicações", first published by IMPA in 1971.

Mathematics Subject Classification (1991):
53-01, 53A05, 58A10, 58Z05, 70Hxx

ISBN 3-540-57618-5 Springer-Verlag Berlin Heidelberg New York
ISBN 0-387-57618-5 Springer-Verlag New York Berlin Heidelberg

Library of Congress Cataloging-in-Publication Data. Carmo, Manfredo. *Formas diferenciais e aplicações*. English. Differential forms and applications / Manfredo P. do Carmo. p. cm. -- (Universitext) Includes bibliographical references and index. ISBN 0-387-57618-5 1. Differential forms. I. Title. QA381.C2813 1994 515'.37--dc20 94-21965

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

Springer-Verlag Berlin Heidelberg New York
a member of Springer Science+Business Media
© Springer-Verlag Berlin Heidelberg 1994

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the Mainland China only and not for export therefrom.

To my friends around the world, without whose help neither this book nor its author would be seeing the light.

Preface

This is a free translation of a set of notes published originally in Portuguese in 1971. They were translated for a course in the College of Differential Geometry, ICTP, Trieste, 1989. In the English translation we omitted a chapter on the Frobenius theorem and an appendix on the nonexistence of a complete hyperbolic plane in euclidean 3-space (Hilbert's theorem). For the present edition, we introduced a chapter on line integrals.

In Chapter 1 we introduce the differential forms in R^n . We only assume an elementary knowledge of calculus, and the chapter can be used as a basis for a course on differential forms for "users" of Mathematics.

In Chapter 2 we start integrating differential forms of degree one along curves in R^n . This already allows some applications of the ideas of Chapter 1. This material is not used in the rest of the book.

In Chapter 3 we present the basic notions of differentiable manifolds. It is useful (but not essential) that the reader be familiar with the notion of a regular surface in R^3 .

In Chapter 4 we introduce the notion of manifold with boundary and prove Stokes theorem and Poincare's lemma.

Starting from this basic material, we could follow any of the possible routes for applications: Topology, Differential Geometry, Mechanics, Lie Groups, etc. We have chosen Differential Geometry. For simplicity, we restricted ourselves to surfaces.

Thus in Chapter 5 we develop the method of moving frames of Elie Cartan for surfaces. We first treat immersed surfaces and next the intrinsic geometry of surfaces

Finally, in Chapter 6, we prove the Gauss-Bonnet theorem for compact orientable surfaces. The proof we present here is essentially due to S.S.Chern. We also prove a relation, due to M. Morse, between the Euler characteristic of such a surface and the critical points of a certain class of differentiable functions on the surface.

As most authors, I am indebted to so many sources that it is hardly possible to acknowledge them all. Let me at least mention that the first four

VIII Preface

chapters were strongly influenced by the writings of my friend and colleague Elon Lima and the last two chapters bear the imprint of my teacher and friend S.S. Chern.

For the present version I am indebted to my colleagues M. Dajczer, L. Rodríguez and W. Santos for reading critically the manuscript and offering a number of useful suggestions. Special thanks are due to Lucio Rodríguez for his care in the camera ready presentation of the final text.

Rio de Janeiro, February 1994.

Manfredo Perdigão do Carmo

Universitext

- Aksoy, A., Khamsi, M. A.: *Methods in Fixed Point Theory*
Aupetit, B.: *A Primer on Spectral Theory*
Bachem, A., Kern, W.: *Linear Programming Duality*
Benedetti, R., Petronio, C.: *Lectures on Hyperbolic Geometry*
Berger, M.: *Geometry I*
Berger, M.: *Geometry II*
Blidtner, J., Hansen, W.: *Potential Theory*
Booss, B., Bleecker, D. D.: *Topology and Analysis*
Carleson, L., Gamelin, T.: *Complex Dynamics*
Carmo, M. P. do: *Differential Forms and Applications*
Cecil, T. E.: *Lie Sphere Geometry: With Applications of Submanifolds*
Chandrasekharan, K.: *Classical Fourier Transforms*
Charlap, L. S.: *Bieberbach Groups and Flat Manifolds*
Chern, S.: *Complex Manifolds without Potential Theory*
Chorin, A. J., Marsden, J. E.: *Mathematical Introduction to Fluid Mechanics*
Cohn, H.: *A Classical Invitation to Algebraic Numbers and Class Fields*
Curtis, M. L.: *Abstract Linear Algebra*
Curtis, M. L.: *Matrix Groups*
Dalen, D. van: *Logic and Structure*
Das, A.: *The Special Theory of Relativity: A Mathematical Exposition*
Devlin, K. J.: *Fundamentals of Contemporary Set Theory*
DiBenedetto, E.: *Degenerate Parabolic Equations*
Dimca, A.: *Singularities and Topology of Hypersurfaces*
Edwards, R. E.: *A Formal Background to Higher Mathematics I a, and I b*
Edwards, R. E.: *A Formal Background to Higher Mathematics II a, and II b*
Emery, M.: *Stochastic Calculus in Manifolds*
Foulds, L. R.: *Graph Theory Applications*
Frauenthal, J. C.: *Mathematical Modeling in Epidemiology*
Fuks, D. B., Rokhlin, V. A.: *Beginner's Course in Topology*
Gallot, S., Hulin, D., Lafontaine, J.: *Riemannian Geometry*
Gardiner, C. F.: *A First Course in Group Theory*
Gårding, L., Tambour, T.: *Algebra for Computer Science*
Godbillon, C.: *Dynamical Systems on Surfaces*
Goldblatt, R.: *Orthogonality and Spacetime Geometry*
Gouvêa, F. Q.: *p -Adic Numbers*
Hahn, A. J.: *Quadratic Algebras, Clifford Algebras, and Arithmetic Witt Groups*
Hájek, P., Havránek, T.: *Mechanizing Hypothesis Formation*
Hlawka, E., Schoißengeier, J., Taschner, R.: *Geometric and Analytic Number Theory*
Holmgren, R. A.: *A First Course in Discrete Dynamical Systems*
Howe, R., Tan, E. Ch.: *Non-Abelian Harmonic Analysis*
Humi, M., Miller, W.: *Second Course in Ordinary Differential Equations
for Scientists and Engineers*
Hurwitz, A., Kritikos, N.: *Lectures on Number Theory*
Iversen, B.: *Cohomology of Sheaves*
Kelly, P., Matthews, G.: *The Non-Euclidean Hyperbolic Plane*
Kempf, G.: *Complex Abelian Varieties and Theta Functions*

- Kloeden, P. E., Platen E., Schurz, H.: Numerical Solution of SDE
Through Computer Experiments
- Kostrikin, A. I.: Introduction to Algebra
- Krasnoselskii, M. A., Pokrovskii, A. V.: Systems with Hysteresis
- Luecking, D. H., Rubel, L. A.: Complex Analysis. A Functional Analysis Approach
- Ma, Zhi-Ming, Roeckner, M.: Dirichlet Forms
- Mac Lane, S., Moerdijk, I.: Sheaves in Geometry and Logic
- Marcus, D. A.: Number Fields
- McCarthy, P. J.: Introduction to Arithmetical Functions
- Meyer, R. M.: Essential Mathematics for Applied Fields
- Meyer-Nieberg, P.: Banach Lattices
- Mines, R., Richman, F., Ruitenberg, W.: A Course in Constructive Algebra
- Moise, E. E.: Introductory Problem Courses in Analysis and Topology
- Montesinos, J. M.: Classical Tessellations and Three-Manifolds
- Nikulin, V. V., Shafarevich, I. R.: Geometries and Groups
- Oden, J. J., Reddy, J. N.: Variational Methods in Theoretical Mechanics
- Øksendal, B.: Stochastic Differential Equations
- Porter, J. R., Woods, R. G.: Extensions and Absolute α -Hausdorff Spaces
- Rees, E. G.: Notes on Geometry
- Reisel, R. B.: Elementary Theory of Metric Spaces
- Rey, W. J. J.: Introduction to Robust and Quasi-Robust Statistical Methods
- Rickart, C. E.: Natural Function Algebras
- Rotman, J. J.: Galois Theory
- Rybakowski, K. P.: The Homotopy Index and Partial Differential Equations
- Samelson, H.: Notes on Lie Algebras
- Schiff, J. L.: Normal Families of Analytic and Mesomorphic Functions
- Shapiro, J. H.: Composition Operators and Classical Function Theory
- Smith, K. T.: Power Series from a Computational Point of View
- Smoryński, C.: Logical Number Theory I: An Introduction
- Smoryński, C.: Self-Reference and Modal Logic
- Stanišić, M. M.: The Mathematical Theory of Turbulence
- Stillwell, J.: Geometry of Surfaces
- Stroock, D. W.: An Introduction to the Theory of Large Deviations
- Sunada, T.: The Fundamental Group and the Laplacian (to appear)
- Sunder, V. S.: An Invitation to von Neumann Algebras
- Tamme, G.: Introduction to Étale Cohomology
- Tondeur, P.: Foliations on Riemannian Manifolds
- Verhulst, F.: Nonlinear Differential Equations and Dynamical Systems
- Zaanen, A. C.: Continuity, Integration and Fourier Theory

Table of Contents

Preface	vii
1. Differential Forms in \mathbb{R}^n	1
2. Line Integrals	17
3. Differentiable Manifolds	33
4. Integration on Manifolds; Stokes Theorem and Poincaré's Lemma	55
1. Integration of Differential Forms	55
2. Stokes Theorem	60
3. Poincaré's Lemma	66
5. Differential Geometry of Surfaces	77
1. The Structure Equations of \mathbb{R}^n	77
2. Surfaces in \mathbb{R}^3	82
3. Intrinsic Geometry of Surfaces	89
6. The Theorem of Gauss-Bonnet and the Theorem of Morse	99
1. The Theorem of Gauss-Bonnet	99
2. The Theorem of Morse	106
References	115
Index	117

1. Differential Forms in \mathbf{R}^n

The goal of this chapter is to define in \mathbf{R}^n “fields of alternate forms” that will be used later to obtain geometric results.

In order to fix the ideas, we will work initially with the three-dimensional space \mathbf{R}^3 .

Let p be a point of \mathbf{R}^3 . The set of vectors $q - p$, $q \in \mathbf{R}^3$ (that have origin at p) will be called the *tangent space of \mathbf{R}^3 at p* and will be denoted by \mathbf{R}_p^3 . The vectors $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$, $e_3 = (0, 0, 1)$ of the canonical basis of \mathbf{R}_0^3 will be identified with their translates $(e_1)_p, (e_2)_p, (e_3)_p$ at the point p .

A *vector field* in \mathbf{R}^3 is a map v that associates to each point $p \in \mathbf{R}^3$ a vector $v(p) \in \mathbf{R}_p^3$. We can write v as

$$v(p) = a_1(p)e_1 + a_2(p)e_2 + a_3(p)e_3,$$

thereby defining three functions $a_i: \mathbf{R}^3 \rightarrow \mathbf{R}$, $i = 1, 2, 3$, that characterize the vector field v . We say that v is *differentiable* if the functions a_i are differentiable.

To each tangent space \mathbf{R}_p^3 we can associate its *dual space* $(\mathbf{R}_p^3)^*$ which is the set of linear maps $\varphi: \mathbf{R}_p^3 \rightarrow \mathbf{R}$. A basis for $(\mathbf{R}_p^3)^*$ is obtained by taking $(dx_i)_p$, $i = 1, 2, 3$, where $x_i: \mathbf{R}^3 \rightarrow \mathbf{R}$ is the map which assigns to each point its i^{th} -coordinate. The set

$$\{(dx_i)_p; i = 1, 2, 3\}$$

is in fact the dual basis of $\{(e_i)_p\}$ since

$$(dx_i)_p(e_j) = \frac{\partial x_i}{\partial x_j} = \begin{cases} 0, & \text{if } i \neq j \\ 1, & \text{if } i = j. \end{cases}$$

Definition 1. A *field of linear forms* (or an *exterior form of degree 1*) in \mathbf{R}^3 is a map ω that associates to each $p \in \mathbf{R}^3$ an element $\omega(p) \in (\mathbf{R}_p^3)^*$; ω can be written as

$$\omega(p) = a_1(p)(dx_1)_p + a_2(p)(dx_2)_p + a_3(p)(dx_3)_p$$

or

$$\omega = \sum_{i=1}^3 a_i dx_i,$$

where a_i are real functions in \mathbf{R}^3 . If the functions a_i are differentiable, ω is called a *differential form of degree 1*.

Now let $\Lambda^2(\mathbf{R}_p^3)^*$ be the set of maps $\varphi: \mathbf{R}_p^3 \times \mathbf{R}_p^3 \rightarrow \mathbf{R}$ that are bilinear (i.e., φ is linear in each variable) and alternate (i.e., $\varphi(v_1, v_2) = -\varphi(v_2, v_1)$). With the usual operations of functions, the set $\Lambda^2(\mathbf{R}_p^3)^*$ becomes a vector space.

When φ_1 and φ_2 belong to $(\mathbf{R}_p^3)^*$, we can obtain an element $\varphi_1 \wedge \varphi_2 \in \Lambda^2(\mathbf{R}_p^3)^*$ by setting

$$(\varphi_1 \wedge \varphi_2)(v_1, v_2) = \det(\varphi_i(v_j))$$

The element $(dx_i)_p \wedge (dx_j)_p \in \Lambda^2(\mathbf{R}_p^3)^*$ will be denoted by $(dx_i \wedge dx_j)_p$. It is easy to see that the set $\{(dx_i \wedge dx_j)_p, i < j\}$ is a basis for $\Lambda^2(\mathbf{R}_p^3)^*$ (this will be proved in a more general setting in Proposition 1 below). Furthermore,

$$(dx_i \wedge dx_j)_p = -(dx_j \wedge dx_i)_p, \quad i \neq j,$$

and

$$(dx_i \wedge dx_i)_p = 0.$$

Definition 2. A *field of bilinear alternating forms* or an *exterior form of degree 2* in \mathbf{R}^3 is a correspondence ω that associates to each $p \in \mathbf{R}^3$ an element $\omega(p) \in \Lambda^2(\mathbf{R}_p^3)^*$; ω can be written in the form

$$\omega(p) = a_{12}(p)(dx_1 \wedge dx_2)_p + a_{13}(p)(dx_1 \wedge dx_3)_p + a_{23}(p)(dx_2 \wedge dx_3)_p$$

or

$$\omega = \sum_{i < j} a_{ij} dx_i \wedge dx_j, \quad i, j = 1, 2, 3,$$

where a_{ij} are real functions in \mathbf{R}^3 . When the functions a_{ij} are differentiable, ω is a *differential form of degree 2*.

We will now generalize the notion of differential form to \mathbf{R}^n . Let $p \in \mathbf{R}^n$, \mathbf{R}_p^n the tangent space of \mathbf{R}^n at p and $(\mathbf{R}_p^n)^*$ its dual space. Let $\Lambda^k(\mathbf{R}_p^n)^*$ be the set of all k -linear alternating maps

$$\varphi: \underbrace{\mathbf{R}_p^n \times \dots \times \mathbf{R}_p^n}_{k \text{ times}} \rightarrow \mathbf{R}$$

(alternating means that φ changes signs with the interchange of two consecutive arguments). With the usual operations, $\Lambda^k(\mathbf{R}_p^n)^*$ is a vector space. Given $\varphi_1, \dots, \varphi_k \in (\mathbf{R}_p^n)^*$, we can obtain an element $\varphi_1 \wedge \varphi_2 \wedge \dots \wedge \varphi_k$ of $\Lambda^k(\mathbf{R}_p^n)^*$ by setting

$$(\varphi_1 \wedge \varphi_2 \wedge \dots \wedge \varphi_k)(v_1, v_2, \dots, v_k) = \det(\varphi_i(v_j)), \quad i, j = 1, \dots, k.$$

It follows from the properties of determinants that $\varphi_1 \wedge \varphi_2 \wedge \dots \wedge \varphi_k$ is in fact k -linear and alternate. In particular $(dx_{i_1})_p \wedge (dx_{i_2})_p \wedge \dots \wedge (dx_{i_k})_p \in \Lambda^k(\mathbb{R}_p^n)^*$, $i_1, i_2, \dots, i_k = 1, \dots, n$. We will denote this element by $(dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k})_p$.

Proposition 1. *The set*

$$\{(dx_{i_1} \wedge \dots \wedge dx_{i_k})_p, \quad i_1 < i_2 < \dots < i_k, \quad i_j \in \{1, \dots, n\}\}$$

is a basis for $\Lambda^k(\mathbb{R}_p^n)^$.*

Proof. The elements of the set are linearly independent. For, if

$$\sum_{i_1 < \dots < i_k} a_{i_1 \dots i_k} dx_{i_1} \wedge \dots \wedge dx_{i_k} = 0,$$

is applied to

$$(e_{j_1}, \dots, e_{j_k}), \quad j_1 < \dots < j_k, \quad j_\ell \in \{1, \dots, n\},$$

we obtain (Exercise 2)

$$\sum_{i_1 < \dots < i_k} a_{i_1 \dots i_k} dx_{i_1} \wedge \dots \wedge dx_{i_k} (e_{j_1}, \dots, e_{j_k}) = a_{j_1 \dots j_k} = 0.$$

We now show that if $f \in \Lambda^k(\mathbb{R}_p^n)^*$, then f is a linear combination of the form

$$f = \sum_{i_1 < \dots < i_k} a_{i_1 \dots i_k} dx_{i_1} \wedge \dots \wedge dx_{i_k}.$$

For that, set

$$g = \sum_{i_1 < \dots < i_k} f(e_{i_1}, \dots, e_{i_k}) dx_{i_1} \wedge \dots \wedge dx_{i_k}.$$

Notice that $g \in \Lambda^k(\mathbb{R}_p^n)^*$ and that

$$g(e_{i_1}, \dots, e_{i_k}) = f(e_{i_1}, \dots, e_{i_k}),$$

for all i_1, \dots, i_k . It follows that $f = g$. Setting $f(e_{i_1}, \dots, e_{i_k}) = a_{i_1 \dots i_k}$, we obtain the above expression for f . \square

Definition 3. An *exterior k -form* in \mathbb{R}^n is a map ω that associates to each $p \in \mathbb{R}^n$ an element $\omega(p) \in \Lambda^k(\mathbb{R}_p^n)^*$; by Proposition 1, ω can be written as

$$\omega(p) = \sum_{i_1 < \dots < i_k} a_{i_1 \dots i_k}(p) (dx_{i_1} \wedge \dots \wedge dx_{i_k})_p, \quad i_j \in \{1, \dots, n\},$$

where a_{i_1, \dots, i_k} are real functions in \mathbb{R}^n . When the a_{i_1, \dots, i_k} are differentiable functions, ω is called a *differential k -form*.

For notational convenience, we will denote by I the k -upla (i_1, \dots, i_k) , $i_1 < \dots < i_k$, $i_j \in \{1, \dots, n\}$, and will use the following notation for ω :

$$\omega = \sum_I a_I dx_I.$$

We also set the convention that a differential 0-form is a differentiable function $f: \mathbb{R}^n \rightarrow \mathbb{R}$.

Example 1. In \mathbb{R}^4 we have the following types of exterior forms (where a_i, a_{ij} , etc., are real functions in \mathbb{R}^4):

0-forms, functions in \mathbb{R}^4 ,

1-forms, $a_1 dx_1 + a_2 dx_2 + a_3 dx_3 + a_4 dx_4$,

2-forms, $a_{12} dx_1 \wedge dx_2 + a_{13} dx_1 \wedge dx_3 + a_{14} dx_1 \wedge dx_4 + a_{23} dx_2 \wedge dx_3 + a_{24} dx_2 \wedge dx_4 + a_{34} dx_3 \wedge dx_4$,

3-forms, $a_{123} dx_1 \wedge dx_2 \wedge dx_3 + a_{124} dx_1 \wedge dx_2 \wedge dx_4 + a_{134} dx_1 \wedge dx_3 \wedge dx_4 + a_{234} dx_2 \wedge dx_3 \wedge dx_4$,

4-forms, $a_{1234} dx_1 \wedge dx_2 \wedge dx_3 \wedge dx_4$.

From now on, we will restrict ourselves to differential k -forms and we will call them simply k -forms.

We are going to define some operations on k -forms in \mathbb{R}^n .

First, if ω and φ are two k -forms:

$$\omega = \sum_I a_I dx_I, \quad \varphi = \sum_I b_I dx_I,$$

we can define their *sum*

$$\omega + \varphi = \sum_I (a_I + b_I) dx_I.$$

Next, if ω is a k -form and φ is an s -form, we can define their *exterior product* $\omega \wedge \varphi$, which is an $(s+k)$ -form, as follows.

Definition 4. Let

$$\omega = \sum_I a_I dx_I, \quad I = (i_1, \dots, i_k), \quad i_1 < \dots < i_k,$$

$$\varphi = \sum_J b_J dx_J, \quad J = (j_1, \dots, j_s), \quad j_1 < \dots < j_s.$$

By definition,

$$\omega \wedge \varphi = \sum_{IJ} a_I b_J dx_I \wedge dx_J.$$

Example 2. Let $\omega = x_1 dx_1 + x_2 dx_2 + x_3 dx_3$ be a 1-form in \mathbf{R}^3 and $\varphi = x_1 dx_1 \wedge dx_2 + dx_1 \wedge dx_3$ be a 2-form in \mathbf{R}^3 . Then, since $dx_i \wedge dx_i = 0$ and $dx_i \wedge dx_j = -dx_j \wedge dx_i$, $i \neq j$, we obtain

$$\begin{aligned}\omega \wedge \varphi &= x_2 dx_2 \wedge dx_1 \wedge dx_3 + x_3 x_1 dx_3 \wedge dx_1 \wedge dx_2 \\ &= (x_1 x_3 - x_2) dx_1 \wedge dx_2 \wedge dx_3.\end{aligned}$$

Remark 1. The definition of exterior product is made in such a way that if $\varphi_1, \dots, \varphi_k$ are 1-forms, then the exterior product $\varphi_1 \wedge \dots \wedge \varphi_k$ agrees with the k -form previously defined by

$$\varphi_1 \wedge \dots \wedge \varphi_k(v_1, \dots, v_k) = \det(\varphi_i(v_j)).$$

This follows immediately from the definition and will be left as an exercise (Exercise 3).

The exterior product of forms in \mathbf{R}^n has the following properties.

Proposition 2. Let ω be a k -form, φ be an s -form and θ be an r -form. Then:

- a) $(\omega \wedge \varphi) \wedge \theta = \omega \wedge (\varphi \wedge \theta)$,
- b) $(\omega \wedge \varphi) = (-1)^{ks}(\varphi \wedge \omega)$,
- c) $\omega \wedge (\varphi + \theta) = \omega \wedge \varphi + \omega \wedge \theta$, if $r = s$.

Proof. (a) and (c) are straightforward. To prove (b), we write

$$\omega = \sum_I a_I dx_I, \quad I = (i_1, \dots, i_k), \quad i_1 < \dots < i_k,$$

$$\varphi = \sum_J b_J dx_J, \quad J = (j_1, \dots, j_s), \quad j_1 < \dots < j_s.$$

Then

$$\begin{aligned}\omega \wedge \varphi &= \sum_{IJ} a_I b_J dx_{i_1} \wedge \dots \wedge dx_{i_k} \wedge dx_{j_1} \wedge \dots \wedge dx_{j_s} \\ &= \sum_{IJ} b_J a_I (-1) dx_{i_1} \wedge \dots \wedge dx_{i_{k-1}} \wedge dx_{j_1} \wedge dx_{i_k} \wedge \dots \wedge dx_{j_s} \\ &= \sum_{IJ} b_J a_I (-1)^k dx_{j_1} \wedge dx_{i_1} \wedge \dots \wedge dx_{i_k} \wedge dx_{j_2} \wedge \dots \wedge dx_{j_s}.\end{aligned}$$

Since J has s elements, we obtain, by repeating the above argument for each dx_{j_ℓ} , $j_\ell \in J$,

$$\begin{aligned}\omega \wedge \varphi &= \sum_{JI} b_J a_I (-1)^{ks} dx_{j_1} \wedge \dots \wedge dx_{j_s} \wedge dx_{i_1} \wedge \dots \wedge dx_{i_k} \\ &= (-1)^{ks} \varphi \wedge \omega.\end{aligned}$$

□

Remark 2. Although $dx_i \wedge dx_i = 0$, it is not true that for any form $\omega \wedge \omega = 0$. For instance, if

$$\omega = x_1 dx_1 \wedge dx_2 + x_2 dx_3 \wedge dx_4,$$

then

$$\omega \wedge \omega = 2x_1 x_2 dx_1 \wedge dx_2 \wedge dx_3 \wedge dx_4.$$

See however Exercise 4.

One of the most important features of differential forms is the way they behave under differentiable maps. Let $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a differentiable map. Then f induces a map f^* that takes k -forms in \mathbf{R}^m into k -forms in \mathbf{R}^n and is defined as follows. Let ω be a k -form in \mathbf{R}^m . By definition, $f^*\omega$ is the k -form in \mathbf{R}^n given by

$$(f^*\omega)(p)(v_1, \dots, v_k) = \omega(f(p))(df_p(v_1), \dots, df_p(v_k)).$$

Here $p \in \mathbf{R}^n$, $v_1, \dots, v_k \in \mathbf{R}_p^n$, and $df_p: \mathbf{R}_p^n \rightarrow \mathbf{R}_{f(p)}^m$ is the differential of the map f at p . We set the convention that if g is a 0-form,

$$f^*(g) = g \circ f.$$

We are going to show that the operation f^* on forms is equivalent to "substitution of variables". Before that, we need some properties of f^* .

Proposition 3. Let $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a differentiable map, ω and φ be k -forms on \mathbf{R}^m and $g: \mathbf{R}^m \rightarrow \mathbf{R}$ be a 0-form on \mathbf{R}^m . Then:

- a) $f^*(\omega + \varphi) = f^*\omega + f^*\varphi$,
- b) $f^*(g\omega) = f^*(g)f^*(\omega)$,
- c) If $\varphi_1, \dots, \varphi_k$ are 1-forms in \mathbf{R}^m , $f^*(\varphi_1 \wedge \dots \wedge \varphi_k) = f^*(\varphi_1) \wedge \dots \wedge f^*(\varphi_k)$.

Proof. The proofs are very simple. Let $p \in \mathbf{R}^n$ and let $v_1, \dots, v_k \in \mathbf{R}_p^n$. Then

- (a) $f^*(\omega + \varphi)(p)(v_1, \dots, v_k) = (\omega + \varphi)(f(p))(df_p(v_1), \dots, df_p(v_k)) = (f^*\omega)(p)(v_1, \dots, v_k) + (f^*\varphi)(p)(v_1, \dots, v_k) = (f^*\omega + f^*\varphi)(p)(v_1, \dots, v_k)$.
- (b) $f^*(g\omega)(p)(v_1, \dots, v_k) = (g\omega)(f(p))(df_p(v_1), \dots, df_p(v_k)) = (g \circ f)(p) \cdot f^*\omega(p)(v_1, \dots, v_k) = f^*g(p) \cdot f^*\omega(p)(v_1, \dots, v_k)$.
- (c) By omitting the indication of the point p , we obtain

$$\begin{aligned} f^*(\varphi_1 \wedge \dots \wedge \varphi_k)(v_1, \dots, v_k) &= (\varphi_1 \wedge \dots \wedge \varphi_k)(df(v_1), \dots, df(v_k)) \\ &= \det(\varphi_i(df(v_j))) = \det(f^*\varphi_i(v_j)) \\ &= (f^*\varphi_1 \wedge \dots \wedge f^*\varphi_k)(v_1, \dots, v_k). \end{aligned}$$

Remark 3. We will show below (See Proposition 4) that (c) holds not only for 1-forms but for k -forms as well.

We can now present the promised interpretation of f^* . Let (x_1, \dots, x_n) be coordinates in \mathbf{R}^n , (y_1, \dots, y_m) be coordinates in \mathbf{R}^m and let $f: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be written as

$$y_1 = f_1(x_1, \dots, x_n), \dots, y_m = f_m(x_1, \dots, x_n). \quad (*)$$

Let $\omega = \sum_I a_I dy_I$ be a k -form in \mathbf{R}^m . By using the above properties of f^* , we obtain

$$f^*\omega = \sum_I f^*(a_I)(f^*dy_{i_1}) \wedge \dots \wedge (f^*dy_{i_k}).$$

Since

$$f^*(dy_i)(v) = dy_i(df(v)) = d(y_i \circ f)(v) = df_i(v),$$

we have

$$f^*\omega = \sum_I a_I(f_1(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n)) df_{i_1} \wedge \dots \wedge df_{i_k},$$

where f_i and df_i are functions of x_j . Thus to apply f^* to ω is equivalent to "substitute" in ω the variables y_i and their differentials by the functions of x_k and dx_k obtained from $(*)$.

Remark 4. In various situations, it is convenient to use differential forms defined only on some open set $U \subset \mathbf{R}^n$ and not on the entire \mathbf{R}^n . It is clear that everything done so far extends trivially to this situation.

Example. (Polar coordinates). Let ω be the 1-form in $\mathbf{R}^2 - \{0, 0\}$ by

$$\omega = -\frac{y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy.$$

Let U be the set in the plane (r, θ) given by

$$U = \{r > 0; 0 < \theta < 2\pi\}$$

and let $f: U \rightarrow \mathbf{R}^2$ be the map

$$f(r, \theta) = \begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$$

Let us compute $f^*\omega$. Since

$$dx = \cos \theta dr - r \sin \theta d\theta,$$

$$dy = \sin \theta dr + r \cos \theta d\theta,$$

we obtain

$$\begin{aligned} f^*\omega &= -\frac{r \sin \theta}{r^2} (\cos \theta dr - r \sin \theta d\theta) + \frac{r \cos \theta}{r^2} (\sin \theta dr + r \cos \theta d\theta) \\ &= d\theta. \end{aligned}$$