


A. C. Davison, D.V. Hinkley

# Bootstrap Methods and their Application

自助法及其应用



CAMBRIDGE

世界图书出版公司  
[www.wpcbj.com.cn](http://www.wpcbj.com.cn)

# Bootstrap methods and their application

A. C. Davison

*Professor of Statistics, Department of Mathematics,  
Swiss Federal Institute of Technology, Lausanne*

D. V. Hinkley

*Professor of Statistics, Department of Statistics and Applied Probability,  
University of California, Santa Barbara*



**图书在版编目 (CIP) 数据**

自助法及其应用 = Bootstrap Methods and Their Application;  
英文 / (瑞士) 戴维森 (Davison, A. C.) 著. —影印本.

—北京: 世界图书出版公司北京公司, 2010. 5

(剑桥概率和统计)

ISBN 978-7-5100-0551-0

I. ①自… II. ①戴… III. ①统计分析—计算机  
模拟—英文 IV. ①C819

中国版本图书馆 CIP 数据核字 (2010) 第 036176 号

---

书 名: Bootstrap Methods and Their Application  
作 者: A. C. Davison, D. V. Hinkley

---

中 译 名: 自助法及其应用  
责任编辑: 高蓉 刘慧

---

出 版 者: 世界图书出版公司北京公司  
印 刷 者: 三河国英印务有限公司  
发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)  
联系电话: 010-64021602, 010-64015659  
电子信箱: kjb@wpcbj.com.cn

---

开 本: 16 开  
印 张: 37  
版 次: 2010 年 04 月  
版权登记: 图字: 01-2009-6575

---

书 号: 978-7-5100-0551-0/O · 766      定 价: 89.00 元

---

---

## Preface

The publication in 1979 of Bradley Efron's first article on bootstrap methods was a major event in Statistics, at once synthesizing some of the earlier resampling ideas and establishing a new framework for simulation-based statistical analysis. The idea of replacing complicated and often inaccurate approximations to biases, variances, and other measures of uncertainty by computer simulations caught the imagination of both theoretical researchers and users of statistical methods. Theoreticians sharpened their pencils and set about establishing mathematical conditions under which the idea could work. Once they had overcome their initial skepticism, applied workers sat down at their terminals and began to amass empirical evidence that the bootstrap often did work better than traditional methods. The early trickle of papers quickly became a torrent, with new additions to the literature appearing every month, and it was hard to see when would be a good moment to try to chart the waters. Then the organizers of COMPSTAT '92 invited us to present a course on the topic, and shortly afterwards we began to write this book.

We decided to try to write a balanced account of resampling methods, to include basic aspects of the theory which underpinned the methods, and to show as many applications as we could in order to illustrate the full potential of the methods — warts and all. We quickly realized that in order for us and others to understand and use the bootstrap, we would need suitable software, and producing it led us further towards a practically oriented treatment. Our view was cemented by two further developments: the appearance of two excellent books, one by Peter Hall on the asymptotic theory and the other on basic methods by Bradley Efron and Robert Tibshirani; and the chance to give further courses that included practicals. Our experience has been that hands-on computing is essential in coming to grips with resampling ideas, so we have included practicals in this book, as well as more theoretical problems.

As the book expanded, we realized that a fully comprehensive treatment was beyond us, and that certain topics could be given only a cursory treatment because too little is known about them. So it is that the reader will find only brief accounts of bootstrap methods for hierarchical data, missing data problems, model selection, robust estimation, nonparametric regression, and complex data. But we do try to point the more ambitious reader in the right direction.

No project of this size is produced in a vacuum. The majority of work on the book was completed while we were at the University of Oxford, and we are very grateful to colleagues and students there, who have helped shape our work in various ways. The experience of trying to teach these methods in Oxford and elsewhere — at the Université de Toulouse I, Université de Neuchâtel, Università degli Studi di Padova, Queensland University of Technology, Universidade de São Paulo, and University of Umeå — has been vital, and we are grateful to participants in these courses for prompting us to think more deeply about the

material. Readers will be grateful to these people also, for unwittingly debugging some of the problems and practicals. We are also grateful to the organizers of COMPSTAT '92 and CLAPEM V for inviting us to give short courses on our work.

While writing this book we have asked many people for access to data, copies of their programs, papers or reprints; some have then been rewarded by our bombarding them with questions, to which the answers have invariably been courteous and informative. We cannot name all those who have helped in this way, but D. R. Brillinger, P. Hall, M. P. Jones, B. D. Ripley, H. O'R. Sternberg and G. A. Young have been especially generous. S. Hutchinson and B. D. Ripley have helped considerably with computing matters.

We are grateful to the mostly anonymous reviewers who commented on an early draft of the book, and to R. Gatto and G. A. Young, who later read various parts in detail. At Cambridge University Press, A. Woollatt and D. Tranah have helped greatly in producing the final version, and their patience has been commendable.

We are particularly indebted to two people. V. Ventura read large portions of the book, and helped with various aspects of the computation. A. J. Canty has turned our version of the bootstrap library functions into reliable working code, checked the book for mistakes, and has made numerous suggestions that have improved it enormously. Both of them have contributed greatly — though of course we take responsibility for any errors that remain in the book. We hope that readers will tell us about them, and we will do our best to correct any future versions of the book; see its WWW page, at URL

<http://statwww.epfl.ch/davison/BMA/>

The book could not have been completed without grants from the UK Engineering and Physical Sciences Research Council, which in addition to providing funding for equipment and research assistantships, supported the work of A. C. Davison through the award of an Advanced Research Fellowship. We also acknowledge support from the US National Science Foundation.

We must also mention the Friday evening sustenance provided at the Eagle and Child, the Lamb and Flag, and the Royal Oak. The projects of many authors have flourished in these amiable establishments.

Finally, we thank our families, friends and colleagues for their patience while this project absorbed our time and energy. Particular thanks are due to Claire Cullen Davison for keeping the Davison family going during the writing of this book.

A. C. Davison and D. V. Hinkley  
Lausanne and Santa Barbara  
May 1997

---

# Contents

<i>Preface</i>	ix
<b>1 Introduction</b>	<b>1</b>
<b>2 The Basic Bootstraps</b>	<b>11</b>
2.1 Introduction	11
2.2 Parametric Simulation	15
2.3 Nonparametric Simulation	22
2.4 Simple Confidence Intervals	27
2.5 Reducing Error	31
2.6 Statistical Issues	37
2.7 Nonparametric Approximations for Variance and Bias	45
2.8 Subsampling Methods	55
2.9 Bibliographic Notes	59
2.10 Problems	60
2.11 Practicals	66
<b>3 Further Ideas</b>	<b>70</b>
3.1 Introduction	70
3.2 Several Samples	71
3.3 Semiparametric Models	77
3.4 Smooth Estimates of $F$	79
3.5 Censoring	82
3.6 Missing Data	88
3.7 Finite Population Sampling	92
3.8 Hierarchical Data	100
3.9 Bootstrapping the Bootstrap	103

3.10	Bootstrap Diagnostics	113
3.11	Choice of Estimator from the Data	120
3.12	Bibliographic Notes	123
3.13	Problems	126
3.14	Practicals	131
<b>4</b>	<b>Tests</b>	<b>136</b>
4.1	Introduction	136
4.2	Resampling for Parametric Tests	140
4.3	Nonparametric Permutation Tests	156
4.4	Nonparametric Bootstrap Tests	161
4.5	Adjusted P-values	175
4.6	Estimating Properties of Tests	180
4.7	Bibliographic Notes	183
4.8	Problems	184
4.9	Practicals	187
<b>5</b>	<b>Confidence Intervals</b>	<b>191</b>
5.1	Introduction	191
5.2	Basic Confidence Limit Methods	193
5.3	Percentile Methods	202
5.4	Theoretical Comparison of Methods	211
5.5	Inversion of Significance Tests	220
5.6	Double Bootstrap Methods	223
5.7	Empirical Comparison of Bootstrap Methods	230
5.8	Multiparameter Methods	231
5.9	Conditional Confidence Regions	238
5.10	Prediction	243
5.11	Bibliographic Notes	246
5.12	Problems	247
5.13	Practicals	251
<b>6</b>	<b>Linear Regression</b>	<b>256</b>
6.1	Introduction	256
6.2	Least Squares Linear Regression	257
6.3	Multiple Linear Regression	273
6.4	Aggregate Prediction Error and Variable Selection	290
6.5	Robust Regression	307
6.6	Bibliographic Notes	315
6.7	Problems	316
6.8	Practicals	321

<b>7</b>	<b>Further Topics in Regression</b>	<b>326</b>
7.1	Introduction	326
7.2	Generalized Linear Models	327
7.3	Survival Data	346
7.4	Other Nonlinear Models	353
7.5	Misclassification Error	358
7.6	Nonparametric Regression	362
7.7	Bibliographic Notes	374
7.8	Problems	376
7.9	Practicals	378
<b>8</b>	<b>Complex Dependence</b>	<b>385</b>
8.1	Introduction	385
8.2	Time Series	385
8.3	Point Processes	415
8.4	Bibliographic Notes	426
8.5	Problems	428
8.6	Practicals	432
<b>9</b>	<b>Improved Calculation</b>	<b>437</b>
9.1	Introduction	437
9.2	Balanced Bootstraps	438
9.3	Control Methods	446
9.4	Importance Resampling	450
9.5	Saddlepoint Approximation	466
9.6	Bibliographic Notes	485
9.7	Problems	487
9.8	Practicals	494
<b>10</b>	<b>Semiparametric Likelihood Inference</b>	<b>499</b>
10.1	Likelihood	499
10.2	Multinomial-Based Likelihoods	500
10.3	Bootstrap Likelihood	507
10.4	Likelihood Based on Confidence Sets	509
10.5	Bayesian Bootstraps	512
10.6	Bibliographic Notes	514
10.7	Problems	516
10.8	Practicals	519



<b>11 Computer Implementation</b>	<b>522</b>
11.1 Introduction	522
11.2 Basic Bootstraps	525
11.3 Further Ideas	531
11.4 Tests	534
11.5 Confidence Intervals	536
11.6 Linear Regression	537
11.7 Further Topics in Regression	540
11.8 Time Series	543
11.9 Improved Simulation	545
11.10 Semiparametric Likelihoods	549
<b>Appendix A. Cumulant Calculations</b>	<b>551</b>
<i>Bibliography</i>	555
<i>Name Index</i>	568
<i>Example index</i>	572
<i>Subject index</i>	575

---

## Introduction

The explicit recognition of uncertainty is central to the statistical sciences. Notions such as prior information, probability models, likelihood, standard errors and confidence limits are all intended to formalize uncertainty and thereby make allowance for it. In simple situations, the uncertainty of an estimate may be gauged by analytical calculation based on an assumed probability model for the available data. But in more complicated problems this approach can be tedious and difficult, and its results are potentially misleading if inappropriate assumptions or simplifications have been made.

For illustration, consider Table 1.1, which is taken from a larger tabulation (Table 7.4) of the numbers of AIDS reports in England and Wales from mid-1983 to the end of 1992. Reports are cross-classified by diagnosis period and length of reporting delay, in three-month intervals. A blank in the table corresponds to an unknown (as yet unreported) entry. The problem was to predict the states of the epidemic in 1991 and 1992, which depend heavily on the values missing at the bottom right of the table.

The data support the assumption that the reporting delay does not depend on the diagnosis period. In this case a simple model is that the number of reports in row  $j$  and column  $k$  of the table has a Poisson distribution with mean  $\mu_{jk} = \exp(\alpha_j + \beta_k)$ . If all the cells of the table are regarded as independent, then the total number of unreported diagnoses in period  $j$  has a Poisson distribution with mean

$$\sum_k \mu_{jk} = \exp(\alpha_j) \sum_k \exp(\beta_k),$$

where the sum is over columns with blanks in row  $j$ . The eventual total of as yet unreported diagnoses from period  $j$  can be estimated by replacing  $\alpha_j$  and  $\beta_k$  by estimates derived from the incomplete table, and thence we obtain the predicted total for period  $j$ . Such predictions are shown by the solid line in

Diagnosis period		Reporting delay interval (quarters):										Total reports to end of 1992
Year	Quarter	0†	1	2	3	4	5	6	...	≥14		
1988	1	31	80	16	9	3	2	8	...	6	174	
	2	26	99	27	9	8	11	3	...	3	211	
	3	31	95	35	13	18	4	6	...	3	224	
	4	36	77	20	26	11	3	8	...	2	205	
1989	1	32	92	32	10	12	19	12	...	2	224	
	2	15	92	14	27	22	21	12	...	1	219	
	3	34	104	29	31	18	8	6	...		253	
	4	38	101	34	18	9	15	6	...		233	
1990	1	31	124	47	24	11	15	8	...		281	
	2	32	132	36	10	9	7	6	...		245	
	3	49	107	51	17	15	8	9	...		260	
	4	44	153	41	16	11	6	5	...		285	
1991	1	41	137	29	33	7	11	6	...		271	
	2	56	124	39	14	12	7	10			263	
	3	53	175	35	17	13	11				306	
	4	63	135	24	23	12					258	
1992	1	71	161	48	25						310	
	2	95	178	39							318	
	3	76	181								273	
	4	67									133	

**Table 1.1** Numbers of AIDS reports in England and Wales to the end of 1992 (De Angelis and Gilks, 1994) extracted from Table 7.4. A † indicates a reporting delay less than one month.


Figure 1.1, together with the observed total reports to the end of 1992. How good are these predictions?

It would be tedious but possible to put pen to paper and estimate the prediction uncertainty through calculations based on the Poisson model. But in fact the data are much more variable than that model would suggest, and by failing to take this into account we would believe that the predictions are more accurate than they really are. Furthermore, a better approach would be to use a semiparametric model to smooth out the evident variability of the increase in diagnoses from quarter to quarter; the corresponding prediction is the dotted line in Figure 1.1. Analytical calculations for this model would be very unpleasant, and a more flexible line of attack is needed. While more than one approach is possible, the one that we shall develop based on computer simulation is both flexible and straightforward.

**Purpose of the Book**

Our central goal is to describe how the computer can be harnessed to obtain reliable standard errors, confidence intervals, and other measures of uncertainty for a wide range of problems. The key idea is to resample from the original data — either directly or via a fitted model — to create replicate datasets, from

Figure 1.1 Predicted quarterly diagnoses from a parametric model (solid) and a semiparametric model (dots) fitted to the AIDS data, together with the actual totals to the end of 1992 (+).



which the variability of the quantities of interest can be assessed without long-winded and error-prone analytical calculation. Because this approach involves repeating the original data analysis procedure with many replicate sets of data, these are sometimes called *computer-intensive methods*. Another name for them is *bootstrap methods*, because to use the data to generate more data seems analogous to a trick used by the fictional Baron Munchausen, who when he found himself at the bottom of a lake got out by pulling himself up by his bootstraps. In the simplest nonparametric problems we do literally sample from the data, and a common initial reaction is that this is a fraud. In fact it is not. It turns out that a wide range of statistical problems can be tackled this way, liberating the investigator from the need to oversimplify complex problems. The approach can also be applied in simple problems, to check the adequacy of standard measures of uncertainty, to relax assumptions, and to give quick approximate solutions. An example of this is random sampling to estimate the permutation distribution of a nonparametric test statistic.

It is of course true that in many applications we can be fairly confident in a particular parametric model and the standard analysis based on that model. Even so, it can still be helpful to see what can be inferred without particular parametric model assumptions. This is in the spirit of *robustness of validity* of the statistical analysis performed. Nonparametric bootstrap analysis allows us to do this.

3	5	7	18	43	85	91	98	100	130	230	487
---	---	---	----	----	----	----	----	-----	-----	-----	-----

**Table 1.2** Service hours between failures of the air-conditioning equipment in a Boeing 720 jet aircraft (Proschan, 1963).

Despite its scope and usefulness, resampling must be carefully applied. Unless certain basic ideas are understood, it is all too easy to produce a solution to the wrong problem, or a bad solution to the right one. Bootstrap methods are intended to help avoid tedious calculations based on questionable assumptions, and this they do. But they cannot replace clear critical thought about the problem, appropriate design of the investigation and data analysis, and incisive presentation of conclusions.

In this book we describe how resampling methods can be used, and evaluate their performance, in a wide range of contexts. Our focus is on the methods and their practical application rather than on the underlying theory, accounts of which are available elsewhere. This book is intended to be useful to the many investigators who want to know how and when the methods can safely be applied, and how to tell when things have gone wrong. The mathematical level of the book reflects this: we have aimed for a clear account of the key ideas without an overload of technical detail.

## Examples

Bootstrap methods can be applied both when there is a well-defined probability model for data and when there is not. In our initial development of the methods we shall make frequent use of two simple examples, one of each type, to illustrate the main points.


**Example 1.1 (Air-conditioning data)** Table 1.2 gives  $n = 12$  times between failures of air-conditioning equipment, for which we wish to estimate the underlying mean or its reciprocal, the failure rate. A simple model for this problem is that the times are sampled from an exponential distribution.

The dotted line in the left panel of Figure 1.2 is the cumulative distribution function (CDF)

$$F_{\mu}(y) = \begin{cases} 0, & y \leq 0, \\ 1 - \exp(-y/\mu), & y > 0, \end{cases}$$

for the fitted exponential distribution with mean  $\mu$  set equal to the sample average,  $\bar{y} = 108.083$ . The solid line on the same plot is the nonparametric equivalent, the empirical distribution function (EDF) for the data, which places equal probabilities  $n^{-1} = 0.08\bar{3}$  at each sample value. Comparison of the two curves suggests that the exponential model fits reasonably well. An alternative view of this is shown in the right panel of the figure, which is an exponential

**Figure 1.2** Summary displays for the air-conditioning data. The left panel shows the EDF for the data,  $\hat{F}$  (solid), and the CDF of a fitted exponential distribution (dots). The right panel shows a plot of the ordered failure times against exponential quantiles, with the fitted exponential model shown as the dotted line.



Q-Q plot — a plot of ordered data values  $y_{(j)}$  against the standard exponential quantiles

$$F_{\mu}^{-1} \left( \frac{j}{n+1} \right) \Big|_{\mu=1} = -\log \left( 1 - \frac{j}{n+1} \right).$$

Although these plots suggest reasonable agreement with the exponential model, the sample is rather too small to have much confidence in this. In the data source the more general gamma model with mean  $\mu$  and index  $\kappa$  is used; its density is

$$f_{\mu,\kappa}(y) = \frac{1}{\Gamma(\kappa)} \left( \frac{\kappa}{\mu} \right)^{\kappa} y^{\kappa-1} \exp(-\kappa y/\mu), \quad y > 0, \quad \mu, \kappa > 0. \quad (1.1)$$

For our sample the estimated index is  $\hat{\kappa} = 0.71$ , which does not differ significantly ( $P = 0.29$ ) from the value  $\kappa = 1$  that corresponds to the exponential model. Our reason for mentioning this will become apparent in Chapter 2.

Basic properties of the estimator  $T = \bar{Y}$  for  $\mu$  are easy to obtain theoretically under the exponential model. For example, it is easy to show that  $T$  is unbiased and has variance  $\mu^2/n$ . Approximate confidence intervals for  $\mu$  can be calculated using these properties in conjunction with a normal approximation for the distribution of  $T$ , although this does not work very well: we can tell this because  $\bar{Y}/\mu$  has an exact gamma distribution, which leads to exact confidence limits. Things are more complicated under the more general gamma model, because the index  $\kappa$  is only estimated, and so in a traditional approach we would use approximations — such as a normal approximation for the distribution of  $T$ , or a chi-squared approximation for the log likelihood ratio statistic.

The parametric simulation methods of Section 2.2 can be used alongside these approximations, to diagnose problems with them, or to replace them entirely. ■

**Example 1.2 (City population data)** Table 1.3 reports  $n = 49$  data pairs, each corresponding to a city in the United States of America, the pair being the 1920 and 1930 populations of the city, which we denote by  $u$  and  $x$ . The data are plotted in Figure 1.3. Interest here is in the ratio of means, because this would enable us to estimate the total population of the USA in 1930 from the 1920 figure. If the cities form a random sample with  $(U, X)$  denoting the pair of population values for a randomly selected city, then the total 1930 population is the product of the total 1920 population and the ratio of expectations  $\theta = E(X)/E(U)$ . This ratio is the parameter of interest.

In this case there is no obvious parametric model for the joint distribution of  $(U, X)$ , so it is natural to estimate  $\theta$  by its empirical analog,  $T = \bar{X}/\bar{U}$ , the ratio of sample averages. We are then concerned with the uncertainty in  $T$ . If we had a plausible parametric model — for example, that the pair  $(U, X)$  has a bivariate lognormal distribution — then theoretical calculations like those in Example 1.1 would lead to bias and variance estimates for use in a normal approximation, which in turn would provide approximate confidence intervals for  $\theta$ . Without such a model we must use nonparametric analysis. It is still possible to estimate the bias and variance of  $T$ , as we shall see, and this makes normal approximation still feasible, as well as more complex approaches to setting confidence intervals. ■

Example 1.1 is special in that an exact distribution is available for the statistic of interest and can be used to calculate confidence limits, at least under the exponential model. But for parametric models in general this will not be true. In Section 2.2 we shall show how to use parametric simulation to obtain approximate distributions, either by approximating moments for use in normal approximations, or — when these are inaccurate — directly.

In Example 1.2 we make no assumptions about the form of the data distribution. But still, as we shall show in Section 2.3, simulation can be used to obtain properties of  $T$ , even to approximate its distribution. Much of Chapter 2 is devoted to this.


## Layout of the Book

Chapter 2 describes the properties of resampling methods for use with single samples from parametric and nonparametric models, discusses practical matters such as the numbers of replicate datasets required, and outlines delta methods for variance approximation based on different forms of jackknife. It

**Table 1.3** Populations in thousands of  $n = 49$  large US cities in 1920 ( $u$ ) and in 1930 ( $x$ ) (Cochran, 1977, p. 152).

$u$	$x$	$u$	$x$	$u$	$x$
138	143	76	80	67	67
93	104	381	464	120	115
61	69	387	459	172	183
179	260	78	106	66	86
48	75	60	57	46	65
37	63	507	634	121	113
29	50	50	64	44	58
23	48	77	89	64	63
30	111	64	77	56	142
2	50	40	60	40	64
38	52	136	139	116	130
46	53	243	291	87	105
71	79	256	288	43	61
25	57	94	85	43	50
298	317	36	46	161	232
74	93	45	53	36	54
50	58				

**Figure 1.3** Populations of 49 large United States cities (in 1000s) in 1920 and 1930.





also contains a basic discussion of confidence intervals and of the ideas that underlie bootstrap methods.

Chapter 3 outlines how the basic ideas are extended to several samples, semiparametric and smooth models, simple cases where data have hierarchical structure or are sampled from a finite population, and to situations where data are incomplete because censored or missing. It goes on to discuss how the simulation output itself may be used to detect problems — so-called bootstrap diagnostics — and how it may be useful to bootstrap the bootstrap.

In Chapter 4 we review the basic principles of significance testing, and then describe Monte Carlo tests, including those using Markov Chain simulation, and parametric bootstrap tests. This is followed by discussion of nonparametric permutation tests, and the more general methods of semi- and nonparametric bootstrap tests. A double bootstrap method is detailed for improved approximation of P-values.

Confidence intervals are the subject of Chapter 5. After outlining basic ideas, we describe how to construct simple confidence intervals based on simulations, and then go on to more complex methods, such as the studentized bootstrap, percentile methods, the double bootstrap and test inversion. The main methods are compared empirically in Section 5.7, then there are brief accounts of confidence regions for multivariate parameters, and of prediction intervals.

The three subsequent chapters deal with more complex problems. Chapter 6 describes how the basic resampling methods may be applied in linear regression problems, including tests for coefficients, prediction analysis, and variable selection. Chapter 7 deals with more complex regression situations: generalized linear models, other nonlinear models, semi- and nonparametric regression, survival analysis, and classification error. Chapter 8 details methods appropriate for time series, spatial data, and point processes.

Chapter 9 describes how variance reduction techniques such as balanced simulation, control variates, and importance sampling can be adapted to yield improved simulations, with the aim of reducing the amount of simulation needed for an answer of given accuracy. It also shows how saddlepoint methods can sometimes be used to avoid simulation entirely.

Chapter 10 describes various semiparametric versions of the likelihood function, the ideas underlying which are closely related to resampling methods. It also briefly outlines a Bayesian version of the bootstrap.

Chapters 2–10 contain problems intended to reinforce the reader's understanding of both methods and theory, and in some cases problems develop topics that could not be included in the text. Some of these demand a knowledge of moments and cumulants, basic facts about which are sketched in the Appendix.

The book also contains practicals that apply resampling routines written in