

原色臺灣藥用植物圖鑑(5)



技 正 邱 年 永 共著
醫學博士 張 光 雄

中國醫藥學院中國藥學研究所

台北 南天書局 出版

797.75
20092
5

港 日 十

原 色 臺灣藥用植物圖鑑(5)

技 正 邱 年 永 共著
醫學博士 張 光 雄

中國醫藥學院中國藥學研究所



台 北 南 天 書 局 發 行

SMC PUBLISHING INC.
P.O. Box 13 342, Taipei 106
Taiwan, Republic of China

**THE
ILLUSTRATED
MEDICINAL PLANTS
OF TAIWAN**

VOL. 5

by

Nien-yung Chiu, Technician &

Kuang-hsiung Chang, M.D.

Institute of Chinese Pharmaceutical Science
China Medical College

250 Color Photographs

SMC PUBLISHING INC.
P.O. Box 13-342, Taipei 106
Taiwan, Republic of China

原色臺灣藥用植物圖鑑／邱年永，張光雄共著。--初

版。--臺北市：南天，民 75-87

冊： 公分。--（南天圖鑑系列：4-8）

參考書目：面

含索引

ISBN 957-638-062-6（一套：精裝）

ISBN 957-638-483-4（第 5 冊：精裝）

1. 藥用植物—圖錄

374.8025

83011746

南天圖鑑系列 ⑧ 原色臺灣藥用植物圖鑑 (5)

新台幣1000元

1998 年 7 月初版一刷發行

2004 年 8 月初版二刷發行

著作權所有
翻印必究

著 者：邱 年 永 · 張 光 雄

發 行人：魏 德 文

發 行 所：南 天 書 局 有 限 公 司

中華民國·台北市羅斯福路3段283巷14弄14號

☎(02)2362-0190 (代表號) Fax: (02)2362-3834

網 址：<http://www.smcbook.com.tw>

電子信箱：e-mail: weitw@smcbook.com.tw

郵政劃撥：01080538 號（南天書局帳戶）

原色製版：名亮彩色製版有限公司

☎(02)2952-5477 板橋市中山路2段400號2樓

原色印刷：皇甫彩藝印刷有限公司

☎(02)2303-5871 台北市長泰街297巷14號

ISBN 957-638-062-6 (套)

ISBN 957-638-483-4 (第 5 冊)

那 序

中國醫藥學院中國藥學研究所邱技正年永與藥學碩士張光雄二君，積數年之功，合著原色臺灣藥用植物圖鑑一書，內容簡明扼要而敘述正確，圖版鮮明，可謂為正統生藥學著作之一。邱君在中國醫藥學院藥學系藥用植物學科及中國藥學研究所從甘教授偉松兄專研藥用植物學，又復在本草學及生藥學之專研上，頗有進境，計在校服務十有餘年來，手不釋卷，專心於研究工作之進行，為人謙和，樂於助人，其人品頗為可敬。張光雄碩士，高雄醫學院藥學系畢業後，嘗服務於臺灣省政府衛生處藥政科，繼考入中國藥學研究所生藥學組，完成其學位論文後，負笈東京醫科大學藥理學教室，攻讀醫學博士(藥理學)學位，並已獲博士學位。二君與余朝夕相處，不僅相與論學，即平日待人處世，亦屬臭味相投，可謂相知最深，二君不辭辛勞，乃係利用公餘，完成此書，竊以為於我國藥用植物學、生藥學之教學研究日趨沒落式微之今日，深以推陳出新，釐正偏頗，而不失其傳統精神，今後生藥學研究之推展，可以預見，應二君之請，而樂為之序。

民國七十五年元月十日

藥學博士 那 琦謹識

甘 序

臺灣位于亞熱帶，氣候溫和，草木繁蔚，自古即視為產藥之區。自1765年，清·趙學敏撰「本草綱目拾遺」，始見咬人狗為本草之學所著錄，中外學人紛紛群起探研，始知臺灣產藥用植物多達兩千種。

邱君年永，來本科任職十餘年，勤奮好學，不但對藥用植物學之鑽研已入堂奧；亦追隨本草學權威那 琦博士勤習本草與生藥之學，已得那氏真傳；且隨植物化學大師魏吉恒教授遊，博覽群書。張君光雄，曾隨盧盛德博士研習生藥化學；復隨那琦教授專攻生藥學兩年，榮獲碩士學位，現負笈東瀛，追隨日本行動藥理學權威蒞谷健博士，修攻博士課程，今已榮獲醫學博士學位，載譽而歸。

今邱、張二君合著「原色臺灣藥用植物圖鑑」，依植物分類系統排列，首為中名（出典），內容計：科名、學名、別名、分布、產地、形態、採集與加工、成分、藥理、效用、方例、臨床，以及彙考等十三項，對臺灣產藥用植物之解說極為詳盡，彩照艷麗，圖文並茂，尤其不朽之作，紙貴洛陽，當可預卜，今慶巨著問世，念其辛勞，因樂為序。

中華民國七十五年元月十日

甘偉松 敬識于

中國醫藥學院藥用植物學科

魏 序

邱技正年永與藥學碩士張光雄君合著「原色臺灣藥用植物圖鑑」第二卷，計收集藥用植物二百五十種，皆為臺灣地區產野生或栽培植物之具有藥用價值者；每種均附有原色植物圖版以資對照，並依中名、學名、別名、分布、產地、形態、採集加工、成分藥理、效用、方例與彙考等項，分別作翔實之敘述；尤以原色圖片皆實地拍攝之五彩幻燈片予以放大精印，栩栩若生，極富真實感，俾讀者於野外採集時得能依地、按圖、恰如「探囊取物」之易於入手。邱技正與余共事近念年，知之最深；伊不分寒暑及星期例假，一直擔任指導學生藥用植物採集、生藥學實驗及本草學研究工作，達於「忘我」之境；曾陸續與那博士、甘教授及臺灣大學植物專家高木村先生發現臺灣新紀錄種植物多種，其對中國醫藥學院研究年報之編輯、出版，由第一期至今之第十二期更著辛勞，譯著尤夥。張光雄學友家學淵源，對生藥學造詣頗深，於中國藥學研究所以優異成績畢業後，即東渡日本，從東京醫科大學藥理學權威澁谷 健先生攻讀藥理學，獲得醫學博士學位。今合二人多年鑽研之心得，合著此書，一經付梓，紙貴可期，裨益中藥研究家與藥用植物同好及國民保健，豈淺鮮哉！余感其治學之勤，修研之誠，因樂予為序。

中華民國七十五年元月十五日

魏 吉 恆 序於中正嶺

許 序

邱年永君臺灣省南投縣人，自民國52年6月起在中國醫藥學院任技正，除擔任學生之藥用植物指導外，熱心研究藥用植物、生藥學及本草學等，有二十餘篇之原著論文並著有單行本：(一)藥用植物栽培法(二)原色野生食用植物圖鑑(三)重要漢方處方口訣集(四)漢方醫學等，對我國醫藥學術界貢獻甚鉅。張光雄碩士臺灣省臺中市人，高雄醫學院藥學系畢業，再考入中國醫藥學院中國藥學研究所，完成其學位論文，後再赴日本在東京醫科大學藥理學研究所攻讀博士課程。業已學成，榮獲醫學博士學位，載譽而歸。

最近邱年永技正與張光雄碩士合著「原色臺灣藥用植物圖鑑」一冊，求序於余，經審閱其內容，對取材及編排，多有獨到之處。從研究心得，搜集中外最新資料，根據基本原則，參照今後發展趨向與新觀念編述。全書收載臺灣常用藥用植物二百五十種並原色照片，分別學名、別名、分布、產地、形態、採集、加工、成分、藥理、效用、方例、彙考等之項目，均有簡明扼要之敘述，內容新穎，文筆清暢，不但在我國罕有之巨著，堪供國人尤以學生、藥用植物研究人士……為良好之參考與借鏡，相信對於國際學術界之貢獻，當非淺鮮，而樂為之序。

中華民國七十五年元月二日

醫學博士 許 喬 木 謹識

於行政院衛生署藥物食品檢驗局

自序

我國醫藥發達絕早，源遠流長，就中國傳統藥學之本草學(Pents'aology)，自漢代神農本草經以降，如魏晉之名醫別錄，南北朝之校定神農本草經、集注神農本草經。至唐之新修本草，宋之開寶本草、嘉祐本草、圖經本草、證類本草等具有一脈相承之體系，世謂之正統本草。其後金元之用藥法象、用藥心法、湯液本草。明清之本草品彙精要及續集、本草綱目、本草備要、本草從新、植物名實圖考及長編等爲本草著作之重心。我國本草始自漢魏、六朝，由早期本草，中經唐宋盛世，復歷金元之演變，至明清近代亘二千多年嬗遞傳承，作家約三百三十家，成書約二千又八十八卷，爲東方藥學之主流。

輒近西洋醫藥東漸，科學益趨昌明，國人趨之若鶩而篤信不移。近年來藥學專家不斷發現合成藥物，其中不乏具有強烈之副作用者，或往往導致人體組織之病變，甚至於有致癌之危機，因而世界各國率皆從事天然藥物之研究。對於我國傳統之本草學，未嘗接觸而認爲無研究之價值，鄙棄而不顧，每遇疾病認爲非服用化學藥品不爲功。惟觀夫阿片、顛茄、毛地黃等均爲植物性藥物，其醫療效果，實非純化學藥品所能代替者，至於當歸、柴胡、麻黃、大黃等藥材，迄今雖已獲悉其部分成分及藥理作用，但僅憑提取其中部分化學成分，無論如何仍不能完全評價其所有之藥效，此類蘊藏極豐之天然藥用資源，確有待於國人科學的求真精神，秉承我國歷代本草之優秀古代紀錄，進行其科學整理，繼以分工合作的精密研究，使國藥現代化，以期其裨益於世界人類之健康。

近年來藥用植物學界基於本草研究之精神，衍化爲現代本草學之內涵，對於醫藥資源之開發，已顯示其重要之地位。本書就臺灣地區野生或栽培之植物，網羅具有藥用價值者計收錄72科250種，分別擷取歷代諸家本草、藥用植物學、藥材學、生藥學、植物化學、藥理學、植物學等主要文獻編著成編，每種藥用植物均附以其植物彩色圖，以資對照易識，進而結合醫學臨床應用。除供醫藥學界研究外，並希望有助於植物學及農林學界之參考。並擬於本輯出版後，陸續完成全書五輯，將臺灣產絕大多數之藥用植物均予著錄，以期完備，顧作者等才疏學淺，勉力成書，謬漏之處，知所難免，尚望先進學者不吝指正爲禱。

本書之完成，承本草學、生藥學教授那琦博士，植物化學教授魏吉恆主任，藥用動物學教授許喬木博士之指導，並蒙分別賜序。復特蒙陳董事長立公資政，郭校長盛助博士，陳主任秘書勝智之勉勵與支持。本所陳忠川博士，以及謝所長明村博士、葉豐次博士、蔡輝彥博士、謝文全博士、張永勳博士、吳金濱博士、林文川博士之長期協助與支援。特別感謝鄭木榮醫師、張春霖道長、蔡吉雄醫師、魏國平醫師提供臨床資料。並承國立中國醫藥研究所高木村先生，本所彭文煌博士、闕甫仔博士、吳啓瑞博士。博士班侯鈺琪、何玉玲、闕甫仙、經總。碩士班陳啓源、林子超研究生之協助校稿。南天書局負責人魏德文藥師及其夫人宋玉蘭女士之爲本書鼎力策劃。作者等內人蔡政純女士，林秀珊女士之全力支持。謹此敬表由衷之謝忱。

中華民國八十七年六月

邱年永・張光雄 誌於
中國醫藥學院中國藥學研究所

PREFACE

Chinese pharmacology reached a developed stage quite early in recorded history. Traditional Chinese pharmacology or pents'aology began with the pharmacopoeia *Shen Nung pen-ts'ao*, which dates from the Han dynasty (206 B.C. - 220 A.D.). The tradition of pharmacopoeias continued to grow until modern times, with contributions being made in every important dynasty. We can now look back over an unbroken tradition of two thousand years, during which some 330 authors have contributed to a science which forms a mainstream of oriental medicine.

Although modern western medicine has come to occupy an unchallenged position throughout the oriental world, many newly discovered synthetic drugs produce harmful side-effects which may disturb the metabolism or even cause cancer. Research in countries all over the world is thus rapidly turning towards the search for natural drugs. Those unaware of traditional Chinese pharmacology as embodied in the pharmacopoeias have tended to neglect it as valueless for research, believing only in the effectiveness of chemical drugs. Yet the fact remains that the medicinal effects of such plant-based drugs as opium, scopolamine and digitalis cannot be reproduced by purely chemical drugs. Furthermore, in the case of such natural drugs as *Angelicae sinensis radix*, *Buplurei radix*, *Ephedrae herba* and *Rhei rhizoma*, only a proportion of their chemical constituents are known, and there is as yet no satisfactory way of evaluating their pharmacological effects as a whole. This rich source of natural medicines must await further co-ordinated research on the basis of the ancient records contained in the Chinese pharmacopoeias, research towards which this book is intended as a small contribution.

The present work brings together 250 species of wild and cultivated plants of medicinal value found in Taiwan. For each plant we have selected relevant information from the traditional pharmacopoeias and from the principal literature in the fields of medicinal botany, pharmacy, pharmacognosy, phytochemistry, pharmacology, botany, etc. A full-colour photograph of each plant is supplied to facilitate checking and recognition, and a summary is given of the plant's clinical functions. We hope that, pharmacologists apart, this information will also prove useful to botanists and to those engaged in agriculture and forestry. We intend eventually to complete a set of five uniform volumes which will attempt to record the medicinal plants of Taiwan in their entirety.

The authors are deeply grateful to a number of individuals for their assistance and advice during the compilation of the work, and our thanks are hereby tendered to all of them. For any mistakes and omissions which remain, however, the authors alone accept entire responsibility, and we earnestly hope that readers will draw our attention to such corrections as may be necessary.

Nien-yung Chiu & Kuang-hsiung Chang

June 1998

凡 例

1. 本書就臺灣地區野生或栽培之藥用植物，收錄93科 250 種。每種藥用植物均附以原色圖片以資對照。並分別就中名、學名、別名、日名、分布、產地、形態、採集、加工、成分、藥理、效用、方例、彙考各項，予以系統敘述。
2. 中名：依據中國歷代諸家本草及採用其相關典籍中最早出典之名稱為主，並附記有出典文獻名稱。其無出典之依據，則採用其通用名稱，並記其出現之地名，以求其名稱之統一。並附以植物名之羅馬拼音。
3. 學名：即拉丁 (Latin) 文植物學名，並附記其科名、屬名及種名。三者均用斜體字，至於命名者 (Author name)則用楷體字。
4. 別名：植物之別名極多而龐雜，故僅就古今名著所記載之植物名及地方性之通用名予以摘錄，均附記其出典及出現之地名。
5. 分布：記述植物之原產地及分布地區。
6. 產地：記述臺灣地區之出產地、分布情形，以及栽培狀態，引種歷史等，並就固有種植物，予以標示。
7. 形態：記述植物外部形態，明記其為木本、藤本或草本，植株之形狀、器官之大小，花之顏色及開花結果時期。
8. 採集·加工：明記其採收季節及方法，藥用部位，調製(修治)，乾燥及加工。
9. 成分·藥理：依有關文獻記述其已知成分，包括一般成分，供營養及藥用等項均予列入，以資參考。藥理則記載包括原生藥、成分之藥理作用機理、效果、毒性及解毒方法等。
10. 效用：列舉歷代諸家本草所錄之藥能以及民間經驗之療效，以及主治。
11. 方例：列舉歷代醫方本草、所錄成方，及地方藥誌等所傳錄之民間驗方，或臨床應用實例。方後並附記出典及地名。
12. 彙考：記述有關植物之出典、事蹟，掌故，考證等事項，以期增加研讀者之興趣。
13. 本書所錄、度量單位長度：如m(公尺)，cm(公分)，mm(公釐)。方例用量單位如兩(=10錢=31.25克)，錢(=10分=3.125克)、分(=10厘=0.3125克)。
14. 藥用植物名稱每有與藥材名相同者，本書就藥用植物學名、中文植物名(含藥材名)、日文植物名，編成三種索引，以便於檢索。
15. 本書之參考文獻甚多，除列舉主要參考文獻略名注解一覽表外，並於植物名、藥材名後附記其出典，以資參考。

INTRODUCTION

This work deals with medicinal plants, both wild and cultivated, found in Taiwan. 250 species are listed, in 93 families. A full-colour photograph of each plant is supplied, together with a systematic and detailed discussion in which the following information is given:

1. Chinese name, in both characters and Romanisation. The name given is that used in the earliest of the Chinese pharmacopoeias in which the plant is mentioned, whose authority is cited in each case. If no authority is available, the most common popular name has been applied, together with the locality where it is used, in an attempt to establish greater standardisation of Chinese plant names.
2. Scientific, or Latin, name, including the name of the family, species and genus to which the plant belongs. These appear in italic type, while the name of the person who named the species appears in standard type.
3. Alternative Chinese names. These are numerous and confused, and only those are recorded that appear in authoritative works or that have wide currency in popular usage in Taiwan. In each case the authority is cited or the locality where a popular name is current is indicated.
4. Japanese name, spelled out in the *katakana* syllabary.
5. Geographical distribution, including the areas to which the plant is native and those where it is found today.
6. Local growing districts, including the areas in Taiwan where the plant is produced, its distribution (e.g. as to elevation), the conditions under which it grows wild or can be cultivated, and the history of its transplantation.
7. Physical appearance. The external physical appearance of the plant is described, and a note is made of whether the plant is a wood, a vine or a grass.
The morphology of the plant, the size of its organs, the colour of the flowers and the times of flowering and fruiting are all noted.
8. Gathering and collection. The gathering season and method of collection of each plant are noted, together with the methods of drying, processing and preparing its medicinal parts.
9. The known medicinal and nutritional components of each plant are listed, according to the learned literature, and also its pharmacological uses, properties, mechanism and effects, including toxicity or effect as an antidote.
10. Medicinal application. The medicinal properties and functions of each plant are listed according to the historical Chinese pharmacopoeias, together with any medicinal uses and functions recognised in popular Chinese folk-medicine.
11. Prescription formulas. These are given according to the historical pharmacopoeias, together with prescriptions commonly applied in popular Chinese folk-medicine and attested to in local pharmacological gazetteers, or examples of clinical applications.
12. References to the learned literature, covering the *locus classicus* of the plant, pertinent historical facts and significant developments, and appropriate cross-references.
13. Weights and measures. Chinese units of weight are used in prescriptions, thus: 1 liang = ch'ien, or 31.25g; 1 ch'ien = 10 fen, or 3.125g; 1 fen = 10 li, or 0.3125g.
Metric units of length are used throughout, thus: m (metre), cm (centimetre), mm (millimetre).
14. Where names of plants are the same as the drugs derived from them, they are indexed separately under scientific name, Chinese botanical name and Japanese name.
15. Apart from a concise bibliography of principal authorities consulted, the relevant authorities are cited in each case after the name of the plant or drug.

First Published 1998
Republished 2004

Copyright © 1998 by SMC Publishing Inc.

SMC PUBLISHING INC.
P.O. Box 13-342, Taipei 106, Taiwan
Tel: (886-2) 2362-0190
Fax: (886-2) 2362-3834
e-mail: weitw@smcbook.com.tw
<http://www.smcbook.com.tw>

All rights reserved. No part of this book may be reproduced in any form, by printed or any other means without written permissions from the publisher.

ISBN 957-638-062-6 (set)
ISBN 957-638-483-4 (vol. 5)

PRINTED IN TAIWAN

原色臺灣藥用植物圖鑑(5)／目次

馬尾藻科

- 1 馬 尾 藻

多孔菌科

- 2 毛 革 蓋 菇

鳳尾蕨科

- 3 大 葉 鳳 尾 草
- 4 劍 葉 鳳 尾 草
- 5 鳳 尾 草
- 6 蜈 蚣 蕨
- 7 三 叉 鳳 尾 蕨

水龍骨科

- 8 石 韋
- 9 柔 軟 石 韋
- 10 掌 葉 石 韋
- 11 廬 山 石 韋

柏科

- 12 紅 檜

楊柳科

- 13 柳
- 14 水 柳

楊梅科

- 15 恆 春 楊 梅
- 16 楊 梅
- 17 銳 葉 楊 梅

胡桃科

- 18 楓 楊

桑科

- 19 天 竺 桑 麻
- 20 桑 麻
- 21 白 肉 榕
- 22 豬 腳 榕
- 23 澀 葉 榕
- 24 盤 龍 藤

蕁麻科

- 25 紫 麻

Sargassaceae

- Sargassum fulvellum* C. Agardh1

Polyporaceae

- Coriolus hirsutus* (Wulf. ex Fr.) Que'l.2

Pteridaceae

- Pteris cretica* Linn.3
Pteris ensiformis Burmann4
Pteris multifida Poiret5
Pteris vittata Linn.7
Pteris wallichiana Agardh8

Polypodiaceae

- Pyrrosia lingua* (Thunb.) Farwell.....9
Pyrrosia mollis (Kunze.) Ching11
Pyrrosia polydactylis (Hance) Ching12
Pyrrosia sheareri (Baker) Ching13

Cupressaceae

- Chamaecyparis formosensis* Matsumura14

Salicaceae

- Salix babylonica* Linn.15
Salix warburgii O. Seemann17

Myricaceae

- Myrica adenophora* Hance var. *kusanoi* Hayata18
Myrica rubra (Lour.) Sieb. et Zucc.19
Myrica rubra Sieb. & Zucc. var. *acuminata* Nakai21

Juglandaceae

- Pterocarya stenoptera* DC.22

Moraceae

- Fatoua pilosa* Gaudin23
Fatoua villosa (Thunb.) Nakai24
Ficus gibbosa Blume25
Ficus tinctoria Forster26
Ficus virgata Reinw. ex Blume27
Malaisia scandens (Lour.) Planchon28

Urticaceae

- Villebrunea frutescens* (Thunb.) Bl.29

山龍眼科	
26	山 龍 眼
檀香科	
27	百 蕊 草
山柑科	
28	山 柑 仔
蓼科	
29	苦 蕎 麥
30	何 首 烏
藜科	
31	臺 灣 沙 藜
32	鰾 蓬
莧科	
33	土 牛 膝
34	紫 莖 牛 膝
35	牛 膝
36	日 本 牛 膝
37	柳 葉 牛 膝
38	南 天 牛 膝
39	杯 莧
紫茉莉科	
40	黃 細 心
番杏科	
41	海 馬 齒 莧
馬齒莧科	
42	馬 齒 莧
43	白 馬 齒 莧
44	四 瓣 馬 齒 莧
石竹科	
45	蚤 綴
大戟科	
46	紫 錦 木
47	續 隨 子
48	大 戟 乳 脂 樹
49	守 宮 木
黃楊科	
50	細 葉 黃 楊
51	臺 灣 黃 楊

Proteaceae	
<i>Helicia formosana</i> Hemsley	30
Santalaceae	
<i>Thesium chinense</i> Turcz.	31
Opiliaceae	
<i>Champereia manillana</i> (Blume) Merr.	32
Polygonaceae	
<i>Fagopyrum tataricum</i> Gaertn	33
<i>Polygonum multiflorum</i> Thunb.	34
Chenopodiaceae	
<i>Atriplex nummularia</i> Lindl.	36
<i>Suaeda nudiflora</i> (Willd.) Moq.	37
Amaranthaceae	
<i>Achyranthes aspera</i> Linn. var. <i>indica</i> Linn.	38
<i>Achyranthes aspera</i> Linn. var. <i>rubrofusaca</i> Hooker f.	40
<i>Achyranthes bidentata</i> Blume	42
<i>Achyranthes japonica</i> (Miq.) Nakai	44
<i>Achyranthes longifolia</i> (Mak.) Makino	45
<i>Achyranthes ogatai</i> Yamamoto	46
<i>Cyathula prostrata</i> (Linn.) Blume	47
Nyctaginaceae	
<i>Boerhaavia diffusa</i> Linn.	48
Aizoaceae	
<i>Sesuvium portulacastrum</i> (L.) Linn.	49
Portulacaceae	
<i>Portulaca oleracea</i> Linn.	50
<i>Portulaca oleracea</i> Linn. var. <i>alba</i> Hong. N. V.	52
<i>Portulaca quadrifida</i> Linn.	53
Caryophyllaceae	
<i>Arenaria serpyllifolia</i> Linn.	54
Euphorbiaceae	
<i>Euphorbia continifolia</i> Linn.	55
<i>Euphorbia lathyris</i> Linn.	56
<i>Euphorbia resinifera</i> Berger	58
<i>Sauropus androgynus</i> Merr.	59
Buxaceae	
<i>Buxus harlandii</i> Hance	60
<i>Buxus microphylla</i> Sieb. et Zucc. var. <i>intermedia</i> (Kaneh) Li	61

木蘭科

- 52 辛 夷
- 53 木 蘭

小蘗科

- 54 細葉十大功勞
- 55 十 大 功 勞
- 56 南 天 竹

防己科

- 57 樟葉木防己
- 58 華南木防己
- 59 木 防 己
- 60 土 防 己
- 61 蘭 嶼 木 防 己
- 62 臺灣土防己
- 63 蓬 萊 藤
- 64 三脈蓬萊藤
- 65 毛 千 金 藤

十字花科

- 66 球 果 蔞 菜

景天科

- 67 伽 藍 菜
- 68 臺灣佛甲草
- 69 佛 甲 草

虎耳草科

- 70 心 基 葉 溲 疏
- 71 大 葉 溲 疏
- 72 臺 灣 溲 疏

海桐科

- 73 海 桐

薔薇科

- 74 日 本 木 瓜
- 75 枇 杷
- 76 黑 星 櫻

豆 科

- 77 蓮 實 藤
- 78 南 天 藤
- 79 臺灣杭子梢
- 80 小 果 刀 豆
- 81 肥 豬 豆

Magnoliaceae

- Magnolia denudata* Desr.62
- Magnolia liliflora* Desrousseauz.64

Berberidaceae

- Mahonia fortunei* (Lindl.) Fedde65
- Mahonia japonica* (Thunb.) DC.66
- Nandina domestica* Thunb.67

Menispermaceae

- Cocculus laurifolius* DC.69
- Cocculus sarmentosus* (Lour.) Diels70
- Cocculus trilobus* (Thunb.) DC.71
- Paracyclea gracillima* (Diels) Yamamoto72
- Paracyclea insularis* Kudo et Yamamoto73
- Paracyclea ochiaiana* Kudo et Yamamoto74
- Pericampylus formosanus* Diels75
- Pericampylus trinervatus* Yamamoto76
- Stephania hispidula* Yamamoto77

Cruciferae

- Rorippa globosa* (Turcz.) Thellung78

Crassulaceae

- Kalanchoe laciniata* (Linn.) DC.79
- Sedum formosanum* N. E. Brown80
- Sedum lineare* Thunb.81

Saxifragaceae

- Deutzia cordatula* Li82
- Deutzia pulchra* Vidal83
- Deutzia taiwanensis* (Maxim.) Schneidr84

Pittosporaceae

- Pittosporum tobira* (Thunb.) Aiton85

Rosaceae

- Chaenomeles japonica* (Thunb.) Lindley86
- Eriobotrya japonica* Linden87
- Prunus phaeosticta* (Hance) Maximowicz89

Leguminosae

- Caesalpinia globulorum* Bakh. F. et van Koyen90
- Caesalpinia nuga* (Linn.) Aiton91
- Campylotropis giraldii* (Schindler) Schindler92
- Canavalia cathartica* Thouin93
- Canavalia lineata* (Thunb.) DC.94

82	濱	刀	豆
83	鐵	刀	木
84	鋪	地	蝙蝠草
85	腎	葉	山螞蝗
86	美	洲	含羞草
87	金	甲	豆
88	黎		豆
89	胡	蘆	巴
水東哥科			
90	水	東	哥
田麻科			
91	小	葉	捕魚木
木棉科			
92	猢	猻	木
93	馬	拉	巴栗
錦葵科			
94	箭	葉	秋葵
95	風	鈴	花
96	臺	灣	磨盤草
97	苘		麻
98	圓	葉	金午時花
99	薄	葉	金午時花
100	東	方	賜米草
芸香科			
101	大	葉	三桠苦
102	金		柑
103	圓	金	柑
104	金		橘
105	四	季	橘
106	山	小	橘
107	小	山	橘
108	食	茱	萸
109	雙	面	刺
漆樹科			
110	埔		鹽
無患子科			
111	荔		枝
112	無	患	樹

<i>Canavalia maritima</i> (Aubl.) Piper	95
<i>Cassia siamea</i> Lamark	96
<i>Christia obcordata</i> (Poir.) Bahn. f.	97
<i>Desmodium renifolium</i> (Linn.) Schindler	98
<i>Mimosa invisa</i> Mart. et Colla.	99
<i>Phaseolus lunatus</i> Linn.	100
<i>Stizolobium hassjoo</i> Piper et Tracy	101
<i>Trigonella foenum-graecum</i> Linn.	102
Saurauiceae	
<i>Saurauia oldhamii</i> Hemsl	104
Tiliaceae	
<i>Grewia piscatorum</i> Hance	105
Bombacaceae	
<i>Adansonia digitata</i> Linn.	106
<i>Pachira macrocarpa</i> Schl	107
Malvaceae	
<i>Abelmoschus sagittifolius</i> (Kurz) Merr.	108
<i>Abutilon striatum</i> Dickson	109
<i>Abutilon taiwanensis</i> S. Y. Hu	110
<i>Abutilon theophrasti</i> Medikus	111
<i>Sida cordifolia</i> Linn.	112
<i>Sida mysorensis</i> Wight et Arnott	113
<i>Sida orientalis</i> Cavanilles	114
Rutaceae	
<i>Evodia triphylla</i> DC.	115
<i>Fortunella crassifolia</i> Swingle	116
<i>Fortunella japonica</i> (Thunb.) Swingle	117
<i>Fortunella margarita</i> (Lour.) Swingle	118
<i>Fortunella obovata</i> Tanaka	119
<i>Glycosmis citrifolia</i> (Willd.) Lindl.	120
<i>Glycosmis cochinchinensis</i> Pierre	121
<i>Zanthoxylum ailanthoides</i> Sieb. et Zucc.	122
<i>Zanthoxylum nitidum</i> (Roxb.) DC.	124
Anacardiaceae	
<i>Rhus semialata</i> Merr. var. <i>roxburghiana</i> DC.	126
Sapindaceae	
<i>Litchi chinensis</i> Sonnerat	128
<i>Sapindus mukorossi</i> Gaertner	130