

ALM 30

Advanced Lectures in Mathematics

Automorphic Forms and L -functions

自守形式与 L -函数

Editor: Jianya Liu



HIGHER EDUCATION PRESS

ALM 30

Advanced Lectures in Mathematics

Automorphic Forms and L -functions

自守形式与 L -函数

Editor: Jianya Liu

Copyright © 2014 by
Higher Education Press
4 Dewai Dajie, Beijing 100120, P. R. China, and
International Press
387 Somerville Ave, Somerville, MA, U. S. A.

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without permission.

图书在版编目 (CIP) 数据

自守形式与 L -函数 = Automorphic Forms and L -functions : 英文 / 刘建亚主编. -- 北京 : 高等教育出版社, 2014. 7

ISBN 978-7-04-039501-3

I. ①自… II. ①刘… III. ①自守形式-英文② L 函数-英文 IV. ①O174.54 ②O174.3

中国版本图书馆 CIP 数据核字 (2014) 第 069815 号

策划编辑 李华英
责任校对 王 雨

责任编辑 李华英
责任印制 张泽业

封面设计 张申申

版式设计 马敬茹

出版发行 高等教育出版社
社 址 北京市西城区德外大街4号
邮政编码 100120
印 刷 北京市四季青双青印刷厂
开 本 787mm×1092mm 1/16
印 张 14
字 数 340 千字
购书热线 010-58581118

咨询电话 400-810-0598
网 址 <http://www.hep.edu.cn>
<http://www.hep.com.cn>
网上订购 <http://www.landaco.com>
<http://www.landaco.com.cn>
版 次 2014 年 7 月第 1 版
印 次 2014 年 7 月第 1 次印刷
定 价 69.00 元

本书如有缺页、倒页、脱页等质量问题, 请到所购图书销售部门联系调换
版权所有 侵权必究
物 料 号 39501-00

ADVANCED LECTURES IN MATHEMATICS

EXECUTIVE EDITORS

Shing-Tung Yau
Harvard University
Cambridge, MA. USA

Lizhen Ji
University of Michigan
Ann Arbor, MI. USA

Kefeng Liu
University of California, Los Angeles
Los Angeles, CA. USA
Zhejiang University
Hangzhou, China

EXECUTIVE BOARD

Chongqing Cheng
Nanjing University
Nanjing, China

Tatsien Li
Fudan University
Shanghai, China

Zhong-Ci Shi
Institute of Computational Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

Zhiying Wen
Tsinghua University
Beijing, China

Zhouping Xin
The Chinese University of Hong Kong
Hong Kong, China

Lo Yang
Institute of Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

Weiping Zhang
Nankai University
Tianjin, China

Xiangyu Zhou
Institute of Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

Xiping Zhu
Sun Yat-sen University
Guangzhou, China



The CIMPA-UNESCO-CHINA Research School 2010: Automorphic Forms and L -functions

Preface

This is a collection of lecture notes from the CIMPA-UNESCO-CHINA Research School 2010: Automorphic Forms and L -functions, held at Shandong University at Weihai, in August 1–14, 2010.

The scientific directors were W. Duke and P. Sarnak (Chair) who provided detailed advice regarding the important aspects of the research school. The invited speakers were J. Cogdell, G. Harcos, E. Lapid, Xiaoqing Li, Wenzhi Luo, P. Michel, A. Reznikov, F. Shahidi, and Yangbo Ye.

Each of the speakers was scheduled to give a series of four lectures. Shahidi, although could not come to Weihai in person, has given full support to the research school by making his notes available, and Cogdell expertly combined his own lecture notes with Shahidi's material to give eight lectures. Not only have the speakers given excellent lectures, but they have also spent time completing their lecture notes – available for us here. Besides the speakers, there were fifteen scholars from all over the world plus ten international graduate students and over sixty Chinese graduate students who participated in various activities offered by the research school. The diligence and enthusiasm, professional skills and hard work of speakers, participants and organizers have contributed to the great success of the research school.

M. Waldschmidt visited Weihai in 2007 and provided clear instructions on how to organize a CIMPA school. M. Jambu, representative of CIMPA, not only participated the whole school, but also conducted lots of administrative duties on behalf of CIMPA before and after the school.

It is a great pleasure to express my sincere thanks to all the friends who have made this research school possible and who have contributed to its success. Thanks are also due to Lizhen Ji, Liping Wang, and Huaying Li for their effort and patience, and to Taiyu Li for helping me prepare the TeX file for publication. This event would not have been possible without the financial support from Shandong University, the NSFC, and CIMPA-UNESCO, and I would like to express my gratitude and thanks to them.

Jianya Liu
School Coordinator
July 2013

Academic Committee

William Duke, UCLA

Peter Sarnak (Chair), Princeton University and IAS

Speakers

Jim Cogdell, The Ohio State University

Gergely Harcos, The Hungarian Academy of Sciences

Erez Lapid, The Hebrew University of Jerusalem

Xiaoqing Li, SUNY at Buffalo

Wenzhi Luo, The Ohio State University

Philippe Michel, Ecole Polytechnique Fédérale de Lausanne

Andre Reznikov, Bar-Ilan University

Freydoon Shahidi, Purdue University

Yangbo Ye, The University of Iowa

School Coordinator

Jianya Liu, Shandong University

ADVANCED LECTURES IN MATHEMATICS

ADVANCED LECTURES IN MATHEMATICS

(Executive Editors: Shing-Tung Yau, Kefeng Liu, Lizhen Ji)

- 30. Automorphic Forms and L -functions (2014)
(Editor: Jianya Liu)
- 28–29. Selected Expository Works of Shing-Tung Yau with Commentary Vol. I, II (2013)
(Editors: Lizhen Ji, Peter Li, Kefeng Liu, Richard Schoen)
- 27. Number Theory and Related Area (2013)
(Editors: Yi Ouyang, Chaoping Xing, Fei Xu, Pu Zhang)
- 24–26. Handbook of Moduli Vol. I, II, III (2012)
(Editors: Gavril Farkas, Ian Morrison)
- 23. Recent Development in Geometry and Analysis (2012)
(Editors: Yuxin Dong, Jixiang Fu, Guozhen Lu, Weimin Sheng, Xiaohua Zhu)
- 22. Differential Geometry (2012)
(Editors: Yibing Shen, Zhongmin Shen, Shing-Tung Yau)
- 21. Advances in Geometric Analysis (2011)
(Editors: Stanisław Janeczko, Jun Li, Duong H. Phong)
- 20. Surveys in Geometric Analysis and Relativity (2011)
(Editors: Hubert L. Bray, William P. Minicozzi II)
- 19. Arithmetic Geometry and Automorphic Forms (2011)
(Editors: James Cogdell, Jens Funke, Michael Rapoport, Tonghai Yang)
- 17–18. Geometry and Analysis Vol. I, II (2010)
(Editor: Lizhen Ji)
- 16. Transformation Groups and Moduli Spaces of Curves (2010)
(Editors: Lizhen Ji, Shing-Tung Yau)
- 15. An Introduction to Groups and Lattices (2010)
(Author: Robert L. Griess, Jr.)
- 13–14. Handbook of Geometric Analysis Vol. II, III (2010)
(Editors: Lizhen Ji, Peter Li, Richard Schoen, Leon Simon)
- 12. Cohomology of Groups and Algebraic K -theory (2009)
(Editors: Lizhen Ji, Kefeng Liu, Shing-Tung Yau)
- 11. Recent Advances in Geometric Analysis (2009)
(Editors: Yng-Ing Lee, Chang-Shou Lin, Mao-Pei Tsui)
- 10. Trends in Partial Differential Equations (2009)
(Editors: Baojun Bian, Shenghong Li, Xu-Jia Wang)
- 9. Automorphic Forms and the Langlands Program (2009)
(Editors: Lizhen Ji, Kefeng Liu, Shing-Tung Yau, Zhujun Zheng)
- 8. Recent Developments in Algebra and Related Areas (2009)
(Editors: Chongying Dong, Fu-An Li)
- 7. Handbook of Geometric Analysis Vol. I (2008)
(Editors: Lizhen Ji, Peter Li, Richard Schoen, Leon Simon)
- 6. Geometry, Analysis and Topology of Discrete Groups (2008)
(Editors: Lizhen Ji, Kefeng Liu, Lo Yang, Shing-Tung Yau)
- 5. Proceedings of The 4th International Congress of Chinese Mathematicians Vol. I, II (2007)
(Editors: Lizhen Ji, Kefeng Liu, Lo Yang, Shing-Tung Yau)
- 4. Variational Principles for Discrete Surfaces (2007)
(Authors: Feng Luo, Xianfeng David Gu, Junfei Dai)

(Continued at the end of this volume)

Contents

<i>L</i>-functions and Functoriality	1
<i>James W. Cogdell</i>	
I <i>L</i> -functions for $GL(n)$ and Converse Theorems	
1 Modular Forms and Automorphic Representations	3
2 <i>L</i> -functions for GL_n and Converse Theorems	8
II <i>L</i> -functions via Eisenstein Series	
3 The Origins: Langlands	15
4 The Method: Langlands-Shahidi	21
5 The Results: Shahidi	26
III Functoriality	
6 Langlands Conjectures and Functoriality	30
7 The Converse Theorem and Functoriality	36
8 Symmetric Powers and Applications	41
References	47
Twisted Hilbert Modular <i>L</i>-functions and Spectral Theory	49
<i>Gergely Harcos</i>	
1 Lecture One: Some Quadratic Forms	49
2 Lecture Two: More Quadratic Forms	53
3 Lecture Three: Preliminaries from Number Theory	56
4 Lecture Four: Subconvexity of Twisted <i>L</i> -functions	61
Acknowledgments	64
References	64
The Voronoi Formula for the Triple Divisor Function	69
<i>Xiaqing Li</i>	
1 Introduction	69
2 Proof of the Main Theorem	71
Acknowledgments	89
References	89

Linnik's Ergodic Method and the Hasse Principle for Ternary**Quadratic Forms** 91*Philippe Michel*

1	Foreword	91
2	Integral Quadratic Forms	92
3	The Hasse Principle	93
4	Quadratic Forms over Lattices	96
5	Equidistribution on Adelic Quotient	101
6	Properties of the Adeles	106
7	The Hasse Integral Principle and Equidistribution of Adelic Orbits	114
8	The Ergodic Method	118
	References	129

Automorphic Periods and Representation Theory..... 131*Andre Reznikov*

1	Automorphic Representations and Frobenius Reciprocity	131
2	Bounds on Periods and Representation Theory	143
	Acknowledgments	147
	References	147

Eisenstein Series, L -functions and Representation Theory..... 149*Freydoon Shahidi*

1	Preliminaries	150
2	L -Groups, L -Functions and Generic Representations	151
3	Eisenstein Series and Intertwining Operators; The Constant Term	154
4	Constant Term and Automorphic L -Functions	156
5	Examples	160
6	Local Coefficients, Nonconstant Term and the Crude Functional Equation	160
7	The Main Induction, Functional Equations and Multiplicativity	162
8	Twists by Highly Ramified Characters, Holomorphy and Boundedness	166
9	Examples of Functoriality with Applications	168

10 Applications to Representation Theory	171
References	173

Lecture Notes on Some Analytic Properties of Automorphic

<i>L</i> -functions for $SL_2(\mathbb{Z})$	179
<i>Yangbo Ye</i>	
1 Introduction	179
2 An Integral Representation and Functional Equation	180
3 A Converse Theorem	188
4 The Phragmén-Lindelöf Principle and Convexity	193
5 The Rankin-Selberg Theory	197
References	202

L-functions and Functoriality

James W. Cogdell*

Abstract

The principle of functoriality is one of the central tenets of the Langlands program. There are two main approaches to functoriality, via the trace formula or via the method of *L*-functions. In this article we discuss the method of *L*-functions, pioneered by Piatetski-Shapiro. This method primarily deals with functoriality in the case where the target group is GL_n , the fundamental tool here being the converse theorem for GL_n . As a vehicle for functoriality, the input to the converse theorem must be checked, and this is done by controlling the *L*-functions of the automorphic representations to be transferred. For the cases presented here these *L*-functions are controlled by the Langlands-Shahidi method, which we outline as well. We discuss most of the major cases of functoriality that have been obtained by combining the converse theorem for GL_n and the Langlands-Shahidi method.

2010 Mathematics Subject Classification: 11F70, 11R39, 22E50, 22E55.

Keywords and Phrases: Automorphic forms, *L*-functions, Converse theorem, Eisenstein series, Langlands-Shahidi method, Functoriality.

The principle of functoriality is one of the central tenets of the Langlands program. It is a purely automorphic avatar of Langlands' vision of a non-abelian class field theory. One can find an outline of this in Section 5 below. There are two main approaches to functoriality. The one envisioned by Langlands is through the Arthur-Selberg trace formula. With the recent proof of the Fundamental Lemma by Ngô, Waldspurger, and others this method is now available and will be the subject of a forthcoming book of Jim Arthur [1]. The second method is that of *L*-functions.

The method of *L*-functions was pioneered by Piatetski-Shapiro. It primarily deals with functoriality in the case where the target group is GL_n . The fundamental tool here is the converse theorem for GL_n , as explained in Section 6 below. As

*Department of Mathematics, Ohio State University, Columbus, OH 43210, USA, cogdell@math.ohio-state.edu. The author was supported in part by NSF Grants DMS-0654017 and DMS-0968505.

Piatetski-Shapiro said “Arthur’s approach is more general, but this approach is much easier.” (see Shahidi’s contribution to [6]). The converse theorem is a way to tell when a representation of $GL_n(\mathbb{A})$ is automorphic based on the analytic properties of its L -functions (see Section 2 below). As a vehicle for functoriality, the input to the converse theorem must be checked, and this is done by controlling the L -functions of the automorphic representations to be functorially transferred. There are two ways to control these L -functions: via integral representations and via Eisenstein series, or the Langlands-Shahidi method. For the examples discussed here, these are controlled by the Langlands-Shahidi method, which we discuss in Sections 3–5. The functorial liftings themselves are obtained in Sections 7 and 8.

While Piatetski-Shapiro viewed this method as being simpler, it is in many ways more flexible and still mysterious. For example, as presented in Section 8, by this method you can obtain the third and fourth symmetric power lifting from GL_2 and these are still not attainable by the trace formula. In general, whenever you can control enough twisted L -functions, you can apply the converse theorem and obtain functoriality to GL_n . Of course controlling these L -functions is hard and for most of them we do not have a way to analyze them: they either fall outside the range of the Langlands-Shahidi method or we do not yet know if we can find integral representations for them. So there is still much work to be done.

These notes represent the lectures I gave at the CIMPA-UNESCO-CHINA Research School on *Automorphic Forms and L-functions* in August of 2010. Their purpose was to present an introduction to the L -function approach to functoriality. My original lectures were to be on GL_n and functoriality. These comprise Sections 1 and 2 here, which present the theory of automorphic representations and L -functions for GL_n , including the converse theorem, and then Sections 6 and 7 which were devoted to an exposition of functoriality via L -functions and then the example of functoriality for the classical groups. However, Shahidi was not able to attend the conference at the last moment, so I also gave an informal introduction to the Langlands-Shahidi method, based on Shahidi’s notes [10]. These appear as Sections 3–5 and 8 here. These sections can be viewed as a “gentle introduction” to [10] and I have tried to indicate where the results here can be found in Shahidi’s contribution. However, having the opportunity to give a single self-contained introduction to our approach to functoriality by the method of L -functions, I took the opportunity to integrate both series of lecture into a single contribution. I hope the reader finds this useful.

A word on references. I have surveyed the material in Sections 1, 2, 6 and 7 many times. Rather than burden this set of informal notes with pages of references, I refer instead to the sources I used for these talks, which are mainly my previous surveys. One can find more extensive references there. Similarly, for Sections 3–5 and 8 I have included in the bibliography the sources that I have used, which were surveys by Kim and Shahidi, Shahidi’s new book, and Shahidi’s contribution to this volume. I hope the reader does not mind this informality.

Finally I would like to thank Jianya Liu, the faculty at Shandong University, and all the students that attended for giving me the opportunity for presenting these lectures.

I L -functions for $GL(n)$ and Converse Theorems

1 Modular Forms and Automorphic Representations

1.1 Classical Modular Forms and Their L -functions

We begin our tale by recalling the classical results of Hecke and Weil.

Let $\mathfrak{H} = \{z = x + iy \mid y > 0\}$ denote the complex upper half-plane. Then a (classical) modular form of weight k for $\Gamma = SL_2(\mathbb{Z})$ is a function $f : \mathfrak{H} \rightarrow \mathbb{C}$ which is holomorphic on $\mathfrak{H}$ and at the cusps of Γ and for each $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ we have

$$f(\gamma z) = f\left(\frac{az + b}{cz + d}\right) = (cz + d)^k f(z).$$

We denote the space of these by $M_k(\Gamma)$. These arise in the theory of theta series, elliptic modular forms, etc. Since $f(z + 1) = f(z)$, by modularity under $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, this will have a Fourier expansion

$$f(z) = \sum_{n=0}^{\infty} a_n e^{2\pi i n z}$$

and is called a *cusp form* if

$$a_0 = \int_0^1 f(x + iy) dx = 0.$$

Recall that $\Gamma \backslash \mathfrak{H}$ is the Riemann sphere with the “cusp at ∞ ” removed, so cusp forms are those modular forms that vanish at the cusps. The subspace of cusp forms of weight k is denoted by $S_k(\Gamma)$.

To each cusp form Hecke associated an L -function given by the Dirichlet series

$$L(s, f) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$$

which is absolutely convergent for $\text{Re}(s) > \frac{k}{2} + 1$. The L -function is related to f through a Mellin transform:

$$\Lambda(s, f) = (2\pi)^{-s} \Gamma(s) L(s, f) = \int_0^{\infty} f(iy) y^s d^\times y.$$

By transferring the analytic properties of f to $L(s, f)$, Hecke showed:

Theorem 1.1. *If $f \in S_k(\Gamma)$ then $\Lambda(s, f)$ is nice, i.e.,*

- (1) $\Lambda(s, f)$ extends to an entire function of s ;
- (2) $\Lambda(s, f)$ is bounded in vertical strips (BVS);
- (3) $\Lambda(s, f)$ satisfied the functional equation (FE)

$$\Lambda(s, f) = (i)^k \Lambda(k - s, f).$$

Note that the functional equation results from the modular transformation law of $f(z)$ under $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, namely $f(Sz) = f\left(\frac{-1}{z}\right) = z^k f(z)$.

Remark 1.1. In general $L(s, f)$ need not have an *Euler product*. Hecke introduced an algebra of operators $\mathcal{H} = \langle T_p \mid p \text{ prime} \rangle$, the original Hecke algebra, such that if f is a simultaneous eigen-function for all T_n , i.e., $T_n f = \lambda_n f$, and if we normalize f such that $a_1 = 1$ then $\lambda_n = a_n$ and we have

$$L(s, f) = \prod_p (1 - a_p p^{-s} + p^{2k-1} p^{-2s})^{-1}.$$

Hecke was able to invert this process, via the inverse Mellin transform, and prove the “Hecke Converse Theorem”.

Theorem 1.2. *Let*

$$D(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$$

be a Dirichlet series that converges in some right half-plane. Set

$$\Lambda(s) = (2\pi)^{-s} \Gamma(s) D(s).$$

If $\Lambda(s)$ is nice as above, i.e., (1) entire continuation, (2) BVS, and (3) satisfies the functional equation $\Lambda(s) = (i)^k \Lambda(k - s)$, then

$$f(z) = \sum_{n=1}^{\infty} a_n e^{2\pi i n z}$$

is a cusp form of weight k for $\Gamma = SL_2(\mathbb{Z})$.

Note that by the Fourier expansion $f(z+1) = f(z)$, so $f(z)$ is modular under the translation matrix T . Since $SL_2(\mathbb{Z})$ is generated by T and the inversion S , we only need the transformation law under S . This follows from the functional equation for $\Lambda(s)$ via Mellin inversion

$$f(iy) = \sum_{n=1}^{\infty} a_n e^{-2\pi n y} = \frac{1}{2\pi i} \int_{\text{Re}(s)=2} \Lambda(s) y^{-s} ds.$$

If f is a cusp form not for all of $SL_2(\mathbb{Z})$ but say for the Hecke congruence group of level N

$$\Gamma = \Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}) \mid c \equiv 0 \pmod{N} \right\}$$

then the inversion S is no longer in Γ . Weil still defined $L(s, f)$ for $f \in S_k(\Gamma)$ and proved they were nice, using now $S_N = \begin{pmatrix} 0 & -1 \\ N & 0 \end{pmatrix}$, which normalizes $\Gamma_0(N)$, in place of S . (Now the functional equation relates f and a form g related to f through S_N .) To invert this process, i.e., to establish the “Weil Converse Theorem”, Weil showed that besides knowing that $D(s)$ was nice, one had to also control twisted Dirichlet series; one needed that the

$$D_\chi(s) = \sum_{n=1}^{\infty} \frac{\chi(n)a_n}{n^s}$$

were *nice* (essentially) for all Dirichlet characters χ of conductor relatively prime to N . The important thing to note here is that one Dirichlet series no longer suffices for inversion, one must control a family of twisted Dirichlet series as well.

1.2 Automorphic Representations of $GL(n)$

The modern theory of automorphic representations is a theory of functions on adèle groups. To make the connection note that

$$\mathfrak{H} = PGL_2^+(\mathbb{R})/PSO_2(\mathbb{R}).$$

So functions of $\mathfrak{H}$ can be lifted to functions on $GL_2(\mathbb{R})$. So we now have functions on a group.

To make an adelic theory, take k a number field ... but for now we can take $k = \mathbb{Q}$. The the adele ring of $\mathbb{Q}$ is

$$\mathbb{A} = \mathbb{A}_{\mathbb{Q}} = \varinjlim_S (\mathbb{R} \times \prod_{p \in S} \mathbb{Q}_p \times \prod_{p \notin S} \mathbb{Z}_p) = \prod'_v \mathbb{Q}_v,$$

where the limit is over all finite subsets S of primes of $\mathbb{Z}$. Then $\mathbb{Q} \hookrightarrow \mathbb{A}$ as a discrete, co-compact subgroup with $\mathbb{Q} \cap (\mathbb{R} \times \prod_p \mathbb{Z}_p) = \mathbb{Z}$, so that $\mathbb{Z} \backslash \mathbb{R} \simeq \mathbb{Q} \backslash \mathbb{A} / \prod_p \mathbb{Z}_p$.

Similarly, for $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$,

$$GL_n(\mathbb{A}) = \varinjlim_S (GL_n(\mathbb{R}) \times \prod_{p \in S} GL_n(\mathbb{Q}_p) \times \prod_{p \notin S} GL_n(\mathbb{Z}_p)) = \prod'_v GL_n(\mathbb{Q}_v)$$

and $GL_n(\mathbb{Q}) \hookrightarrow GL_n(\mathbb{A})$ as a canonical discrete subgroup. While the quotient $GL_n(\mathbb{Q}) \backslash GL_n(\mathbb{A})$ is no longer compact, it is finite volume modulo the center $Z_n(\mathbb{A}) \simeq \mathbb{A}^\times$.

The analogue of the space of classical modular forms is the space of automorphic forms

$$\mathcal{A}(GL_n(\mathbb{Q}) \backslash GL_n(\mathbb{A}); \omega)$$

consisting of functions $\varphi : GL_n(\mathbb{A}) \rightarrow \mathbb{C}$ such that $\varphi(\gamma g) = \varphi(g)$ for all $\gamma \in GL_n(\mathbb{Q})$ plus regularity and growth conditions to match the classical conditions of holomorphy and holomorphy at the cusps. Here ω is a fixed unitary central character, i.e., φ satisfies $\varphi(zg) = \omega(z)\varphi(g)$ for $z \in Z_n(\mathbb{A})$. $GL_n(\mathbb{A})$ acts on the