

Norman Steenrod

Princeton Landmarks in Mathematics

The Topology of Fibre Bundles

纤维丛拓扑学

PRINCETON UNIVERSITY PRESS

世界图书出版公司

www.wpcbj.com.cn

THE TOPOLOGY OF FIBRE BUNDLES

By
NORMAN STEENROD

PRINCETON UNIVERSITY PRESS

PRINCETON, NEW JERSEY

图书在版编目 (CIP) 数据

纤维丛拓扑学: 英文/ (美) 斯廷罗德 (Steenrod, N.) 著.
—影印本. —北京: 世界图书出版公司北京公司, 2011. 1
书名原文: The Topology of Fibre Bundles
ISBN 978-7-5100-2956-1

I. ①纤… II. ①斯… III. ①纤维丛—拓扑—英文
IV. ①O189.3

中国版本图书馆 CIP 数据核字 (2010) 第 212307 号

书 名: The Topology of Fibre Bundles

作 者: Norman Steenrod

中 译 名: 纤维丛拓扑学

责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河国英印务有限公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64021602, 010-64015659

电子信箱: kjb@wpcbj.com.cn

开 本: 24 开

印 张: 10

版 次: 2011 年 01 月

版权登记: 图字: 01-2010-4884

书 号: 978-7-5100-2956-1/O · 862

定 价: 35.00 元

THE TOPOLOGY OF FIBRE BUNDLES

PRINCETON LANDMARKS
IN MATHEMATICS AND PHYSICS

Non-standard Analysis, *by Abraham Robinson*

General Theory of Relativity, *by P.A.M. Dirac*

Angular Momentum in Quantum Mechanics,
by A. R. Edmonds

Mathematical Foundations of Quantum Mechanics,
by John von Neumann

Introduction to Mathematical Logic, *by Alonzo Church*

Convex Analysis, *by R. Tyrrell Rockafellar*

Riemannian Geometry, *by Luther Pfahler Eisenhart*

The Classical Groups, *by Hermann Weyl*

Topology from the Differentiable Viewpoint,
by John W. Milnor

Algebraic Theory of Numbers, *by Hermann Weyl*

Continuous Geometry, *by John von Neumann*

Linear Programming and Extensions, *by George B. Dantzig*

Operator Techniques in Atomic Spectroscopy, *by Brian R. Judd*

Preface

The recognition of the domain of mathematics called fibre bundles took place in the period 1935–1940. The first general definitions were given by H. Whitney. His work and that of H. Hopf and E. Stiefel demonstrated the importance of the subject for the applications of topology to differential geometry. Since then, some seventy odd papers dealing with bundles have appeared. The subject has attracted general interest, for it contains some of the finest applications of topology to other fields, and gives promise of many more. It also marks a return of algebraic topology to its origin; and, after many years of introspective development, a revitalization of the subject from its roots in the study of classical manifolds.

No exposition of fibre bundles has appeared. The literature is in a state of partial confusion, due mainly to the experimentation with a variety of definitions of “fibre bundle.” It has not been clear that any one definition would suffice for all results. The derivations of analogous conclusions from differing hypotheses have produced much overlapping. Many “known” results have not been published. It has been realized that certain standard theorems of topology are special cases of propositions about bundles, but the generalized forms have not been given.

The present treatment is an initial attempt at an organization. It grew out of lectures which I gave at the University of Michigan in 1947, and at Princeton University in 1948. The informed reader will find little here that is essentially new. Only such improvements and fresh applications are made as must accompany any reasonably successful organization.

The book is divided into three parts according to the demands made on the reader’s knowledge of topology. The first part presupposes only a minimum of point set theory and closes with two articles dealing with covering spaces and the fundamental group. Part II makes extensive use of the homotopy groups of Hurewicz. Since no treatment of these has appeared in book form, Part II opens with a survey of the subject. Definitions and results are stated in detail; some proofs are given, and others are indicated. In Part III we make use of cohomology theory. Here, again, a survey is required because

the standard treatments do not include the generalized form we must use. A reader who is familiar with the elements of homology theory will have little difficulty.

I must acknowledge my gratitude to Professor Sze-tsen Hu and Dr. R. L. Taylor who read the manuscript and suggested many improvements.

I wish to acknowledge also the aid of the National Academy of Sciences in support of publication of this volume.

Numbers enclosed in brackets refer to the bibliography.

NORMAN STEENROD

May, 1950
Princeton University

Copyright © 1951 by Princeton University Press
Published by Princeton University Press, 41 William Street,
Princeton New Jersey 08540
In the United Kingdom: Princeton University Press,
Chichester, West Sussex

All Rights Reserved

Second printing 1957, appendix added
Sixth printing 1967
Seventh printing, and first paperback printing, 1999

Steenrod, Norman Earl, 1910–1971.
The topology of fibre bundles / by Norman E. Steenrod.
p. cm. — (Princeton landmarks in mathematics
and physics) Originally published: Princeton :
Princeton University Press, 1951.

Includes bibliographical references and index.

ISBN 0-691-00548-6 (pbk. : alk. paper)

1. Fibre bundles (Mathematics) I. Title. II. Series.

QA 612.6.S74 1999

514'.224—dc21

99-17187

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without permission in writing from the Publisher.

<http://pup.princeton.edu>

Printed in the United States of America

1 3 5 7 9 10 8 6 4 2

Contents

Part I. THE GENERAL THEORY OF BUNDLES

1. Introduction	3
2. Coordinate bundles and fibre bundles	6
3. Construction of a bundle from coordinate transformations	14
4. The product bundle	16
5. The Ehresmann-Feldbau definition of bundle.	18
6. Differentiable manifolds and tensor bundles	20
7. Factor spaces of groups	28
8. The principal bundle and the principal map	35
9. Associated bundles and relative bundles	43
10. The induced bundle	47
11. Homotopies of maps of bundles.	49
12. Construction of cross-sections	54
13. Bundles having a totally disconnected group.	59
14. Covering spaces	67

Part II. THE HOMOTOPY THEORY OF BUNDLES

15. Homotopy groups	72
16. The operations of π_1 on π_n	83
17. The homotopy sequence of a bundle	90
18. The classification of bundles over the n -sphere	96
19. Universal bundles and the classification theorem.	100
20. The fibering of spheres by spheres	105
21. The homotopy groups of spheres	110
22. Homotopy groups of the orthogonal groups	114
23. A characteristic map for the bundle R_{n+1} over S^n	118
24. A characteristic map for the bundle U_n over S^{2n-1}	124
25. The homotopy groups of miscellaneous manifolds	131
26. Sphere bundles over spheres.	134
27. The tangent bundle of S^n	140
28. On the non-existence of fiberings of spheres by spheres	144

Part III. THE COHOMOLOGY THEORY OF BUNDLES

29. The stepwise extension of a cross-section	148
30. Bundles of coefficients	151

31. Cohomology groups based on a bundle of coefficients . . .	155
32. The obstruction cocycle	166
33. The difference cochain	169
34. Extension and deformation theorems	174
35. The primary obstruction and the characteristic cohomology class	177
36. The primary difference of two cross-sections	181
37. Extensions of functions, and the homotopy classification of maps.	184
38. The Whitney characteristic classes of a sphere bundle . . .	190
39. The Stiefel characteristic classes of differentiable manifolds .	199
40. Quadratic forms on manifolds	204
41. Complex analytic manifolds and exterior forms of degree 2.	209
 Appendix	 218
 Bibliography	 223
 Index	 228

THE TOPOLOGY OF FIBRE BUNDLES

Part I. The General Theory of Bundles

§1. INTRODUCTION

1.1. Provisional definition. A fibre bundle \mathfrak{B} consists, at least, of the following: (i) a topological space B called the *bundle space* (or, simply, bundle), (ii) a topological space X called the *base space*, (iii) a continuous map

$$p: B \rightarrow X$$

of B onto X called the *projection*, and (iv) a space Y called the *fibre*. The set Y_x , defined by

$$Y_x = p^{-1}(x),$$

is called the *fibre over the point x of X* . It is required that each Y_x be homeomorphic to Y . Finally, for each x of X , there is a neighborhood V of x and a homeomorphism

$$\phi: V \times Y \rightarrow p^{-1}(V)$$

such that

$$p\phi(x', y) = x' \quad x' \in V, y \in Y.$$

A *cross-section* of a bundle is a continuous map $f: X \rightarrow B$ such that $pf(x) = x$ for each $x \in X$.

The above definition of bundle is not sufficiently restrictive. A bundle will be required to carry additional structure involving a group G of homeomorphisms of Y called the *group of the bundle*. Before imposing the additional requirements, consideration of a collection of examples will show the need for these. The discussion of these examples will be brief and intuitive; each will be treated later in detail.

1.2. The product bundle. The first example is the *product bundle* or product space $B = X \times Y$. In this case, the projection is given by $p(x, y) = x$. Taking $V = X$ and $\phi =$ the identity, the last condition is fulfilled. The cross-sections of B are just the graphs of maps $X \rightarrow Y$. The fibres are, of course, all homeomorphic, however there is a natural unique homeomorphism $Y_x \rightarrow Y$ given by $(x, y) \rightarrow y$. As will be seen, this is equivalent to the statement that the group G of the bundle consists of the identity alone.

1.3. The Möbius band. The second example is the Möbius band. The base space X is a circle obtained from a line segment L (as indicated in Fig. 1) by identifying its ends. The fibre Y is a line segment. The

bundle B is obtained from the product $L \times Y$ by matching the two ends with a twist. The projection $L \times Y \rightarrow L$ carries over under this matching into a projection $p: B \rightarrow X$. There are numerous cross-sections; any curve as indicated with end points that match provides a cross-section. It is clear that any two cross-sections must agree on at least one point. There is no natural unique homeomorphism of Y_x




FIG. 1.

with Y . However there are two such which differ by the map g of Y on itself obtained by reflecting in its midpoint. In this case the group G is the cyclic group of order 2 generated by g .

1.4. The Klein bottle. The third example is the Klein bottle. The preceding construction is modified by replacing the fibre by a circle (Fig. 2). The ends of the cylinder $L \times Y$ are identified, as indicated, by reflecting in the diameter de . Again, the group G is the cyclic group




FIG. 2.

of order 2 generated by this reflection. (It is impossible to visualize this example in complete detail since the Klein bottle cannot be imbedded topologically in euclidean 3-space.)

1.5. The twisted torus. The fourth example, we will call the *twisted torus*. The construction is the same as for the Klein bottle except that reflection in the diameter de is replaced by reflection in the center of the circle (or rotation through 180°). As before, the group G is

cyclic of order 2. In contrast to the preceding two examples, this bundle is homeomorphic to the product space $X \times Y$ and in such a way as to preserve fibres. However to achieve this one must use homeomorphisms $Y \rightarrow Y_x$ other than the two natural ones. But they need not differ from these by more than rotations of Y . This behavior is expressed by saying that the twisted torus is not a product bundle, but it is equivalent to one in the full group of rotations of Y .

1.6. Covering spaces. A covering space B of a space X is another example of a bundle. The projection $p: B \rightarrow X$ is the covering map. The usual definition of a covering space is the definition of bundle, in §1.1, modified by requiring that each Y_x is a discrete subspace of B , and that ϕ is a homeomorphism of $V \times Y_x$ with $p^{-1}(V)$ so that $\phi(x, y) = y$. If, in addition, it is supposed that X is arcwise connected, motion of a point x along a curve C in X from x_1 to x_2 can be covered by a continuous motion of Y_x in B from Y_{x_1} to Y_{x_2} . Choosing a base point x_0 , each Y_x can be put in 1-1 correspondence with $Y = Y_{x_0}$, using a curve in X . This correspondence depends only on the homotopy class of the curve. Considering the action on Y of closed curves from x_0 to x_0 , the fundamental group $\pi_1(X)$ appears as a group of permutations on Y . Any two correspondences of Y_x with Y differ by a permutation corresponding to an element of $\pi_1(X)$. Thus, *for covering spaces, the group of the bundle is a factor group of the fundamental group of the base space.*

1.7. Coset spaces. Another example of a bundle is a Lie group B operating as a transitive group of transformations on a manifold X . The projection is defined by selecting a point $x_0 \in X$ and defining $p(b) = b(x_0)$. If Y is the subgroup of B which leaves x_0 fixed, then the fibres are just the left cosets of Y in B . There are many natural correspondences $Y \rightarrow Y_x$, any $b \in Y_x$ defines one by $y \rightarrow b \cdot y$. However any two such $y \rightarrow b \cdot y$, $y \rightarrow b' \cdot y$ differ by the left translation of Y corresponding to $b^{-1}b'$. Thus the group G of the bundle coincides with the fibre Y and acts on Y by left translations. Finding a cross-section for such a bundle is just the problem of constructing in B a simply-transitive continuous family of transformations.

1.8. The tangent bundle of a manifold. As a final example let X be an n -dimensional differentiable manifold, let B be the set of all tangent vectors at all points of X , and let p assign to each vector its initial point. Then Y_x is the tangent plane at x . It is a linear space. Choosing a single representative Y , linear correspondences $Y_x \rightarrow Y$ can be constructed (using chains of coordinate neighborhoods in X), but not uniquely. In this case the group G of the bundle is the full linear group operating on Y . A cross-section here is just a vector field over X . The entire bundle is called the tangent bundle of X .

1.9. Generalizations of product spaces. It is to be observed that all the preceding examples of bundles are very much like product spaces. The language and notation has been designed to reflect this fact. A bundle is a generalization of a product space. The study of two spaces X and Y and maps $f: X \rightarrow Y$ is equivalent to the study of the product space $X \times Y$, its projections into X and Y , and graphs of maps f . This is broadened by replacing $X \times Y$ by a bundle space B , sacrificing the projection into Y , but replacing it, for each x , by a family of maps $Y_x \rightarrow Y$ any two of which differ by an element of a group G operating on Y . The graphs of continuous functions $f: X \rightarrow Y$ are replaced by cross-sections of the bundle.

This point of view would lead one to expect that most of the concepts of topology connected with pairs of spaces and their maps should generalize in some form. This is sustained in all that follows. For example, the Hopf theorem on the classification of maps of an n -complex into an n -sphere generalizes into the theory of the characteristic cohomology classes of a sphere-bundle.

The problems connected with bundles are of various types. The simplest question is the one of existence of a cross-section. This is of importance in differential geometry where a tensor field with prescribed algebraic properties is to be constructed. Is the bundle equivalent to a product bundle? If so, there exist many cross-sections. What are the relations connecting the homology and homotopy groups of the base space, bundle, fibre, and group? Can the bundle be simplified by replacing the group G by a smaller one? For given X, Y, G , what are the possible distinct bundles B ? This last is the classification problem.

§2. COORDINATE BUNDLES AND FIBRE BUNDLES

2.1. The examples of §1 show that a bundle carries, as part of its structure, a group G of transformations of the fibre Y . In the last two examples, the group G has a topology. It is necessary to weave G and its topology into the definition of the bundle. This will be achieved through the intermediate notion of a *fibre bundle with coordinate systems* (briefly: "coordinate bundle"). The coordinate systems are eliminated by a notion of equivalence of coordinate bundles, and a passage to equivalence classes.

2.2. Transformation groups. A *topological group* G is a set which has a group structure and a topology such that (a) g^{-1} is continuous for g in G , and (b) $g_1 g_2$ is continuous simultaneously in g_1 and g_2 , i.e. the map $G \times G \rightarrow G$ given by $(g_1, g_2) \rightarrow g_1 g_2$ is continuous when $G \times G$ has the usual topology of a product space.

If G is a topological group, and Y is a topological space, we say

that G is a topological transformation group of Y relative to a map $\eta: G \times Y \rightarrow Y$ if (i) η is continuous, (ii) $\eta(e, y) = y$ where e is the identity of G , and (iii) $\eta(g_1 g_2, y) = \eta(g_1, \eta(g_2, y))$ for all g_1, g_2 in G and y in Y .

As we shall rarely consider more than one such η , we shall abbreviate $\eta(g, y)$ by $g \cdot y$. Then (ii) becomes $e \cdot y = y$ and (iii) becomes $(g_1 g_2) \cdot y = g_1 \cdot (g_2 \cdot y)$. For any fixed g , $y \rightarrow g \cdot y$ is a homeomorphism of Y onto itself; for it has the continuous inverse $y \rightarrow g^{-1} \cdot y$. In this way η provides a homomorphism of G into the group of homeomorphisms of Y .

We shall say that G is *effective* if $g \cdot y = y$, for all y , implies $g = e$. Then G is isomorphic to a group of homeomorphisms of Y . In this case one might identify G with the group of homeomorphisms, however we shall frequently allow the same G to operate on several spaces.

Unless otherwise stated, a topological transformation group will be assumed to be effective.

2.3. Definition of coordinate bundle. A *coordinate bundle* \mathcal{B} is a collection as follows:

- (1) A space B called the *bundle space*,
- (2) a space X called the *base space*,
- (3) a map $p: B \rightarrow X$ of B onto X called the *projection*,
- (4) a space Y called the *fibre*,
- (5) an effective topological transformation group G of Y called the *group of the bundle*,
- (6) a family $\{V_j\}$ of open sets covering X indexed by a set J , the V_j 's are called *coordinate neighborhoods*, and
- (7) for each j in J , a homeomorphism

$$\phi_j: V_j \times Y \rightarrow p^{-1}(V_j)$$

called the *coordinate function*.

The coordinate functions are required to satisfy the following conditions:

$$(8) \quad p\phi_j(x, y) = x, \quad \text{for } x \in V_j, y \in Y,$$

(9) if the map $\phi_{j,x}: Y \rightarrow p^{-1}(x)$ is defined by setting

$$\phi_{j,x}(y) = \phi_j(x, y),$$

then, for each pair i, j in J , and each $x \in V_i \cap V_j$, the homeomorphism

$$\phi_{j,x}^{-1}\phi_{i,x}: Y \rightarrow Y$$

coincides with the operation of an element of G (it is unique since G is