


Lecture Notes in Mathematics

1586

Rodney Nillsen

Difference Spaces and Invariant Linear Forms



Springer-Verlag

Rodney Nillsen

Difference Spaces and Invariant Linear Forms

Springer-Verlag

Berlin Heidelberg New York

London Paris Tokyo

Hong Kong Barcelona

Budapest

Author

Rodney Nillsen

Department of Mathematics

University of Wollongong

Northfields Avenue

Wollongong NSW 2522, Australia

E-Mail: r.nillsen@uow.edu.au

The figure on the cover illustrates the graphs of the following functions: the first derivative $-2xe^{-x^2}$ of e^{-x^2} ; the first order difference $e^{-(x-1)^2} - e^{-(x+1)^2}$ of $e^{-(x-1)^2}$; the second derivative $2(2x^2 - 1)e^{-x^2}$ of e^{-x^2} ; and the second order difference $e^{-x^2} - 2^{-1}(e^{-(x-1)^2} + e^{-(x+1)^2})$ of e^{-x^2} . The general relationship between such functions is described in Corollary 3.7 of Chapter III.

Mathematics Subject Classification (1991): 43-02, 42-02, 42B10, 42B20, 43A15, 47F05

ISBN 3-540-58323-8 Springer-Verlag Berlin Heidelberg New York

ISBN 0-387-58323-8 Springer-Verlag New York Berlin Heidelberg

CIP-Data applied for

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

© Springer-Verlag Berlin Heidelberg 1994

Printed in Germany

Typesetting: Camera ready by author

46/3140-543210 - Printed on acid-free paper

Editorial Policy

§ 1. Lecture Notes aim to report new developments - quickly, informally, and at a high level. The texts should be reasonably self-contained and rounded off. Thus they may, and often will, present not only results of the author but also related work by other people. Furthermore, the manuscripts should provide sufficient motivation, examples and applications. This clearly distinguishes Lecture Notes manuscripts from journal articles which normally are very concise. Articles intended for a journal but too long to be accepted by most journals, usually do not have this “lecture notes” character. For similar reasons it is unusual for Ph. D. theses to be accepted for the Lecture Notes series.

§ 2. Manuscripts or plans for Lecture Notes volumes should be submitted (preferably in duplicate) either to one of the series editors or to Springer- Verlag, Heidelberg . These proposals are then refereed. A final decision concerning publication can only be made on the basis of the complete manuscript, but a preliminary decision can often be based on partial information: a fairly detailed outline describing the planned contents of each chapter, and an indication of the estimated length, a bibliography, and one or two sample chapters - or a first draft of the manuscript. The editors will try to make the preliminary decision as definite as they can on the basis of the available information.

§ 3. Final manuscripts should preferably be in English. They should contain at least 100 pages of scientific text and should include

- a table of contents;
- an informative introduction, perhaps with some historical remarks: it should be accessible to a reader not particularly familiar with the topic treated;
- a subject index: as a rule this is genuinely helpful for the reader.

Further remarks and relevant addresses at the back of this book.

Editors:

A. Dold, Heidelberg

B. Eckmann, Zürich

F. Takens, Groningen



To Sylvia

“ That is not said right,” said the Caterpillar.
“ Not *quite* right, I’m afraid,” said Alice, timidly;
“ some of the words have got altered.”

LEWIS CARROLL,
in *Alice’s Adventures in Wonderland*

PREFACE

The ideas in this work have had a long period of gestation, although the fuller development and expression of them has occurred over a relatively short and intense period. The dominant ideas emerged in an endeavour to answer the following question: if f is a function in $L^2(\mathbb{R}^n)$, how can the behaviour of its Fourier transform near the origin of \mathbb{R}^n be described and characterized? The corresponding question for the circle group had been given a satisfactory answer by Gary Meisters and Wolfgang Schmidt in 1972, so the work also can be regarded as arising from an attempt to extend their result from the compact case of the circle group to the non-compact case of \mathbb{R}^n . The answers presented to these and related questions have implications for other areas of analysis; the notable ones being the ranges of partial differential operators, and the behaviour of some of the singular integral operators of classical analysis.

I have received a great deal of help and encouragement from many individuals, many of whom are probably not aware of the positive effect they have had upon my carrying out this work. Their input has been too multifarious and diverse for me to record them all by name or to list all of the many ways in which their ideas or suggestions or actions have been of aid to me.

There are, however, some for whom it is incumbent upon me to record their very substantial and specific assistance. One of these was Igor Kluvánek, who sustained an interest in and an encouragement for this work over an extended period – in particular, since late in 1990, when in an extended conversation in a Randwick restaurant, we discussed the general question of the behaviour of the Fourier transform near the origin. As well, to the extent that the present work is considered to emphasise clarity and precision of thought, and is considered to be written in a way which shows respect and consideration for the reader, I would like to think that the writing of it shows his positive influence which extends back over a much longer period. His death in 1994 meant the loss of a close friend who had also been a great intellectual influence upon me. Among my colleagues at Wollongong, I am particularly indebted to Graham Williams and Keith Tognetti. The former has provided me with invaluable opportunities for discussing many of the ideas in this work, which is all the more appreciated because of the time which was freely given, but which often could be ill-afforded. The latter has had an infectious enthusiasm and openness to new ideas, which has also had an effect on my own more sceptical nature, and this has been all to my own good. I am also indebted to Wai Lok Lo, who detected several errors in various earlier versions of the manuscript.

As well, I am aware that my mathematical colleagues at Wollongong have played a very positive, if indirect, rôle in this work, by maintaining an harmonious and intellectually alive environment amongst themselves which has contributed in no small measure to my capacity and opportunity to carry this work through to completion. Also, I have been fortunate in receiving support directly from the Analysis Research Group at my University, and indirectly from the Graduate Faculty. All of these have sustained me during a period in which changing attitudes towards universities, and within them, have

created difficulties of acceptance, understanding and appreciation of those disciplines and researches primarily concerned with the basic understanding of phenomena and abstract concepts.

I have been extremely fortunate with the dedication and attention to detail shown by Carolyn Silveri in what has proved to be a long and demanding task of word-processing. Her many suggestions have greatly added to the presentation of the completed work. I am also indebted to Kerrie Gamble, who originally prepared the draft of Chapter IV.

The work has been completed in the face of many competing tasks, and the tolerance, understanding, encouragement and support of my wife have been immeasurably important. The good humour with which my children have borne both my absences and unavailability for sufficiently many pastimes has also played its part in my completing this work. I am also very grateful to my parents, who have invariably provided me with encouragement in my work over many years.

Notwithstanding the help I have received from so many people, for any errors, omissions or deficiencies in the work, the author should be considered responsible.

Rodney Nillsen,

August, 1994

CONTENTS

INTRODUCTION

1. Difference spaces	1
2. Differentiation, differences and the behaviour of the Fourier transform near the origin	2
3. Multiplication spaces	4
4. Arrangement of the work	4
5. Notation and conventions	5
6. Notes	7

CHAPTER I. GENERAL AND PREPARATORY RESULTS

1. Introduction: further aspects of the multiplication spaces and difference spaces	9
2. The spaces of abstract distributions	10
3. Characterization of the multiplication and difference spaces	12
4. Multiplication sets and difference sets	20
5. Difference spaces and invariant linear forms	24
6. Measures with prescribed Fourier transform	35
7. Notes	42

CHAPTER II. MULTIPLICATION AND DIFFERENCE SPACES ON \mathbb{R}^n

1. Introduction : the growth function and its significance	44
2. The finite integral theorem	45
3. The characterization of multiplication sets	50
4. The Fourier transform interpretation: the characterization of difference sets	67
5. Some particular spaces, interpolation and applications to invariant linear forms	77
6. Sharpness aspects of the characterizations of the difference and multiplication spaces	86
7. Spaces of negative index and dual spaces	98
8. Difference spaces, centroids and moments	102
9. Connections with wavelets	110
10. Notes	114

CHAPTER III. APPLICATIONS TO DIFFERENTIAL AND SINGULAR INTEGRAL OPERATORS

1. Introduction : operators and difference spaces	118
2. Sobolev-type spaces and multiplier operators	119
3. Applications to differential operators	125
4. Applications to singular integral operators	137
5. Notes	150

CHAPTER IV. RESULTS FOR L^p SPACES ON GENERAL GROUPS

1. Introduction: the averaging method of Woodward	152
2. Difference spaces and invariant linear forms for $L^1(G)$	153
3. Difference spaces and invariant linear forms for $L^p(G)$, $1 < p \leq \infty$	162
4. Notes	173

BIBLIOGRAPHY	175
--------------	-----

INDEX OF AUTHORS	178
------------------	-----

INDEX OF SUBJECTS	179
-------------------	-----

INDEX OF SYMBOLS	182
------------------	-----

INTRODUCTION

1. Difference spaces

Let G be a Hausdorff locally compact group and let $M(G)$ denote the space of Radon measures on G . Let $*$ denote the usual convolution of functions, measures and distributions on G . Let $1 \leq p < \infty$ and let X be a subspace of $L^p(G)$ or a space of distribution on G with the property that $M(G) * X \subseteq X$. If $x \in G$ let $\delta_x \in M(G)$ denote the Dirac measure at x . Then if $\emptyset \neq S \subseteq M(G)$, the vector space $\mathcal{D}(X, S)$ is defined to be the subspace of X consisting of all functions or distributions f in X for which there exist $m \in \mathbb{N}$, $\mu_1, \mu_2, \dots, \mu_m \in S$ and $f_1, f_2, \dots, f_m \in X$ such that $f = \sum_{j=1}^m (f_j - \mu_j * f_j)$. The space $\mathcal{D}(X, S)$ is called a *difference space* of X . In the case when $S = \{\delta_x : x \in G\}$, the difference space $\mathcal{D}(X, S)$ is denoted by $\mathcal{D}(X)$.

A linear form L in the algebraic dual X' of X is said to be S -invariant if $L(\mu * f) = L(f)$ for all $f \in X$ and $\mu \in S$. Equivalently, L is S -invariant if and only if L vanishes identically on the subspace $\mathcal{D}(X, S)$ of X . If $x \in G$ and $f \in X$, $\delta_x * f$ is called the (left) *translation* of f by x . Note that if $f \in L^p(G)$, $(\delta_x * f)(t) = f(x^{-1}t)$, for almost all $t \in G$. A linear form L on X is said to be *translation invariant* if $L(\delta_x * f) = L(f)$ for all $x \in G$ and $f \in X$. Thus, L is translation invariant if and only if L vanishes identically on $\mathcal{D}(X)$. In fact, it is easy to prove, using a Hamel basis argument, that

$$\mathcal{D}(X, S) = \bigcap \left\{ \text{kernel of } L : L \in X' \text{ and } L \text{ is } S\text{-invariant} \right\}. \quad (1.1)$$

The preceding remarks make it clear that there is a close relationship between invariant linear forms on X and the corresponding difference space of X , and this is a main reason for interest in the difference spaces. Historically, interest seems to have centred more on the invariant forms associated with difference spaces, rather than with the difference spaces themselves. This is perhaps because of the importance of invariant integration in many different areas – for example, the Haar measure on G is the measure arising from a positive linear form which vanishes identically on a difference space of continuous functions which have compact support.

In particular, there is a large body of work on invariant linear forms on $L^\infty(G)$. If μ is a positive, continuous element of $L^\infty(G)'$ and $\|\mu\| = 1$, then μ is called a *mean* on $L^\infty(G)$, and G is said to be *amenable* if $L^\infty(G)$ has a translation invariant mean. The theory of amenability on groups and other structures is now very extensive, and is the subject of recent books by Paterson [49] and Pier [50].

The emphasis in this work is more upon invariant linear forms on $L^p(G)$ for $1 \leq p < \infty$, and upon invariant forms on spaces $\mathcal{F}_p(G)$ of abstract distributions on G which are characterized by the property that their Fourier transforms are in $L^p(\hat{G})$, where \hat{G} is the dual of G when G is abelian. Even more, the emphasis is upon the corresponding difference spaces and upon closely related spaces known as generalized difference spaces. In certain cases where G is abelian, the main results characterize, for example, a difference space $\mathcal{D}(L^2(G), S)$ as being isomorphic under the Fourier transform to a space $L^2(\hat{G}, \mu)$,

where μ is a non-negative measure on \hat{G} . Such characterizations of difference spaces suggest that these spaces are of interest quite apart from their connection with invariant linear forms. A further reason for interest in these spaces, and the generalized difference spaces, is that they also serve to characterize the ranges of certain types of differential operators, including such familiar ones as the Laplace and wave operators. Also, they shed light on the behaviour of some of the singular integral operators of classical analysis such as the Riesz potential operators.

2. Differentiation, differences and the behaviour of the Fourier transform near the origin

In numerical analysis, finite differences have long been used to give local approximations to derivatives. For example, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable and if its derivative $D(f) \in L^2(\mathbb{R})$, consider $x, y \in \mathbb{R}$ with y near the origin, and let $g \in L^2(\mathbb{R})$ be the function given by $g(u) = -f(u)/y$. Then $D(f)(x)$ may be approximated by

$$\frac{f(x+y) - f(x)}{y} = \frac{(\delta_{-y} * f - f)(x)}{y} = (g - \delta_{-y} * g)(x).$$

This suggests that, even globally, there may be a relationship between $D(f)$ and differences of the form $h - \delta_{-y} * h$, for $h \in L^2(\mathbb{R})$ and $y \in \mathbb{R}$.

A major motivation for the n -dimensional results presented here was to develop a theory of difference spaces, or a theory of spaces related to such spaces, which would suffice to characterize the ranges of a reasonably large class of partial differential operators. Each of these operators was envisaged as being defined on a suitable Sobolev-type subspace of $L^2(\mathbb{R}^n)$ (say), and the range of each one was envisaged as being a difference space or some closely related space. The connection between these spaces and differential operators comes via the Fourier transform and, because of the centrality of this idea to this work, it is now described in more detail. The Fourier transform of f is denoted by \hat{f} or f^\wedge .

If $f \in L^2(\mathbb{R})$ let $D(f)$ denote the derivative of f in the sense of Schwartz distributions. Consider $f \in L^2(\mathbb{R})$ such that $D(f) \in L^2(\mathbb{R})$. Then $\hat{f} \in L^2(\mathbb{R})$ and $D(f)^\wedge(x) = ix\hat{f}(x)$, for almost all $x \in \mathbb{R}$. Plancherel's theorem then shows that $D(f)^\wedge$ is small near the origin in the sense that

$$\int_{-\infty}^{\infty} \frac{|D(f)^\wedge(x)|^2}{|x|^2} dx < \infty.$$

In fact, it is easy to see that

$$D(L^2(\mathbb{R})) \cap L^2(\mathbb{R}) = \left\{ g : g \in L^2(\mathbb{R}) \text{ and } \int_{-\infty}^{\infty} \frac{|\hat{g}(x)|^2}{|x|^2} dx < \infty \right\}. \quad (2.1)$$

On the other hand, if $f \in L^2(\mathbb{R})$ and $y \in \mathbb{R}$, $(f - \delta_y * f)^\wedge(x) = (1 - e^{-ixy})\hat{f}(x)$,

for almost all $x \in \mathbb{R}$, so that

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{|(f - \delta_y * f)^\wedge(x)|^2}{|x|^2} dx &= \int_{-\infty}^{\infty} |\hat{f}(x)|^2 \left| \frac{1 - e^{-ixy}}{x} \right|^2 dx, \\ &= y^2 \int_{-\infty}^{\infty} \left(\frac{\sin \frac{xy}{2}}{\frac{xy}{2}} \right)^2 |\hat{f}(x)|^2 dx, \\ &\leq y^2 \int_{-\infty}^{\infty} |\hat{f}(x)|^2 dx, \text{ as } |\sin x| \leq |x|, \\ &< \infty. \end{aligned}$$

It follows that for all $h \in \mathcal{D}(L^2(\mathbb{R}))$,

$$\int_{-\infty}^{\infty} \frac{|\hat{h}(x)|^2}{|x|^2} dx < \infty,$$

an observation originally made in 1973 by Meisters, who deduced from it that $\mathcal{D}(L^2(\mathbb{R}))$ is a proper, dense subspace of $L^2(\mathbb{R})$ and that there are discontinuous linear forms on $L^2(\mathbb{R})$ which are translation invariant. More recently, the author has proved that, in fact,

$$\mathcal{D}(L^2(\mathbb{R})) = \left\{ h : h \in L^2(\mathbb{R}) \text{ and } \int_{-\infty}^{\infty} \frac{|\hat{h}(x)|^2}{|x|^2} dx < \infty \right\}. \quad (2.2)$$

This shows that the functions in $L^2(\mathbb{R})$ which are in $\mathcal{D}(L^2(\mathbb{R}))$ are completely characterized by the behaviour of their Fourier transforms near the origin. Equation (2.2) also shows that $\mathcal{D}(L^2(\mathbb{R}))$ is a Hilbert space in the norm $||| \cdot |||$ given by $|||f||| = \left(\int_{-\infty}^{\infty} |\hat{f}(x)|^2 (1 + |x|^{-2}) dx \right)^{1/2}$.

A comparison of (2.1) and (2.2) shows that $\mathcal{D}(L^2(\mathbb{R})) = D(L^2(\mathbb{R})) \cap L^2(\mathbb{R})$, a result which describes $\mathcal{D}(L^2(\mathbb{R}))$ as the range of the differentiation operator D on the Sobolev space consisting of those functions in $L^2(\mathbb{R})$ whose derivatives are in $L^2(\mathbb{R})$. This Sobolev space is a Hilbert space in the norm $|| \cdot ||$ which is given by $||f|| = \left(\int_{-\infty}^{\infty} |\hat{f}(x)|^2 (1 + |x|^2) dx \right)^{1/2}$, and the differentiation operator is an isometry from this space onto the space $\mathcal{D}(L^2(\mathbb{R}))$ in the norm $||| \cdot |||$. The theory of generalized multiplication spaces and generalized difference spaces presented in Chapter II was motivated by an attempt to extend this description of the range of the differentiation operator, and its description as an isometry, to more general differential operators. Such results may be regarded as putting the idea of approximating derivatives by finite differences within the framework of functional analysis and operator theory.

The preceding discussion also establishes a connection between the range of the differentiation operator D and invariant linear forms. For, since the range of D , when D is restricted to the appropriate Sobolev subspace of $L^2(\mathbb{R})$, is equal to $\mathcal{D}(L^2(\mathbb{R}))$, it follows from (1.1) that

$$D(L^2(\mathbb{R})) \cap L^2(\mathbb{R}) = \bigcap \left\{ \text{kernel of } L : L \in L^2(\mathbb{R})' \text{ and } L \text{ is translation invariant} \right\}.$$

A similar description is also valid for the ranges of differential operators belonging to a class of operators which includes the Laplace and wave operators. To the extent that one regards amenability as the study of invariant linear forms and associated concepts, such results may be considered to establish a connection between amenability and the theory of partial differential operators.

3. Multiplication spaces

Let G be a Hausdorff locally compact abelian group with dual \hat{G} . Let F be a family of complex valued Borel measurable functions on G . Then if $1 \leq p \leq \infty$, the vector space $\mathcal{M}(L^p(G), F)$ is defined to consist of those functions f defined almost everywhere on G for which there exist $m \in \mathbb{N}$, $f_1, f_2, \dots, f_m \in F$ and $g_1, g_2, \dots, g_m \in L^p(G)$ such that $f = \sum_{j=1}^m f_j g_j$. The space $\mathcal{M}(L^p(G), F)$ is called a *multiplication space* of $L^p(G)$. The Fourier transform changes convolutions into multiplications, so it follows that if $S \subseteq M(G)$, if $F = \{1 - \hat{\mu} : \mu \in S\}$ and if $1 \leq p < \infty$, then the Fourier transform is a bijection from $\mathcal{D}(\mathcal{F}_p(G), S)$ onto $\mathcal{M}(L^p(\hat{G}), F)$.

Corresponding to the generalized difference spaces are the generalized multiplication spaces. The basic problem is to characterize a multiplication space, or a generalized multiplication space of $L^p(G)$, as a space $L^p(G, \nu)$ for some measure ν on G .

Characterizing a multiplication space of $L^p(G)$ in this way is usually easier than the corresponding problem for difference spaces of $L^p(G)$. This is because characterizing a space $\mathcal{M}(L^p(G), F)$ is usually no harder than characterizing $\mathcal{M}(L^2(G), F)$, but the Fourier transform is not generally a bijection from $L^p(G)$ onto a space $L^q(\hat{G})$, a fact familiar from considerations arising from the Hausdorff-Young theorem (see [29, p.146], for example). This difficulty is the motivation for introducing the spaces $\mathcal{F}_p(G)$ of abstract distributions on G , and for considering difference spaces of $\mathcal{F}_p(G)$.

4. Arrangement of the work

Chapter I is largely concerned with multiplication spaces and sets, with difference spaces and sets, and with the characterization of these in terms of the finiteness of certain integrals. The relationship between difference spaces and invariant linear forms is discussed, the emphasis being on how the two types of object are related to each other. The relationship between the multiplication and difference spaces is given by the Fourier transform, and this raises the problem of constructing measures having a certain type of prescribed Fourier transform. This is discussed for subsequent application.

In Chapter II are presented the basic characterizations of generalized multiplication and difference spaces on \mathbb{R}^n . This leads to deeper and more useful characterizations of these sets than was possible in the more general setting in Chapter I. Some sharpness aspects of these characterizations are presented, duality aspects are discussed, and the chapter concludes with discussions of other representations of some of the difference spaces, and with establishing a connection between these spaces and the theory of

wavelets.

Chapter III is concerned with applications to describing the ranges of various types of partial differential operators in terms of the corresponding difference spaces. In this formulation the operators are isometries from an appropriate Sobolev-type space onto a corresponding generalized difference space. Applications to various singular integral operators are discussed, including the Hilbert and Riesz transforms and the Riesz potential operators.

Difference spaces of $L^p(G)$, where G is a locally compact group (not necessarily abelian) and $1 \leq p \leq \infty$, are discussed in Chapter IV. Owing to the difficulty associated with the Fourier transform in this more general setting, the results concentrate on comparison of difference spaces with each other, rather than with explicit characterizations as were possible for \mathbb{R}^n . These results may be interpreted equally as results about various types of invariant linear forms on $L^p(G)$.

Detailed references and comments are kept for the notes section at the end of each chapter, except where they are considered essential for elucidating the main text.

The main aim of the work has been to present new results on difference spaces, multiplication spaces, invariant linear forms and associated applications. However, every effort has been made to make the whole work, and each chapter, as self contained as seemed reasonable. It is hoped that this will enhance readability of the work. A reader familiar with integration theory, Fourier theory and the theory of distributions on \mathbb{R}^n , should find Chapters II and III quite accessible. A reader familiar with basic harmonic analysis on locally compact groups should find Chapters I and IV readily accessible. It is hoped that the work as a whole will be of interest to graduate students and researchers in the areas of Fourier analysis, abstract harmonic analysis, partial differential equations and singular integral operators.

5. Notations and conventions

In this section are introduced some of the notations and conventions to be used. The set of natural numbers is denoted by \mathbb{N} , the integers by \mathbb{Z} , the real numbers by \mathbb{R} , the complex numbers by \mathbb{C} , and the circle group of complex numbers of modulus one by \mathbb{T} . The non-negative integers are denoted by \mathbb{Z}_+ , and the non-negative reals by \mathbb{R}_+ . The complement of a set A is denoted by A^c . If A_1, A_2, \dots, A_n are sets, their Cartesian product is denoted by $\prod_{j=1}^n A_j$, and this is abbreviated to A^n in the case that $A_1 = A_2 = \dots = A_n = A$. The cardinality of the continuum is denoted by c . If f is a given function, it is occasionally convenient to denote it alternatively by $f(\cdot)$. The complex conjugate of a function f is denoted by \bar{f} .

If G is a Hausdorff locally compact group, μ_G denotes a non-zero left invariant Haar measure on G . If G is also abelian, \hat{G} denotes the dual group of G . The identity of G is generally denoted by e , and the identity of \hat{G} (when G is abelian) is generally denoted by \hat{e} . If G is abelian, the Fourier transform on $L^1(G)$, relative to some choice of a non-zero Haar measure μ_G , is given by $\hat{f}(\gamma) = \int \bar{\gamma} f d\mu_G$, for $f \in L^1(G)$ and $\gamma \in \hat{G}$. In general, the Fourier transform of a function, distribution or measure μ on G is denoted by $\hat{\mu}$ or

μ^\wedge . The inverse Fourier transform of a function, distribution or measure μ is denoted by μ^\vee .

If $n \in \mathbb{N}$, \mathbb{R}^n is the corresponding n -dimensional Euclidean space. If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ and $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, the inner product of x and y is $\langle x, y \rangle = \sum_{j=1}^n x_j y_j$. Also, $|x|$ is the Euclidean norm of x , $|x| = \langle x, x \rangle^{1/2}$. The usual Lebesgue measure on \mathbb{R}^n is denoted by μ_n (thus $\mu_n = "dx"$), and the Fourier transform on \mathbb{R}^n is taken relative to μ_n . Thus, if $f \in L^1(\mathbb{R}^n)$, \hat{f} is given by $\hat{f}(x) = \int_{\mathbb{R}^n} e^{-i\langle \cdot, x \rangle} f d\mu_n$, for all $x \in \mathbb{R}^n$.

Plancherel's Theorem says that the Fourier transform may be extended from $L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ to all of $L^2(\mathbb{R}^n)$ so that it becomes a linear bijection on $L^2(\mathbb{R}^n)$ such that $\int_{\mathbb{R}^n} f \bar{g} d\mu_n = (2\pi)^{-n} \int_{\mathbb{R}^n} \hat{f} \overline{\hat{g}} d\mu_n$ for all $f, g \in L^2(\mathbb{R}^n)$.

If $\mu_1, \mu_2, \dots, \mu_n$ are measures, their product is denoted by $\prod_{j=1}^n \mu_j$, abbreviated to μ^n in the case $\mu_1 = \mu_2 = \dots = \mu_n = \mu$. Also, if a_1, a_2, \dots, a_n are numbers, $\prod_{j=1}^n a_j$ denotes their product. The symbol \prod may therefore stand for any one of several different products of sets, measures or numbers.

The space of infinitely differentiable complex valued functions on \mathbb{R}^n is denoted by $C^\infty(\mathbb{R}^n)$, and a function $f \in C^\infty(\mathbb{R}^n)$ is said to be *rapidly decreasing* if $\varphi \frac{\partial}{\partial x_{i_1}} \frac{\partial}{\partial x_{i_2}} \dots \frac{\partial}{\partial x_{i_\ell}}(f) \in L^\infty(\mathbb{R}^n)$ for all polynomials φ and all $i_1, i_2, \dots, i_\ell \in \mathbb{N}$. The space of rapidly decreasing functions is denoted by $\mathcal{S}(\mathbb{R}^n)$. If $m, i_1, i_2, \dots, i_\ell \in \mathbb{Z}_+$, the mapping

$$f \mapsto \left\| (1 + |x|^2)^m \frac{\partial}{\partial x_{i_1}} \frac{\partial}{\partial x_{i_2}} \dots \frac{\partial}{\partial x_{i_\ell}}(f) \right\|_\infty$$

defines a seminorm on $\mathcal{S}(\mathbb{R}^n)$. There is a countable number of such seminorms, and $\mathcal{S}(\mathbb{R}^n)$ is given the coarsest topology in which all such seminorms are continuous. This makes $\mathcal{S}(\mathbb{R}^n)$ a Fréchet space whose continuous dual $\mathcal{S}(\mathbb{R}^n)^*$ is the space of *tempered distributions* on \mathbb{R}^n . If $1 \leq p \leq \infty$ and $f \in L^p(\mathbb{R}^n)$, f can be identified with the element of $\mathcal{S}(\mathbb{R}^n)^*$ given by $\varphi \mapsto \int_{\mathbb{R}^n} f \varphi d\mu_n$, for $\varphi \in \mathcal{S}(\mathbb{R}^n)$. Thus each space $L^p(\mathbb{R}^n)$, for $1 \leq p \leq \infty$, can be regarded as a set of tempered distributions.

The Fourier transform is a linear homeomorphism on $\mathcal{S}(\mathbb{R}^n)$. If $f \in \mathcal{S}(\mathbb{R}^n)^*$, the Fourier transform \hat{f} of f is in $\mathcal{S}(\mathbb{R}^n)^*$ and is given by the equation

$$\hat{f}(\varphi) = f(\hat{\varphi}), \text{ for } \varphi \in \mathcal{S}(\mathbb{R}^n).$$

In the case where $f \in L^1(\mathbb{R}^n)$, the previous definition of \hat{f} is consistent with the present definition of \hat{f} when f is regarded as an element of $\mathcal{S}(\mathbb{R}^n)^*$.

Further notations will be introduced as required. The book by Hewitt and Ross [23] is a suitable reference for harmonic analysis on groups, and its notation is closely followed here. The books by Rudin [54] and Schwartz [58] are very useful references for the theory of distributions and Fourier analysis on \mathbb{R}^n . The book of Rudin is followed more closely here as far as notation is concerned.

A reference in the text to Theorem 2.3, for example, refers to Theorem 2.3 in the same chapter as the reference. Otherwise, the chapter is explicitly mentioned. The end of a proof or example is indicated by \square .