


HYDRODYNAMIC STABILITY

P. G. DRAZIN

Professor of Applied Mathematics, University of Bristol

W. H. REID

Professor of Applied Mathematics, University of Chicago



CAMBRIDGE UNIVERSITY PRESS

CAMBRIDGE

LONDON NEW YORK NEW ROCHELLE

MELBOURNE SYDNEY

Published by the Press Syndicate of the University of Cambridge
The Pitt Building, Trumpington Street, Cambridge CB2 1RP
32 East 57th Street, New York, NY 10022, USA
296 Beaconsfield Parade, Middle Park, Melbourne 3206, Australia

© Cambridge University Press 1981

First published 1981
First paperback edition 1982

Reprinted 1984

Printed in the United States of America
British Library cataloguing in publication data

Drazin, P G
Hydrodynamic stability.—

(Cambridge monographs on mechanics and applied mathematics).

1. Hydrodynamics 2. Stability

I. Title II. Reid, William Hill III. Series
532'.51 QC151 80-40273

ISBN 0 521 22798 4 hard covers
ISBN 0 521 28980 7 paperback

TO OUR PARENTS

Isaac and Leah Drazin

William and Edna Reid

PREFACE

For nearly a century now, hydrodynamic stability has been recognized as one of the central problems of fluid mechanics. It is concerned with when and how laminar flows break down, their subsequent development, and their eventual transition to turbulence. It has many applications in engineering, in meteorology and oceanography, and in astrophysics and geophysics. Some of these applications are mentioned, but the book is written from the point of view intrinsic to fluid mechanics and applied mathematics. Thus, although we have emphasized the analytical aspects of the theory, we have also tried, wherever possible, to relate the theory to experimental and numerical results.

Our aim in writing this book has been twofold. Firstly, in Chapters 1–4, to describe the fundamental ideas, methods, and results in three major areas of the subject: thermal convection, rotating and curved flows, and parallel shear flows. Secondly, to provide an introduction to some aspects of the subject which are of current research interest. These include some of the more recent developments in the asymptotic theory of the Orr–Sommerfeld equation in Chapter 5, some applications of the linear stability theory in Chapter 6 and finally, in Chapter 7, a discussion of some of the fundamental ideas involved in current work on the nonlinear theory of hydrodynamic stability.

Each chapter ends with a number of problems which often extend or supplement the main text as well as provide exercises to help the reader understand the topics. An asterisk is used to indicate those problems which we judge to be relatively long or difficult. Some hints and references are given to help in the solution of many of the problems. We have also prepared answers to the problems which may be obtained from either author for a nominal charge to cover reproduction and postage costs.

Thus this is a textbook suitable for a graduate course on the fundamental ideas and methods and on the major applications of the theory of hydrodynamic stability. It also leads the reader up to the frontiers of research on selected topics. In general we have assumed that the reader is familiar with whatever mathematical methods are needed, notably in the theories of ordinary and of partial differential equations and in the theory of functions of a complex variable. But we have explained some specialized and modern mathematical points at length where it seems that they are likely to be unfamiliar to most readers.

We are grateful to our many colleagues throughout the world who have responded so generously to our various inquiries. In particular, we thank A. Davey, T. H. Hughes, and L. M. Mack for providing new or unpublished numerical results, R. J. Donnelly, E. L. Koschmieder and S. A. Thorpe for providing photographs, J. P. Cleave for advice on some mathematical points, L. C. Woods for advice on the presentation of the material, B. S. Ng for detailed comments on Chapters 1-5, and A. Davey and J. T. Stuart for constructive criticisms of a draft of Chapter 7. For help with the typing of the manuscript we also thank N. Thorp in Bristol and M. Bowie, F. Flowers, L. Henley, and M. Newman in Chicago. We are especially indebted to S. Chandrasekhar and C. C. Lin, who have contributed so much to the theory of hydrodynamic stability; through their papers and books, and through our personal contacts with them, they have greatly influenced our work on the subject. One of us (W.H.R.) also wishes to acknowledge with thanks the generous support provided over the years by the U.S. National Science Foundation, most recently under grant no. MCS 78-01249.

And, finally, we should like to thank G. K. Batchelor not only for his help as editor of this series but also for his kindness during an early stage in our careers when it was our good fortune to be associated with him.

Bristol

P.G.D.

Chicago

W.H.R.

August 1979

For this new impression we have corrected some misprints and other errors, and we have inserted Problem 1.12 on p. 31 and an addendum on pp. 479-80.

March 1982

P.G.D.

W.H.R.

CONTENTS

<i>Preface</i>	page xiii
----------------	--------------

1 INTRODUCTION

1 Introduction	1
2 Mechanisms of instability	4
3 Fundamental concepts of hydrodynamic stability	8
4 Kelvin-Helmholtz instability	14
5 Break-up of a liquid jet in air	22
Problems for chapter 1	27

2 THERMAL INSTABILITY

6 Introduction	32
7 The equations of motion	34
The exact equations, 34; The Boussinesq equations, 35	
8 The stability problem	37
The linearized equations, 37; The boundary conditions, 40; Normal modes, 42	
9 General stability characteristics	44
Exchange of stabilities, 44; A variational principle, 45	
10 Particular stability characteristics	50
Free-free boundaries, 50; Rigid-rigid boundaries, 51; free-rigid boundaries, 52	
11 The cells	52
12 Experimental results	59
13 Some applications	62
Problems for chapter 2	63

3 CENTRIFUGAL INSTABILITY

14	Introduction	69
15	Instability of an inviscid fluid	71
	Three-dimensional disturbances, 73; Axisymmetric disturbances, 77; Two-dimensional disturbances, 80	
16	Instability of Couette flow of an inviscid fluid	82
17	The Taylor problem	88
	Axisymmetric disturbances, 90; Two-dimensional disturbances, 103; Three-dimensional disturbances, 104; Some experimental results, 104	
18	The Dean problem	108
	The Dean problem, 108; The Taylor-Dean problem, 113	
19	The Görtler problem	116
	Problems for chapter 3	121

4 PARALLEL SHEAR FLOWS

20	Introduction	124
	The inviscid theory	
21	The governing equations	126
22	General criteria for instability	131
23	Flows with piecewise-linear velocity profiles	144
	Unbounded vortex sheet, 145; Unbounded shear layer, 146; Bounded shear layer, 147	
24	The initial-value problem	147
	The viscous theory	
25	The governing equations	153
26	The eigenvalue spectrum for small Reynolds numbers	158
	A perturbation expansion, 159; Sufficient conditions for stability, 161	
27	Heuristic methods of approximation	164
	The reduced equation and the inviscid approximations, 165; The boundary-layer approximation near a rigid wall, 167; The WKBJ approximations, 167; The local turning-point approximations,	

	171; The truncated equation and Tollmien's improved viscous approximations, 175; The viscous correction to the singular inviscid solution, 177	
28	Approximations to the eigenvalue relation	180
	Symmetrical flows in a channel, 181; Flows of the boundary-layer type, 183; The boundary-layer approximation to $\phi_3(z)$, 184; The WKBJ approximation to $\phi_3(z)$, 185; The local turning-point approximation to $\phi_3(z)$, 188; Tollmien's improved approximation to $\phi_3(z)$, 191	
29	The long-wave approximation for unbounded flows	196
30	Numerical methods of solution	202
	Expansions in orthogonal functions, 203; Finite-difference methods, 206; Initial-value methods (shooting), 207	
31	Stability characteristics of various basic flows	211
	Plane Couette flow, 212; Poiseuille flow in a circular pipe, 216; Plane Poiseuille flow, 221; Combined plane Couette and plane Poiseuille flow, 223; The Blasius boundary-layer profile, 224; The asymptotic suction boundary-layer profile, 227; Boundary layers at separation, 229; The Falkner-Skan profiles, 231; The Bickley jet, 233; The hyperbolic-tangent shear layer, 237	
32	Experimental results	239
	Problems for chapter 4	245

5 UNIFORM ASYMPTOTIC APPROXIMATIONS

33	Introduction	251
	Plane Couette flow	
34	The integral representations of the solutions	256
35	The differential equation method	263
	General velocity profiles	
36	A preliminary transformation	265

37	The inner and outer expansions	267
	The inner expansions, 268; The outer expansions, 271; The central matching problem, 276; Composite approximations, 278	
38	Uniform approximations	280
	The solution of well-balanced type, 280; The solutions of balanced type, 280; The solutions of dominant-recessive type, 283	
39	A comparison with Lin's theory	285
40	Preliminary simplification of the eigenvalue relation	290
41	The uniform approximation to the eigenvalue relation	295
	A computational form of the first approximation to the eigenvalue relation, 299; Results for plane Poiseuille flow, 301	
42	A comparison with the heuristic approximations to the eigenvalue relation	305
	The local turning-point approximation to $\phi_3(z)$, 305; Tollmien's improved approximation to $\phi_3(z)$, 306; The uniform approximation to $\phi_3(z)$ based on the truncated equation, 308; The uniform approximation to $\phi_3(z)$ based on the Orr-Sommerfeld equation, 309	
43	A numerical treatment of the Orr-Sommerfeld problem using compound matrices	311
	Symmetrical flows in a channel, 315; Boundary-layer flows, 316	
	Problems for chapter 5	317

6 ADDITIONAL TOPICS IN LINEAR STABILITY THEORY

44	Instability of parallel flow of a stratified fluid	320
	Introduction, 320; Internal gravity waves and Rayleigh-Taylor instability, 324; Kelvin-Helmholtz instability, 325	
45	Baroclinic instability	333
46	Instability of the pinch	339
47	Development of linear instability in time and space	345
	Initial-value problems, 345; Spatially growing modes, 349	

48	Instability of unsteady flows	353
	Introduction, 353; Instability of periodic flows, 354;	
	Instability of other unsteady basic flows, 361	
	Problems for chapter 6	363

7 NONLINEAR STABILITY

49	Introduction	370
	Landau's theory, 370; Discussion, 376	
50	The derivation of ordinary differential systems governing stability	380
51	Resonant wave interactions	387
	Internal resonance of a double pendulum, 387;	
	Resonant wave interactions, 392	
52	Fundamental concepts of nonlinear stability	398
	Introduction to ordinary differential equations, 398;	
	Introduction to bifurcation theory, 402; Structural stability, 407; Spatial development of nonlinear stability, 416; Critical layers in parallel flow, 420	
53	Additional fundamental concepts of nonlinear stability	423
	The energy method, 424; Maximum and minimum energy in vortex motion, 432; Application of boundary-layer theory to cellular instability, 434	
54	Some applications of the nonlinear theory	435
	Bénard convection, 435; Couette flow, 442;	
	Parallel shear flows, 450	
	Problems for chapter 7	458

APPENDIX. A CLASS OF GENERALIZED AIRY FUNCTIONS

A1	The Airy functions $A_k(z)$	465
A2	The functions $A_k(z, p)$, $B_0(z, p)$ and $B_k(z, p)$	466
A3	The functions $A_k(z, p, q)$ and $B_k(z, p, q)$	472
A4	The zeros of $A_1(z, p)$	477
	Addendum: Weakly non-parallel theories for the Blasius boundary layer	479
	<i>Bibliography and author index</i>	481
	<i>Motion picture index</i>	517
	<i>Subject index</i>	519

CHAPTER 1

INTRODUCTION

Yet not every solution of the equations of motion, even if it is exact, can actually occur in Nature. The flows that occur in Nature must not only obey the equations of fluid dynamics, but also be stable.

— L. D. Landau & E. M. Lifshitz (1959)

1 Introduction

The essential problems of hydrodynamic stability were recognized and formulated in the nineteenth century, notably by Helmholtz, Kelvin, Rayleigh and Reynolds. It is difficult to introduce these problems more clearly than in Osborne Reynolds's (1883) own description of his classic series of experiments on the instability of flow in a pipe.

The ... experiments were made on three tubes ... The diameters of these were nearly 1 inch, $\frac{1}{2}$ inch and $\frac{1}{4}$ inch. They were all ... fitted with trumpet mouthpieces, so that the water might enter without disturbance. The water was drawn through the tubes out of a large glass tank, in which the tubes were immersed, arrangements being made so that a streak or streaks of highly coloured water entered the tubes with the clear water.

The general results were as follows:—

(1) When the velocities were sufficiently low, the streak of colour extended in a beautiful straight line through the tube, Fig. 1.1(a).

(2) If the water in the tank had not quite settled to rest, at sufficiently low velocities, the streak would shift about the tube, but there was no appearance of sinuosity.

(3) As the velocity was increased by small stages, at some point in the tube, always at a considerable distance from the trumpet or intake, the colour band would all at once mix up with the surrounding water, and fill the rest of the tube with a mass of coloured water, as in Fig. 1.1(b). Any increase in the velocity caused the point of break down to approach the trumpet, but with no velocities that were tried did it reach this. On viewing the tube by the light of an electric spark, the mass of colour resolved itself into a mass of more or less distinct curls, showing eddies, as in Fig. 1.1(c).

Reynolds went on to show that the *laminar flow*, the smooth flow he described in paragraph (1), breaks down when Va/ν exceeds a certain critical value, V being the maximum velocity of the water in the tube, a the radius of the tube, and ν the kinematic viscosity of




Fig. 1.1. (a) Laminar flow in a pipe. (b) Transition to turbulent flow in a pipe. (c) Transition to turbulent flow as seen when illuminated by a spark. (From Reynolds 1883.)

water at the appropriate temperature. This dimensionless number Va/ν , now called the *Reynolds number*, specifies any class of dynamically similar flows through a pipe; here we shall denote the number by R . The series of experiments gave the critical value R_c of the Reynolds number as nearly 13 000. However,

the critical velocity was very sensitive to disturbance in the water before entering the tubes This at once suggested the idea that the condition might be one of instability for disturbance of a certain magnitude and stable for smaller disturbances.

At the critical velocity

another phenomenon . . . was the intermittent character of the disturbance. The disturbance would suddenly come on through a certain length of tube and pass away and then come on again, giving the appearance of flashes, and these flashes would often commence successively at one point in the pipe. The appearance when the flashes succeeded each other rapidly was as shown . . .

in Fig. 1.2. Such 'flashes' are now called *turbulent spots* or *turbulent bursts*. Below the critical Reynolds number there was laminar Poiseuille flow with a parabolic velocity profile, the resistance of the




Fig. 1.2. Turbulent spots in a pipe. (From Reynolds 1883.)

pipe to the flow of water being proportional to the mean velocity. As the velocity increased above its critical value, Reynolds found that the flow became *turbulent*, with a chaotic motion that strongly diffused the dye throughout the water in the tube. The resistance of the pipe to turbulent flow grew in proportion to the square of the mean velocity.

Later experimentalists have introduced disturbances of finite amplitude at the intake or used tubes with roughened walls to find R_c as low as 2000, and have used such regular flows and such smooth-walled tubes that R_c was 40 000 or even more.

Reynolds's description illustrates the aims of the study of hydrodynamic stability: to find whether a given laminar flow is unstable and, if so, to find how it breaks down into turbulence or some other laminar flow.

Methods of analysing the stability of flows were formulated in Reynolds's time. The method of normal modes for studying the oscillations and instability of a dynamical system of particles and rigid bodies was already highly developed. A known solution of Newton's or Lagrange's equations of motion for the system was perturbed. The equations were linearized by neglecting products of the perturbations. It was further assumed that the perturbation of each quantity could be resolved into independent components or modes varying with time t like e^{st} for some constant s , which is in general complex. The values of s for the modes were calculated from the linearized equations. If the real part of s was found to be positive for any mode, the system was deemed unstable because a general initial small perturbation of the system would grow exponentially in time until it was no longer small. Stokes, Kelvin and Rayleigh adapted this method of normal modes to fluid dynamics. The essential mathematical difference between fluid and particle dynamics is that the equations of motion are partial rather than ordinary differential equations. This difference leads to many

technical difficulties in hydrodynamic stability, which, to this day, have been overcome for only a few classes of flows with very simple configurations.

Indeed, Reynolds's experiment itself is still imperfectly understood. However, we can explain qualitatively the transition from laminar flow to turbulence with some confidence. Poiseuille flow with a parabolic profile is stable to infinitesimal disturbances at all Reynolds numbers. Some way below the observed critical Reynolds number a finite disturbance may grow if it is not too small. Above the critical Reynolds number quite small disturbances, perhaps introduced at the inlet or by an irregularity of the wall of the tube, grow rapidly with a sinuous motion. Soon they grow so much that nonlinearity becomes strong and large eddies (Fig. 1.1(c)) or turbulent spots (Fig. 1.2) form. At high Reynolds numbers turbulence ensues at once and the flow becomes random and strongly nonlinear everywhere.[†] This instability of Poiseuille flow may be contrasted with that of plane Poiseuille flow, which is unstable to infinitesimal disturbances at sufficiently large values of the Reynolds number. This explanation is supported by the treatment of the theory of the linear stability of Poiseuille flow in § 31 and of the nonlinear stability in § 49.1.

The physical mechanisms of Reynolds's experiments on instability of Poiseuille flow in a pipe are vividly illustrated by a film loop made by Stewart (FL 1968) for the Educational Development Center. This loop consists of edited excerpts from his longer film on *Turbulence* (Stewart F 1968). Details of these and other motion pictures on hydrodynamic stability may be found at the end of the bibliography.

2 Mechanisms of instability

Few laminar flows correspond to known solutions of the nonlinear equations of motion. Fewer are simple enough to allow detailed analysis of their instability. Consequently research on hydrodynamic stability has been deep but narrow. We are forced to study

[†] Many of the features of the transition from laminar to turbulent flow can easily be seen by observing the smoke from a cigarette. Light the cigarette, point the burning tip upwards in still air, and observe the smoke as it rises from rest at the hot tip.

the stability of a few classes of simple laminar flows, mostly with planar, axial or spherical symmetry. Unfortunately, their simplicity obscures some general aspects of instability, especially three-dimensional ones, such as flow with stretching of vortex lines. To gain a wider understanding of hydrodynamic stability and to put these simple flows in perspective as prototypes, it is helpful to sketch the important physical mechanisms of instability.

Broadly speaking, one may say that instability occurs because there is some disturbance of the equilibrium of the external forces, inertia and viscous stresses of a fluid. We shall discuss the external forces first. External forces of interest are buoyancy in a fluid of variable density, surface tension, magnetohydrodynamic forces, etc. It is also convenient to regard centrifugal and Coriolis forces as external forces when there is rotation of the whole system in which the fluid moves. If heavy fluid rests above light fluid it is clear that the fluids tend to overturn under the action of gravity. A similar instability occurs on the free surface of a container of liquid when it is moved downwards at a uniform acceleration greater than the gravitational acceleration. There is in fact a close analogy between the problem of instability of a fluid of variable density, namely Bénard convection bounded by horizontal planes, and the problem of instability of axisymmetric swirling flow of homogeneous fluid, namely Taylor vortices bounded by two coaxial rotating cylinders. The analogue of the density turns out to be the square of the circulation or swirl. If in Couette flow the circulation around the inner cylinder is greater in magnitude than that around the outer, the centrifugal force tends to throw out the fluid near the inner cylinder as an overturning instability. This centrifugal instability may occur also in flows along a rigid curved surface such as a concave wall of a channel. Surface tension resists the increase of area of a surface and so exerts a stabilizing influence, particularly on disturbances of small length scale. A magnetic field can inhibit the motion of an electrically conducting fluid across the magnetic lines of force and thereby usually tends to stabilize flows.

In the absence of any external force or of viscosity, a fluid moves according to the equilibrium between its inertia and internal stresses of pressure. A small disturbance may upset this equilibrium. The tendency of fluid to move down pressure gradients can

be seen to amplify disturbances of certain flows and thereby create instability. This instability can be depicted more reliably in terms of interactions of the vortex lines, which are convected and stretched by the motion of the fluid.

An obvious effect of viscosity is to dissipate the energy of any disturbance and thereby stabilize a flow. Indeed, for this reason any bounded flow is stable if the viscosity is large enough. So, by and large, viscosity has a stabilizing influence. Viscosity has also the more complicated effect of diffusing momentum. This can make some flows, notably parallel shear flows, *unstable* although the same flows of an inviscid fluid are stable.

Thermal conductivity, or molecular diffusion of heat, has some effects similar to those of viscosity, or molecular diffusion of momentum. It tends to smooth out the temperature differences of a disturbance and is usually a stabilizing influence.

It is natural to consider the stability of primarily steady flows, but unsteady ones are also of some practical importance. The acceleration of a laminar flow plays an identifiable role in its stability. Analysis is difficult in general, but it emerges that acceleration of a laminar flow has a stabilizing and deceleration a destabilizing tendency. Flows that oscillate in time, such as Poiseuille flow driven through a circular pipe by an oscillatory pressure gradient, have intricate stability characteristics. Parametric stability or instability may occur, whereby the free oscillations of disturbances of the mean flow resonate with the forced oscillations of the flow.

Finally, the boundaries of a flow are an important factor. They constrain the development of a disturbance and usually the closer they are together the more stable the flow. However, they sometimes give rise to strong shear in boundary layers which is diffused outwards by viscosity and so leads to instability of the flow.

In a typical flow more than one of these mechanisms may act. For example, in plane Poiseuille flow the dual effects of viscosity, the inertia and the boundaries all influence the instability. Plane Poiseuille flow of inviscid fluid is stable. At large but finite Reynolds numbers the diffusion of momentum from thin shear layers near the walls leads to instability. At small Reynolds numbers the dissipative role of viscosity is dominant and there is stability. This leads to a critical value of the dimensionless number R representing the ratio

of the magnitudes of the destabilizing forces of shear and stabilizing viscous forces for which their effects may be said to balance.

This summary of mechanisms of hydrodynamic stability will be given substance by detailed problems in this and later chapters. But the necessary details of the instability of any particular flow should not obscure these general mechanisms, whose recognition helps to classify as well as to understand problems. Instability arising from an upset of the equilibrium between an external force and dissipative effects is usually simpler than inertial instability. Prototype problems of this simpler instability will be analysed first, the linear instability of a fluid heated from below in Chapter 2 and centrifugal instability in Chapter 3. If fluid lies at rest between two horizontal plates, the lower one being hotter than the upper, then we have light fluid below heavy fluid. The buoyancy tends to overturn the fluid. This tendency is countered by the dissipative and diffusive effects of viscosity and thermal conductivity. The dimensionless number typifying the ratio of the destabilizing buoyancy to the stabilizing diffusive forces is called the *Rayleigh number*; its critical value is calculated and related to experiments. In this flow, viscosity and thermal conductivity have only stabilizing effects. The instability of Couette flow between rotating cylinders is analogous, viscosity tending only to stabilize the centrifugal instability. The linear instability of parallel shear flows is treated in Chapter 4, where it is shown that viscosity typically plays the dual roles of stabilizer and destabilizer. There is an imbalance between the inertia and both the dissipative and diffusive effects of viscosity. The physical mechanism of this instability and its mathematical description is more difficult to understand than that of instability due to an external force, and will be explained in detail in Chapter 4. The most difficult mathematical topics, involving the asymptotic theory of the solutions of the Orr-Sommerfeld equation which governs the problem, will be elaborated in Chapter 5. Chapter 5 may be ignored by the less mathematically inclined reader who is interested in stability characteristics rather than in deducing them. More complicated flows whose stability is governed by more than two of the mechanisms will be presented in Chapter 6. Their mathematical difficulties are so formidable that only a few illustrative examples will be given. Also a few other problems will be discussed, in