
UNIVERSAL & ALGEBRA SYMPOSIUM

Proceedings of the V Universal Algebra Symposium.



Turawa, Poland

3-7 May, 1988

Edited by

Katarzyna Hałkowska Bogusław Stawski

Institute of Mathematics Pedagogical University of Opole Poland

Published by

World Scientific Publishing Co. Pte. Ltd. P O Box 128, Farrer Road, Singapore 9128

USA office: World Scientific Publishing Co., Inc. 687 Hartwell Street, Teaneck, NJ 07666, USA

UK office: World Scientific Publishing Co. Pte. Ltd.73 Lynton Mead, Totteridge, London N20 8DH, England

Library of Congress Cataloging-in-Publication data is available.

Proceedings of the V Universal Algebra Symposium UNIVERSAL AND APPLIED ALGEBRA

Copyright © 1989 by World Scientific Publishing Co. Pte. Ltd.

All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the Publisher.

ISBN 9971-50-837-0


Printed in Singapore by JBW Printers & Binders Pte. Ltd.

ORGANIZING COMMITTEE

- K. Hałkowska (chairman); W. Chromik; B Kujawa;
- Z. Majcher; J Płonka; C. Sitarczyk; B. Stawski;
 A. Walendziak; T. Wesołowski.

EDITORIAL BOARD

M. Kolibiar; J. Płonka; J. Słomiński; T. Traczyk;
J. Varlet



此为试读,需要完整PDF请访问: www.ertongbook.com

FOREWORD

Every year since 1984 the conference of Universal and Applied Algebra has held Poland. It been in has Departament of Universal organized cyclically by the Applied Algebra of the Institute of Mathematics of Nicholas Copernicus University in Torun. the Departament in the Institute of Mathematics in Algebra Pedagogical University of Opole and the Departament of Algebraic Methodology of the Istitute of Mathematics the Warsaw Technical University.

The material contained in these Proceedings represents rather faithfully the scope of the V Symposium. What is not mirrored here is the interactive and friendly atmosphere of the Symposium.

We wish to thank all the lecturers and all the participants for creating this atmosphere. We would like to extend special thanks to Professor Stanisław Kochman - the Rector of the Pedagogical University of Opole and Professor Witold Wacławek vice -Rector of the University of Opole for their constant help in the organization of the Conference.

Finally, this book could not have been produced without the active co-operation of the members of the Editorial Board particulary Professor Milan Kolibiar for his contribution to having this volume printed.

Katarzyna Hałkowska Bogusław Stawski

CONTENTS

Foreword	vii
Termal Algebras—an Approach to Inference on Incomplete Data W. Bartol	1.
Completion and Cantor Extension of Cyclically Ordered Groups S. Cemák	13
Boolean Clones and Hyperidentities in Universal Algebras K. Denecke	23
Perfect Distributive Median Algebras H. Draškovičová and J. Hedlíková	46
Medial N-groups and Skew Elements W. A. Dudek	55
On Some Lattices Adequate for Intuitionistic Predicate Logic W. Dzik	81
On the Stability of Marriages T. Fofanova	87
The Tensor Product on a Variety J. Galanová	92
Implicative Orthoposets R. Godowski, T Traczyk and J. Varlet	99
On Normal and Regular Identities and Hyperidentities E. Graczyńska	107
Absolute Independence in Atomless Algebras J. Grygiel	136
On Free Algebras in Varieties Defined by Externally Compatible Identities K. Halkowska	143

On Certain Classes of Categories and Monoids Constructed from	
Abstract Mal'cev Clones, I	149
HJ. Hoehnke	
Categorical Characterization of Universally Axiomatizable Classes	
of Finitary Relational Systems	17
G. Jarzembski	
On a Class of Median Groups	190
M. Kolibiar	
Generalized Pascal Triangles	198
I. Korec	
Determining Congruences by Some System of Their Classes	219
B. Kujawa and K. D. Thanh	-1
Minimal Generics of Regular Varieties J. Płonka	227
On the Structure of Some Bilattices A. Romanowska and A. Trakul	235
On Some Varieties of Symmetric Idempotent Entropic Groupoids B. Roszkowska	254
The Cone and Forgetful Cone Varieties and the Connections Between Hypervarieties and Usual Varieties J. Stomiński	275
On Some Realization of Grassmann Algebra B. Stawski	296
Nilpotency in Universal Algebras R. Šulka	308
Infinite θ -Decompositions in Modular Lattices A. Walendziak	321
On the Join of Some Varieties of Algebras having the Semilattice Structure T. Wesołowski	334

	X
The Sum of an (F_1, F_2) System of Disjoint Algebras A. Wojtunik	346
Enriched Categories and General Algebra H. J. Vogel	353
The Lattice of Varieties of Left Normal Bandoids E. Zając	366
The Sequent Axiomatization of Some Three-Valued Propositional Logic A. Zbrzezny	378
List of Participants	380

391

List of Addresses

TERMAL ALGEBRAS - AN APPROACH TO INFERENCE ON INCOMPLETE DATA

Wiktor Bartol¹
Institute of Mathematics
University of Warsaw
00-901 Warsaw, Poland

Abstract. The notion of a termal (partial) algebra (introduced informally by I.Fleischer) is defined in an intent to create a general algebraic model systems of inference (or action)over incomplete data. We introduce the notion of a consistent and strongly consistent termal algebra and we prove that inconsistency is a local property of finite character i.e. a termal algebra is inconsistent iff it contains an inconsistent finite relative subalgebra. We also t.he t.hat. poset of all consistent congruences of a termal algebra (i.e.of those congruences which yield consistent quotients) is an algebraic lattice.

In [F11] and [F12] I.Fleischer considered informally defined partial algebras with operations induced by terms rather than by fundamental operation symbols. Such algebras (called termal algebras here) could be viewed on a very general level as algebraic models of systems of inference (or actions) on incomplete data. In this interpretation the fundamental operations correspond to primary deduction rules (or elementary actions), while term operations

¹This research has been supported by the Polish Government Grant CPBP 01.01

correspond to secondary deduction rules (or complex actions), which can yield a result even when not all the intermediate steps are defined. For instance, the result of an application of if α then t_1 else t_2 could be defined as t_1 for α true, even when t_2 is undefined at a given valuation of variables. In this approach one may ask whether such an extension of rules is consistent i.e.whether there is some world of reference in which all values (including the "default" ones) can be interpreted. In an abstract setting total (termal) algebras seem to be good candidates for such worlds of reference. The existence of an embedding of a termal algebra into a total one could be understood as possibility of completing the partial knowledge represented by the termal algebra or, putting it in other words, as the existence of a complete context in which this knowledge is valid. This leads to the definition of a consistent termal algebra, which is the main subject of this paper.

Basic Definitions

Let (F,n) be an algebraic type i.e. F is a set of fundamental operation symbols and $n:F\longrightarrow N$ is the arity function into the set N of all non-negative integers (for convenience we consider finitary operations only). Let $T_F(X)$ be the set of all F-terms over a fixed countably infinite set X of variables and denote by T_F the quotient of $T_F(X)$ obtained via an identification of terms which differ only by a permutation on the set X. This means that we identify two terms $t_1, t_2 \in T_F(X)$ with $t_1 = t_1(x_1, ..., x_n)$ iff there is a bijection $n: X \longrightarrow X$ such that t_2 is obtained from t_1 by substituting $n(x_i)$ for x_i for i=1,...,n. Thus any $t\in T_F$ represents an "abstract" construction of a term, independent of the names of variables. We shall use objects in T_F as if they were just terms.

此为试读,需要完整PDF请访问: www.ertongbook.com

Note that the arity function n of a type (F,n) induces a new function $n^*:T_r \longrightarrow N$ such that for any $t \in T_r$, $n^*(t)=|var(t)|$ (the number of different variables in t) is the arity of the term t. We may now consider a new type (T_F, n^*) , which in turn induces a new set of terms T_T (X) (which shall be abbreviated to $T_T(X)$) Clearly any term $\tau \in T_{T_{-}}^{+}(X)$ has a unique decomposition into a term $t \in T_{F}^{-}(X)$, which will be denoted by $r^{\#}$. Hence e.g. if F consists of a single unary symbol f, then T_F can be identified with the set $(f^n(x): n \in N)$, where $f^o(x)=x$, $f^{n+1}(x)=f(f^n(x))$ for $n \in N$. Further.

 $T_T(X) = (f^k(f^l(\ldots(f^m(x))\ldots):k,l,\ldots,m\in\mathbb{N},\ x\in X))$ and, for instance,

 $f^{4}(x)^{\#} = f^{2}(f^{2}(x))^{\#} = f^{3}(f(x))^{\#} = f(f(f(f(x)))) \neq f^{3}(f(y))$ (if $x \neq y$)

We are now ready to define the basic notion: given a type (F,n), a partial algebra $\mathfrak{V}=(A,(\mathbf{t}^{\mathfrak{V}})_{t\in T_{\mathbf{F}}})$ of $(T_{\mathbf{r}}, n^*)$ is a termal algebra iff all equations of the form

hold weakly in $\mathfrak U$ for all $\tau, o \in T_T(X)$ with $\tau^\# = \sigma^\#$. Recall that for any $t \in T_F$, $t^{\mathfrak U}$ denotes a partial $n^\#$ -ary function on A (an interpretation of the operation symbol t in the algebra U) and then for any $\tau \in T_T(X)$, τ^U will be the induced term function on A. An equation $\tau \approx \sigma$ holds weakly in a partial algebra u iff $r^{u}(a) = \sigma^{u}(a)$ for any valuation a of variables, which belongs to the domain of both 7 and

The following example illustrates the notion of a termal algebra:

Ex.1 Let F=(f) with n(f)=1. Then a termal algebra of type (T_F, n^*) is a partial algebra $\mathfrak{V}=(A, (f^{n\mathfrak{V}})_{n\in\mathbb{N}})$, where for each $n \in \mathbb{N}$, f^{NU} is a unary partial operation on U and all the equations of the form

 $f^{k}(f^{l}(...(f^{k}(x))...))=f^{p}(f^{l}(...(f^{s}(x))...))$ with k+l+...+m=p+r+...+s (k,l,...,m,p,r,...s \in N) and $x\in X$ hold weakly in U.

In particular, let A=(a,b,c,d) be a four-element set and let $\mathfrak{A}=(A,(f^{n\mathfrak{A}})_{n\in\mathbb{N}})$ be a unary partial algebra such that:

dom
$$f^{\mathfrak{A}} = \text{dom } f^{\mathfrak{A}\mathfrak{A}} = \text{dom } f^{\mathfrak{B}\mathfrak{A}} = \{a\}$$

dom $f^{\mathfrak{A}\mathfrak{A}} = \emptyset$ for $n \neq 1,3,5$
 $f^{\mathfrak{A}}(a) = b$, $f^{\mathfrak{A}\mathfrak{A}}(a) = c$, $f^{\mathfrak{B}\mathfrak{A}}(a) = d$

Consider now any equation of the form (1). Then it can be easily verified that for any valuation \underline{a} of variables in A either both sides are identical or at least one of them is undefined. Thus all equations of the form (1) hold weakly in $\mathfrak A$ and therefore $\mathfrak A$ is a termal algebra.

A natural class of termal algebras is obtained in the following way: take any partial algebra' $\mathbb{U}=(A,(f^{\mathfrak{U}})_{f\in F})$ of type (F,n). For any $t\in T_F$ define $t^{\mathfrak{U}}$ as the term function induced in the usual way by the fundamental operations of \mathfrak{U} . Then we get a termal algebra $(A,(t^{\mathfrak{U}})_{t\in T_F})$, which shall be called the canonical termal algebra over \mathfrak{U} . Observe that a total termal algebra (of type (T_F,n^*)) is always a canonical termal algebra over its F-reduct. We shall usually omit the distinction between these two algebras.

Basic algebraic notions carry over from "ordinary" partial algebras to termal algebras in a natural way. We only recall some of them for the sake of completeness. A mapping $h: A \longrightarrow B$ is a homomorphism of a termal algebra $\mathfrak{V}=(A,(t^{\mathfrak{V}})_{t\in T_F})$ into a similar termal algebra $\mathfrak{B}=(B,(t^{\mathfrak{B}})_{t\in T_F})$ iff for any $t\in T_F$ and $a\in \text{dom } t^{\mathfrak{V}}$, $h\cdot a\in \text{dom } t^{\mathfrak{B}}$ and then $h(t^{\mathfrak{V}}(a))=t^{\mathfrak{B}}(h\cdot a)$. If moreover $h\cdot a\in \text{dom } t^{\mathfrak{B}}$ implies $a\in \text{dom } t^{\mathfrak{V}}$, then h is a closed homomorphism. An injective homomorphism will be called an embedding.

For any termal algebras $\mathbb{U}=(A,(t^{\mathbb{U}})_{t\in T_F})$ and $\mathbb{B}=(B,(t^{\mathbb{B}})_{t\in T_F})$ with $A\subseteq B$, \mathbb{U} is a weak subalgebra of \mathbb{B} iff $t^{\mathbb{U}}\subseteq t^{\mathbb{B}}$ for all $t\in T_F$. \mathbb{U} is a relative subalgebra of \mathbb{B} iff $t^{\mathbb{U}}=t^{\mathbb{B}}\cap A^{n}(t)+1$ for all $t\in T_F$. Finally, \mathbb{U} is a (closed)

subalgebra of B iff $t^{\mathfrak{A}} = t^{\mathfrak{B}} \cap (A^{n^{\bullet}(t)} \times B)$ for all $t \in T_{F}$.

If U is a weak subalgebra of B, A=B and B is total then we shall call B an inner completion of U.

Algebraic constructions used in this paper such as direct product, direct limit, homomorphic image etc.ar generalized similarly (for precise definitions se e.g.[Burl).


Consistency

Let U be a termal algebra. U is said to be consistent iff it can be embedded into a total termal algebra. It is strongly consistent iff it has an inner completion (to a total termal algebra). U is inconsistent iff it is not consistent. Hence, any canonical termal algebra is both consistent and strongly consistent. Any strongly consistent termal algebra is consistent. The following example shows that the converse may not be true.

 $\underline{\text{Ex}}$ 2 Let $A=\{a,b\}$ and let f be a unary operation symbol. Define a termal algebra $\mathfrak{V}=(A,(f^n\mathfrak{V})_{f\in F})$ so that $\text{dom } f^{\mathfrak{V}}=\{a\}$ and $\text{dom } f^{\mathfrak{S}}=\{b\}$

m
$$f^{\mathcal{U}} = \{a\}$$
 and dom $f^{\mathcal{U}} = \{l\}$
dom $f^{n\mathcal{U}} = \emptyset$ for $n \neq l, 2$
 $f^{\mathcal{U}}(a) = b$ and $f^{\mathcal{U}}(b) = a$


As in the previous example it can be easily seen that $\mathfrak U$ is a termal algebra indeed. It is consistent, since it can be embedded into a total algebra as in the following diagram(the dotted lines represent the completion of $f^{\mathfrak U}$):



On the other hand, there is no completion on the set $\{\varepsilon,b\}$, which would extend all the operations f^{nV} for $n \in \mathbb{N}$. Indeed,

if $G=((a,b),f^G)$ were a total algebra completing U, then for any possible value of $f^G(b)$ we would have $f^{2G}(b)=b\neq f^{2U}(b)$. Hence U is not strongly consistent.

Ex.3 Let U be the algebra of Ex.1. It is consistent, since it can be embedded into a total algebra as indicated below:



However, if we modify U into U only by identifying b and c:



then ${\bf U}'$ becomes inconsistent, since for any total algebra ${\bf C}$ which extends ${\bf U}'$ we would have both

$$f^{2a}(c) = c$$

(since $f^{20}(c) = f^{20}(f^{0}(a)) = f^{30}(a) = f^{30}(a) = c$) and

(since $f^{26}(c) = f^{26}(f^{36}(a) = f^{56}(a) = f^{54}(a) = d$).

Observe that the mapping $h:(a,b,c,d) \longrightarrow (a,-d)$ such that h(b)=c and h(x)=x for $x\neq b$ is a closed homomorphism of the consistent termal algebra $\mathfrak U$ onto its inconsistent modification $\mathfrak U$. This proves that the class of all consistent termal algebras of a fixed type is not closed with respect to homomorphic images (even if only closed homomorphisms are considered). However, the following holds true:

Theorem 1 The class of all consistent termal algebras

of a fixed type is closed with respect to

- (i) isomorphic images
- (ii) weak subalgebras
- (iii) direct products.

Proof.(i) and (ii): Let $\mathfrak U$ be a consistent termal algebra and h an embedding of $\mathfrak U$ into a total algebra $\mathfrak B$. If j is an isomorphism (or an identical embedding, respectively) of $\mathfrak C$ into $\mathfrak U$, then $h\cdot j$ is an embedding of $\mathfrak C$ into $\mathfrak B$.

(iii): Let $\mathfrak{U}_{i'}, i \in I$, be a family of consistent termal algebras. Then any family of embeddings h_i of \mathfrak{U}_i into a total algebra \mathfrak{B}_i for $i \in I$ induces an embedding of the direct product $\Pi \mathfrak{U}_i$ into the total direct product $\Pi \mathfrak{B}_i$.

Observe that the proofs of (i) and (iii) in the above theorem carry over to the case of strongly consistent termal algebras. Thus

 $\underline{\text{Theorem}}$ $\underline{2}$ The class of all strongly consistent termal algebras of a fixed type is closed with respect to

- (i) isomorphic images
- (ii) direct products.

A weak subalgebra (nor even a closed one) of a strongly consistent termal algebra need not be strongly consistent (though it will always be consistent):


Ex.4 Let $\mathfrak U$ be the algebra of Ex.2. If $\mathfrak B=(\{a,b,c\},f^{\mathfrak B})$ differs from $\mathfrak U$ only in having one additional (isolated) element c in the universe, then $\mathfrak U$ is a subalgebra of $\mathfrak B$. Moreover, $\mathfrak B$ is strongly consistent, while $\mathfrak U$ is not.

The Compactness Theorem

We shall prove here that the consistency of a termal algebra depends on its finite fragments or - equivalently - that a termal algebra is inconsistent iff it contains an inconsistent finite substructure.

Let X be a class of similar partial algebras and let ${\mathfrak A}$

be a partial algebra of the same type as all the algebras in K. A partial algebra $\mathfrak{B} \in K$ is a K-reflection of \mathfrak{U} iff there exists a homomorphism $h: \mathfrak{U} \longrightarrow \mathfrak{B}$ such that for any algebra $\mathfrak{C} \in K$ and any homomorphism $g: \mathfrak{U} \longrightarrow \mathfrak{C}$ there exists a homomorphism (unique) $j: \mathfrak{B} \longrightarrow \mathfrak{C}$ such that the following diagram is commutative:



An algebra $\mathfrak{B} \in \mathcal{K}$ is a \mathcal{K} -monoreflection of \mathfrak{U} iff the homomorphism h in the above definition can be chosen to be injective.

A subclass M of K is (mono)reflective in K iff every algebra U in K has an M-(mono)reflection.

The following theorem gives sufficient conditions for M to be reflective in any class of partial algebras which contains it:

Theorem 3 ([Sch]) Let ${\tt U}$ be an arbitrary partial algebra and let ${\tt M}$ be a class of partial algebras of the same type as ${\tt U}$ such that ${\tt M}$ is closed with respect to

- (i) isomorphic images
- (ii) (closed) subalgebras
- (iii) direct products.

Then U has an M-reflection.

Now let 7 be the class of all termal algebras of some fixed type and let 67 be the subclass of all consistent termal algebras, 77 - the class of all total termal algebras of the same type. Then from Theorems 1 and 3 we deduce the following

Corollary The classes 67 and 77 are both reflective in the class 7. $\hfill\Box$

Moreover, a simple argument leads to the following property of consistent termal algebras:

此为试读,需要完整PDF请访问: www.ertongbook.com