

Max Karoubi

K-Theory

An Introduction

Max Karoubi

K-Theory

An Introduction

With 26 Figures



Springer-Verlag
Berlin Heidelberg New York 1978

Max Karoubi

**Université Paris VII, U.E.R. de Mathématiques, Tour 45-55,
5^e Etage, 2 Place Jussieu
F-75221 Paris Cedex 05**

AMS Subject Classification (1970)

Primary: 55B15, 18F25, 55F45, 55F10, 55D20

**Secondary: 10C05, 16A54, 46H25, 55E20, 55E50, 55E10, 55J25,
55G25**

**ISBN 3-540-08090-2 Springer-Verlag Berlin Heidelberg New York
ISBN 0-387-08090-2 Springer-Verlag New York Heidelberg Berlin**

Library of Congress Cataloging in Publication Data. Karoubi, Max. K-theory. (Grundlehren der mathematischen Wissenschaften; 226). Bibliography: p. Includes index. I. K-theory. I. Title. II. Series: Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen; 226. QA612.33.K373. 514'.23. 77-23162.

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks. Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher.

© by Springer-Verlag Berlin Heidelberg 1978.
Printed in Germany.

Typesetting: William Clowes & Sons Limited, London, Beccles and Colchester. Printing and bookbinding: K. Triltsch, Würzburg.
2141/3140-543210

Foreword

K-theory was introduced by A. Grothendieck in his formulation of the Riemann–Roch theorem (cf. Borel and Serre [2]). For each projective algebraic variety, Grothendieck constructed a group from the category of coherent algebraic sheaves, and showed that it had many nice properties. Atiyah and Hirzebruch [3] considered a topological analog defined for any compact space X , a group $K(X)$ constructed from the category of vector bundles on X . It is this “topological *K*-theory” that this book will study.

Topological *K*-theory has become an important tool in topology. Using *K*-theory, Adams and Atiyah were able to give a simple proof that the only spheres which can be provided with H -space structures are S^1 , S^3 and S^7 . Moreover, it is possible to derive a substantial part of stable homotopy theory from *K*-theory (cf. J. F. Adams [2]). Further applications to analysis and algebra are found in the work of Atiyah–Singer [2], Bass [1], Quillen [1], and others. A key factor in these applications is Bott periodicity (Bott [2]).

The purpose of this book is to provide advanced students and mathematicians in other fields with the fundamental material in this subject. In addition, several applications of the type described above are included. In general we have tried to make this book self-contained, beginning with elementary concepts wherever possible; however, we assume that the reader is familiar with the basic definitions of homotopy theory: homotopy classes of maps and homotopy groups (cf. collection of spaces including projective spaces, flag bundles, and Grassmannians. Hilton [1] or Hu [1] for instance). Ordinary cohomology theory is used, but not until the end of Chapter V. Thus this book might be regarded as a fairly self-contained introduction to a “generalized cohomology theory”.

The first two chapters (“Vector bundles” and “First notions in *K*-theory”) are chiefly expository; for the reader who is familiar with this material, a brief glance will serve to acquaint him with the notation and approach used. Chapter III is devoted to proving the Bott periodicity theorems. We employ various techniques following the proofs given by Atiyah and Bott [1], Wood [1] and the author [2], using a combination of functional analysis and “algebraic *K*-theory”.

Chapter IV deals with the computation of particular *K*-groups of a large The version of the “Thom isomorphism” in Section IV.5 is mainly due to Atiyah, Bott and Shapiro [1] (in fact they were responsible for the introduction of Clifford algebras in *K*-theory, one of the techniques which we employ in Chapter III).

Chapter V describes some applications of K -theory to the question of H -space structures on the sphere and the Hopf invariant (Adams and Atiyah [1]), and to the solution of the vector field problem (Adams [1]). We also present a sketch of the theory of characteristic classes, which we apply in the proof of the Atiyah–Hirzebruch integrality theorems [1]. In the last section we use K -theory to make some computations on the stable homotopy groups of spheres, via the groups $J(X)$ (cf. Adams [2], Atiyah [1], and Kervaire–Milnor [1]).

In spite of its relative length, this book is certainly not exhaustive in its coverage of K -theory. We have omitted some important topics, particularly those which are presented in detail in the literature. For instance, the Atiyah–Singer index theorem is proved in Cartan–Schwartz [1], Palais [1], and Atiyah–Singer [2] (see also appendix 3 in Hirzebruch [2] for the concepts involved). The relationship between other cohomology theories and K -theory is only sketched in Sections V.3 and V.4. A more complete treatment can be found in Conner–Floyd [1] and Hilton [2] (Atiyah–Hirzebruch spectral sequence). Finally algebraic K -theory is a field which is also growing very quickly at present. Some of the standard references at this time are Bass’s book [1] and the Springer Lecture Notes in Mathematics, Vol. 341, 342, and 343.

I would like to close this foreword with sincere thanks to Maria Klawe, who greatly helped me in the translation of the original manuscript from French to English.

Paris, Summer 1977

Max Karoubi

Remarks on Notation and Terminology


The following notation is used throughout the book: \mathbb{Z} integers, \mathbb{Q} rational numbers, \mathbb{R} real numbers, \mathbb{C} complex numbers, \mathbb{H} quaternions; $GL_n(A)$ denotes the group of invertible $n \times n$ matrices with coefficients in the ring A . The notation $* \cdots *$ signifies an assertion in the text which is not a direct consequence of the theorems proved in this book, but which may be found in the literature; these assertions are not referred to again, except occasionally in exercises.

If \mathcal{C} is a category, and if E and F are objects of \mathcal{C} , then the symbol $\mathcal{C}(E, F)$ or $\text{Hom}_{\mathcal{C}}(E, F)$ means the set of morphisms from E to F .

More specific notation is listed at the end of the book.

A reference to another part of the book is usually given by two numbers (e.g. 5.21) if it is in the same chapter, or by three numbers (e.g. IV.6.7) if it is in a different chapter.

Interdependence of Chapters and Sections



Summary of the Book by Sections

Chapter I. Vector Bundles

1. *Quasi-vector bundles.* This section covers the general concepts and definitions necessary to introduce Section 2. Theorem 1.12 is particularly important in the sequel.

2. *Vector bundles.* The “vector bundles” considered here are locally trivial vector bundles whose fibers are finite dimensional vector spaces over \mathbb{R} or \mathbb{C} . To be mentioned: Proposition 2.7 and Examples 2.3 and 2.4 will be referred to in the sequel.

3. *Clutching theorems.* This technical section is necessary to provide a bridge between the theory of vector bundles and the theory of “coordinate bundles” of N. Steenrod [1]. The clutching theorems are useful in the construction of the tangent bundle of a differentiable manifold (3.18) and in the description of vector bundles over spheres (3.9; see also I.7.6).

4. *Operations on vector bundles.* Certain “continuous” operations on finite dimensional vector spaces: direct sum, tensor product, duality, exterior powers, etc. . . . can be also defined on the category of vector bundles.

5. *Sections of vector bundles.* Only continuous sections are considered here. The major topic concerns the solution of problems involving extensions of sections over paracompact spaces.

6. *Algebraic properties of the category of vector bundles.* In this section we prove that the category $\mathcal{E}(X)$ of vector bundles over a compact space X , is a “pseudo-abelian additive” category. Essentially this means that one has direct sums of vector bundles (the “Whitney sum”), and that every projection operator has an image. From this categorical description (6.13), we deduce the theorem of Serre and Swan (6.18): The category $\mathcal{E}(X)$ is equivalent to the category $\mathcal{P}(A)$, where A is the ring of continuous functions on X , and $\mathcal{P}(A)$ is the category of finitely generated projective modules over A .

7. *Homotopy and representability theorems.* This section is essential for the following chapters. We prove that the problem of classification of vector bundles

with compact base X depends only on the homotopy type of X (7.2). We also prove that $\Phi_n^k(X)$ (the set of isomorphism classes of k -vector bundles, over X of rank n for $k = \mathbb{R}$ or \mathbb{C}), considered as a functor of X , is a direct limit of representable functors. This takes the concrete form of Theorems 7.10 and 7.14.

8. *Metrics and forms on vector bundles.* It is sometimes important to have some additional structure on vector bundles, such as bilinear forms, Hermitian forms, etc. With the exception of Theorem 8.7, this section is not used in the following chapters (except in the exercises).

Chapter II. First Notions of K -Theory

1. *The Grothendieck group of an additive category. The group $K(X)$.* Starting with the simple notion of symmetrization of an abelian monoid, we define the group $K(\mathcal{C})$ of an additive category using the monoid of isomorphism classes of objects of \mathcal{C} . Considering the case where \mathcal{C} is $\mathcal{E}(X)$ and X is compact, we obtain the group $K(X)$ (actually $K_{\mathbb{R}}(X)$ or $K_{\mathbb{C}}(X)$ according to which theory of vector bundles is considered). We prove that $K_{\mathbb{R}}(X) \approx [X, \mathbb{Z} \times BO]$ and $K_{\mathbb{C}}(X) \approx [X, \mathbb{Z} \times BU]$ (1.33).

2. *The Grothendieck group of an additive functor. The group $K(X, Y)$.* In order to obtain a "reasonable" definition of the Grothendieck group $K(\varphi)$ for an additive functor $\varphi: \mathcal{C} \rightarrow \mathcal{C}'$, which generalizes the definition of $K(\mathcal{C})$ when $\mathcal{C}' = 0$, we assume some topological conditions on the categories \mathcal{C} and \mathcal{C}' and on the functor φ (2.6). Since these conditions are satisfied by the "restriction" functor $\mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ where Y is closed in X , we then define the "relative group" $K(X, Y)$ to be the K -group of this functor. In fact, $K(X, Y) \approx \tilde{K}(X/Y)$ (2.35). This isomorphism shows that essentially we do not obtain a new group; however, the groups $K(\varphi)$ and $K(X, Y)$ will be important technical tools later on.

3. *The group K^{-1} of a Banach category. The group $K^{-1}(X)$.* This section represents the first step towards the construction of a cohomology theory h^* where the term h^0 is the group $K(X, Y)$ (also denoted by $K^0(X, Y)$) considered in II.2. The group $K^{-1}(\mathcal{C})$, where \mathcal{C} is a Banach category, is obtained from the automorphisms of objects of \mathcal{C} . Again, if we consider the case where \mathcal{C} is $\mathcal{E}(X)$, we obtain the group called $K^{-1}(X)$. We prove that if Y is a closed subspace of X then the sequence

$$K^{-1}(X) \rightarrow K^{-1}(Y) \rightarrow K(X, Y) \rightarrow K(X) \rightarrow K(Y) \text{ is exact.}$$

We also prove that $K_{\mathbb{R}}^{-1}(X) \approx [X, 0]$ and $K_{\mathbb{C}}^{-1}(X) \approx [X, U]$ (3.19).

4. *The groups $K^{-n}(X)$ and $K^{-n}(X, Y)$.* The aim of this section is to define the groups $K^{-n}(X, Y)$ for $n \geq 2$ and to establish the exact sequence

$$K^{-n-1}(X) \rightarrow K^{-n-1}(Y) \rightarrow K^{-n}(X, Y) \rightarrow K^{-n}(X) \rightarrow K^{-n}(Y), \quad \text{for } n \geq 1$$

One possible definition is $K^{-n}(X, Y) = \tilde{K}(S^n(X/Y))$ (4.12). We prove some "Mayer-Vietoris exact sequences" (4.18 and 4.19) which will be very useful later on.

5. Multiplicative structures. The tensor product of vector bundles provides the group $K(X)$ with a ring structure. It is more difficult to define a "cup-product"

$$K(X, Y) \times K(X', Y') \rightarrow K(X \times X', X \times Y' \cup Y \times X')$$

or more generally

$$K^{-n}(X, Y) \times K^{-n'}(X', Y') \rightarrow K^{-n-n'}(X \times X', X \times Y' \cup Y \times X')$$

when Y and Y' are non-empty. This is accomplished in a theoretical sense in proposition 5.6; however, in applications it is often useful to have more explicit formulas. For this we introduce another definition of the group $K(X, Y)$ by putting metrics on the vector bundles involved (5.16). This will not be used before Chapter IV. The existence of such cup-products shows that there is a direct splitting $K(X) \approx H^0(X; \mathbb{Z}) \oplus K'(X)$ where $K'(X)$ is a nil ideal (cf. 5.9; note that $K'(X) \approx \tilde{K}(X)$ if X is connected).

Chapter III. Bott Periodicity

1. Periodicity in complex K -theory. In this section we define an isomorphism $K_{\mathbb{C}}^{-n}(X, Y) \approx K_{\mathbb{C}}^{-n-2}(X, Y)$. The method (due to Atiyah, Bott, and Wood) is to reduce this isomorphism for general n , to a theorem on Banach algebras (1.11): If A is a complex Banach algebra, the group $K(A)$ (defined as $K(\mathcal{P}(A))$) is naturally isomorphic to $\pi_1(\mathrm{GL}(A))$ where $\mathrm{GL}(A) = \mathrm{inj} \lim \mathrm{GL}_n(A)$. This theorem is proved using the Fourier series of a continuous function with values in a complex Banach space, and some classical results in Algebraic K -theory on Laurent polynomials. The original theorem follows when we let A be the ring of complex continuous functions on a compact space.

2. First applications of Bott periodicity theorem in the complex case. As a first application we obtain the classical theorem of Bott: for $n > i/2$, we have $\pi_i(\cup(n)) \approx \mathbb{Z}$ if i is odd and $\pi_i(\cup(n)) = 0$ for i even. We also prove that real K -theory is periodic of period 4 mod. 2-torsion: $K_{\mathbb{R}}^{-n}(X, Y) \otimes_{\mathbb{Z}} \mathbb{Z}' \approx K_{\mathbb{R}}^{-n-4}(X, Y) \otimes_{\mathbb{Z}} \mathbb{Z}'$, where $\mathbb{Z}' = \mathbb{Z}[\frac{1}{2}]$. This theorem will be strengthened in III.5.

3. Clifford algebras. These algebras play an important role in real K -theory and will be used in Chapter IV in both real and complex K -theory. This section is purely algebraic. The essential result is Theorem 3.21, which establishes a kind of periodicity for Clifford algebras. This "algebraic" periodicity will be effectively used in III.5 to prove the "topological" periodicity of real K -theory and at the same time give another proof of the periodicity of complex K -theory.

4. The functors $K^{p,q}(\mathcal{C})$ and $K^{p,q}(X)$. The idea of this section is to use the Clifford algebras $C^{p,q}$ to algebraically define new functors $K^n(X) = K^{p,q}(X)$ for $n = p - q \in \mathbb{Z}$. We prove that these functors are *by definition* periodic, of period 8 in the real case, and of period 2 in the complex case, and that $K^0(X)$ and $K^{-1}(X)$ are indeed the functors defined in Chapter II. Bott periodicity will then be proved if we show that the two definitions of $K^n(X)$ agree for negative values of n . This is done in the next two sections.

5. The functors $K^{p,q}(X, Y)$ and the isomorphism t . Periodicity in real K -theory. After some preliminaries introducing the relative groups $K^{p,q}(X, Y)$ we present the fundamental theorem of this chapter: The groups $K^{p,q+1}(X, Y)$ and $K^{p,q}(X \times B^1, X \times S^0 \cup Y \times B^1)$ are isomorphic. Assuming this theorem (the proof follows in Section III.6), we prove that $K_{\mathbb{R}}^{-n}(X, Y) \approx K_{\mathbb{R}}^{-n-8}(X, Y)$ with the definitions of Chapter II. At the same time we prove the periodicity in complex K -theory (5.17) once more. Moreover, using Propositions 4.29 and 4.30 we prove the existence of weak homotopy equivalences between the iterated loop spaces $\Omega^i(0)$ and certain homogeneous spaces (5.22). We also compute the homotopy groups $\pi_i(0(n))$ for $n > i + 1$ (5.19) with the help of Clifford algebras.

6. Proof of the fundamental theorem. The pattern of this section is analogous to that of Section III.1, since the main theorem is likewise a consequence of a general theorem on Banach algebras (6.12). Moreover the proof of this general theorem uses the same ideas as the proof of Theorem 1.11.

Chapter IV. Computations of Some K -Groups

1. The Thom isomorphism in complex K -theory for complex vector bundles. The purpose of this section is to compute the complex K -theory of the Thom space of a complex vector bundle (1.9). In this computation a key role is played by bundles of exterior algebras. Theorem 1.3. is particularly important in the sequel.

2. Complex K -theory of complex projective spaces and complex projective bundles. In this section (classical in style), we construct a method which may also be used for ordinary cohomology (see V.3). Using the technical Proposition 2.4 we are able to compute the K -theory of $P_n = P(\mathbb{C}^{n+1})$ and more generally of $P(V)$ where V is a complex vector bundle (2.13). The “splitting principle” (2.15) is used frequently later on. With this principle we are able to make the multiplicative structure of $K^*(P(V))$ explicit (2.16).

3. Complex K -theory of flag bundles and Grassmann bundles. K -theory of a product. This section is also classical in style, but is not essential to the sequel. We explicitly compute $K^*(F(V))$ where $F(V)$ is the flag bundle of a complex vector

bundle V . We also compute $K^*(G_p(V))$ where $G_p(V)$ is the fiber bundle of p -subspaces in V (3.12). These results are used to compute $\mathcal{K}(BU(n)) = \text{proj lim } K(G_p(\mathbb{C}^n))$ (3.22), and the K -theory of a product (3.27).

4. Complements in Clifford algebras. The concept of “spinors” was not introduced in Section III.3, since it is not essential in proving Bott periodicity. However we now need this concept to prove the analog of Thom’s theorem in K -theory (for real or complex vector bundles). After some algebraic preliminaries we study the possibilities of lifting the structural group of a real vector bundle to the spinorial group $\text{Spin}(n)$ or $\text{Spin}^c(n)$. Theorem 4.22 is particularly important for our purpose.

5. The Thom isomorphism in real and complex K -theory for real vector bundles. As in IV.1, the purpose of this section is to compute the K -theory of the Thom space of a vector bundle, but now the vector bundle is real, and the K -theory used is real or complex. With an additional spinorial hypothesis, we prove that $K(V) \approx K^{-n}(X)$ if n is the rank of V . If the base is compact and n is a multiple of 8 (of 2 in complex K -theory), we prove that $K(V)$ is a $K(X)$ -module of rank one generated by the “Thom class” T_V . Finally, if $f: X \rightarrow Y$ is a proper continuous map between differentiable manifolds and if $\text{Dim}(Y) - \text{Dim}(X) = 0 \pmod{8}$ ($\pmod{2}$ in the complex case), we define, with an additional spinorial hypothesis, a “Gysin homomorphism” $f_*: K(X) \rightarrow K(Y)$ which is analogous to the Gysin homomorphism in ordinary cohomology. This homomorphism is only used in V.4.

6. Real and complex K -theory of real projective spaces and real projective bundles. This section is much more technical than the others (the results are only used in V.2). After some easy but tedious lemmas making systematic use of Clifford algebras, we are able to compute (up to extension) the real and complex K -theory of a real projective bundle (6.40 and 6.42). In the case of real projective spaces, the K -theory is completely determined (6.46 and 6.47).

7. Operations in K -theory. One of the charms of K -theory is that we are able to define some very nice operations. For example, there are the exterior power operations λ^k (due to Grothendieck). By a method due to Atiyah we determine all the operations in complex K -theory. With this method we show that the “Adams operations” ψ^k are the only ring operations in complex K -theory (7.13). They will be very useful in applications.

The operations λ^k and ψ^k may also be defined in real K -theory. However, their properties are more difficult to prove. We must refer to Adams [3] or Exercise 8.5 for a complete proof. From the operations ψ^k , we obtain the operations ρ^k , which will be very useful in V.2 and V.5.

Chapter V. Some Applications of K -Theory

1. H -space structures on spheres and the Hopf invariant. Using the Adams operations in complex K -theory, we prove that the only spheres which admit an H -space

structure are S^1 , S^3 , and S^7 . In fact, we prove more: if $f: S^{2n-1} \rightarrow S^n$ is a map of odd Hopf invariant, then n must be 2, 4 or 8.

2. The solution of the vector field problem on the sphere. Let us write every integer t in the form $(2\alpha - 1) \cdot 2^\beta$, for $\beta = \gamma + 4\delta$ with $0 \leq \gamma \leq 3$, and define $\rho(t) = 2^\gamma + 8\delta$. Then the maximum number of independent vector fields on the sphere S^{t-1} is exactly $\rho(t) - 1$ (2.10). The proof of this classical theorem is "elementary" (in the context of this book) and uses essentially the operations ρ^k in the real K -theory of real projective spaces.

3. Characteristic classes and the Chern character. For each complex vector bundle V , we define "Chern classes" $c_i(V) \in H^{2i}(X; \mathbb{Z})$ in an axiomatic way (3.15). The construction of these classes is analogous to the construction of classes done in Section IV.3. By means of these classes, we construct a fundamental homomorphism, the "Chern character", from $K_{\mathbb{C}}(X)$ to $H^{\text{even}}(X; \mathbb{Q})$. The Chern character induces an isomorphism between $K_{\mathbb{C}}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$ and $H^{\text{even}}(X; \mathbb{Q})$ for every compact X .

4. The Riemann–Roch theorem and integrality theorems. To each complex stable vector bundle (resp. "spinorial real stable bundle") we associate an important characteristic class $\tau(V)$, called the Todd class (resp. $A(V)$, called the Atiyah–Hirzebruch class). These classes play an important role in the "differentiable Riemann–Roch theorem": For each suitably continuous map $f: X \rightarrow Y$ and for each element x of $K_{\mathbb{C}}(X)$, we have the formula $ch(f_*^K(x)) = f_*^H(A(v_f) \cdot ch(x))$ where $A(v_f)$ denotes the Atiyah–Hirzebruch class of the stable bundle $f^*(TY) - TX$ (assuming that $\text{Dim}(Y) = \text{Dim}(X) \bmod 2$ and that there is a stable "spinorial structure on v_f "). From this theorem we obtain integral theorems for characteristic classes (4.21) and the homotopy invariance of certain characteristic classes (4.24).

5. Applications of K -theory to stable homotopy. In this section we explain how K -theory may be applied to obtain some interesting information about the stable homotopy groups of spheres. We only include those partial results which can be obtained from the material in this book. More complete results are found in the series of J. F. Adams on the groups $J(X)$ [2], and in Husemoller's book [1].

Table of Contents

Summary of the Book by Sections	XIII
---	------

Chapter I. Vector Bundles

1. Quasi-Vector Bundles	1
2. Vector Bundles	4
3. Clutching Theorems	7
4. Operations on Vector Bundles	18
5. Sections of Vector Bundles	21
6. Algebraic Properties of the Category of Vector Bundles	26
7. Homotopy Theory of Vector Bundles	35
8. Metrics and Forms on Vector Bundles	40
9. Exercises	44
10. Historical Note	51

Chapter II. First Notions of K -Theory

1. The Grothendieck Group of a Category. The Group $K(X)$	52
2. The Grothendieck Group of a Functor. The Group $K(X, Y)$	59
3. The Group K^{-1} of a Banach Category. The Group $K^{-1}(X)$	71
4. The Groups $K^{-n}(X)$ and $K^{-n}(X, Y)$	82
5. Multiplicative Structures	92
6. Exercises	105
7. Historical Note	111

Chapter III. Bott Periodicity

1. Periodicity in Complex K -Theory	113
2. First Applications of Bott Periodicity Theorem in Complex K -Theory	123
3. Clifford Algebras	127
4. The Functors $K^{p,q}(\mathcal{C})$ and $K^{p,q}(X)$	137

5. The Functors $K^{p,q}(X, Y)$ and the Isomorphism t . Periodicity in Real K -Theory	148
6. Proof of the Fundamental Theorem	161
7. Exercises	174
8. Historical Note	179

Chapter IV. Computation of Some K -Groups

1. The Thom Isomorphism in Complex K -Theory for Complex Vector Bundles	180
2. Complex K -Theory of Complex Projective Spaces and Complex Projective Bundles	188
3. Complex K -Theory of Flag Bundles and Grassmann Bundles. K -Theory of a Product	196
4. Complements in Clifford Algebras	208
5. The Thom Isomorphism in Real and Complex K -Theory for Real Vector Bundles	221
6. Real and Complex K -Theory of Real Projective Spaces and Real Projective Bundles	234
7. Operations in K -Theory	252
8. Exercises	265
9. Historical Note	268

Chapter V. Some Applications of K -Theory

1. H -Space Structures on Spheres and the Hopf Invariant	270
2. The Solution of the Vector Field Problem on the Sphere	273
3. Characteristic Classes and the Chern Character	277
4. The Riemann–Roch Theorem and Integrality Theorems	284
5. Applications of K -Theory to Stable Homotopy	293
6. Historical Note	299

Bibliography	301
-------------------------------	-----

List of Notation	304
-----------------------------------	-----

Index	307
------------------------	-----

Vector Bundles

1. Quasi-Vector Bundles

1.1. Let k be the field of real numbers or complex numbers¹⁾, and let X be a topological space.

1.2. Definition. A *quasi-vector bundle* with base X is given by

- 1) a finite dimensional k -vector space E_x for every point x of X ,
- 2) a topology on the disjoint union $E = \bigsqcup E_x$ which induces the natural topology on each E_x , such that the obvious projection $\pi: E \rightarrow X$ is continuous.

1.3. Example. Let X be the sphere $S^n = \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\}$. For every point x of S^n we choose E_x to be the vector space orthogonal to x . Then $E = \bigsqcup E_x$ is naturally a subspace of $S^n \times \mathbb{R}^{n+1}$ and may be provided with the induced topology.

1.4. Example. Starting from the preceding example, let us arbitrarily choose a vector space $F_x \subset E_x$ for each $x \in S^n$; then if F is given the induced topology again we have a quasi-vector bundle on X .

More examples are given in the following sections.

1.5. A quasi-vector bundle is denoted by $\xi = (E, \pi, X)$ or simply by E if there is no risk of confusion. The space E is the *total space* of ξ and E_x is the *fiber* of ξ at the point x .

1.6. Let $\xi = (E, \pi, X)$ and $\xi' = (E', \pi', X')$ be quasi-vector bundles. A *general morphism* from ξ to ξ' is given by a pair (f, g) of continuous maps $f: X \rightarrow X'$ and $g: E \rightarrow E'$ such that

- 1) the diagram

$$\begin{array}{ccc} E & \xrightarrow{g} & E' \\ \pi \downarrow & & \downarrow \pi' \\ X & \xrightarrow{f} & X' \end{array}$$

is commutative.

¹⁾ In general, these are the most interesting cases; however, sometimes we will use the field of quaternions \mathbb{H} .

2) The map $g_x: E_x \rightarrow E'_{f(x)}$ induced by g is k -linear.

General morphisms can be composed in an obvious way. In this way we construct a category whose objects are quasi-vector bundles and whose arrows are general morphisms.

1.7. If ξ and ξ' have the same base $X = X'$, a *morphism* between ξ and ξ' is a general morphism (f, g) such that $f = \text{Id}_X$. Such a morphism will be simply called g in the sequel. The quasi-vector bundles with the same base X are the objects of a subcategory, whose arrows are the morphisms we have just defined.

1.8. Example. Let us return to Example 1.3, and let $n=1$. Let $\xi' = (E', \pi', X')$ where $X = X' = S^1$, and $E' = S^1 \times \mathbb{R}$ with the product topology. If we identify \mathbb{R}^2 with the complex numbers as usual, we can define a continuous map $g: E \rightarrow E'$ by the formula $g(x, z) = (x, iz/x)$ (this is well defined because x is orthogonal to z in $\mathbb{R}^2 = \mathbb{C}$). In fact g is an isomorphism between E and E' in the category described in 1.7.

1.9. Example. Let E'' be the quotient of $E' = S^1 \times \mathbb{R}$ by the equivalence relation $(x, t) \sim (y, u)$ if $y = \varepsilon x$ and $u = \varepsilon t$ with $\varepsilon = \pm 1$. Then E'' is the total space of a quasi-vector bundle over $P_1(\mathbb{R})$ called the *infinite Moebius band*. By identifying $P_1(\mathbb{R})$ with S^1 by the map $z \mapsto z^2$, we see easily that E'' is also the quotient of $I \times \mathbb{R}$ by the equivalence relation which identifies $(0, u)$ with $(1, -u)$. If we restrict u to have norm less than 1, we obtain the classical Moebius band.

We claim that the bundles E' and E'' over S^1 are not isomorphic. Suppose there exists an isomorphism $g: E' \rightarrow E''$; then we must have $E' - X'$ homeomorphic to $E'' - X''$ where X' (and X'') denote the set of points of the form $(x, 0)$ with $x \in S^1$ (note that $X'' \approx X'$). But $E'' - X''$ is connected and $E' - X'$ is not.

1.10. Let V be a finite dimensional vector space (as always over k). The preceding examples show the importance of quasi-vector bundles of the form $E = X \times V$, as models. To be more precise, $E_x = V$ and the total space may be identified with $X \times V$ provided with the product topology. Such bundles are called *trivial quasi-vector bundles* or simply *trivial vector bundles*.

1.11. Let $E = X \times V$ and $E' = X \times V'$ be trivial vector bundles with base X . We want to explicitly describe the morphisms from E to E' (again in the category defined in 1.7). Since the diagram

$$\begin{array}{ccc} X \times V & \longrightarrow & X \times V' \\ & \searrow & \swarrow \\ & X & \end{array}$$

is commutative, for each point x of X , g induces a linear map $g_x: V \rightarrow V'$. Let $\check{g}: X \rightarrow \mathcal{L}(V, V')$ be the map defined by $\check{g}(x) = g_x$.