

**McGRAW-HILL
BOOK COMPANY**

New York

St. Louis

San Francisco

Düsseldorf

Johannesburg

Kuala Lumpur

London

Mexico

Montreal

New Delhi

Panama

Paris

São Paulo

Singapore

Sydney

Tokyo

Toronto

WALTER RUDIN

Professor of Mathematics

University of Wisconsin, Madison

Real and Complex Analysis

SECOND EDITION

**McGRAW-HILL
BOOK COMPANY**

New York

St. Louis

San Francisco

Düsseldorf

Johannesburg

Kuala Lumpur

London

Mexico

Montreal

New Delhi

Panama

Paris

São Paulo

Singapore

Sydney

Tokyo

Toronto

WALTER RUDIN

Professor of Mathematics

University of Wisconsin, Madison

Real and Complex Analysis

SECOND EDITION

This book was set in Times Roman by Composition Technology, Inc.
The editors were Jack L. Farnsworth and M. E. Margolies;
the production supervisor was Thomas J. LoPinto.
The Maple Press Company was printer and binder.

Library of Congress Cataloging in Publication Data

Rudin, Walter, date
Real and complex analysis.

(McGraw-Hill series in higher mathematics)

Bibliography: p.

1. Mathematical analysis. I. Title.

QA300.R82 1974 515 73-15743

ISBN 0-07-054233-3

**REAL AND
COMPLEX ANALYSIS**

Copyright © 1966, 1974 by McGraw-Hill, Inc. All rights reserved.
Printed in the United States of America. No part of this publication may be reproduced,
stored in a retrieval system, or transmitted, in any form or by any means,
electronic, mechanical, photocopying, recording, or otherwise,
without the prior written permission of the publisher.

1234567890MAMM79876543

PREFACE

This book contains a first-year graduate course in which the basic techniques and theorems of analysis are presented in such a way that the intimate connections between its various branches are strongly emphasized. The traditionally separate subjects of “real analysis” and “complex analysis” are thus united; some of the basic ideas from functional analysis are also included.

Here are some examples of the way in which these connections are demonstrated and exploited. The Riesz representation theorem and the Hahn-Banach theorem allow one to “guess” the Poisson integral formula. They team up in the proof of Runge’s theorem. They combine with Blaschke’s theorem on the zeros of bounded holomorphic functions to give a proof of the Müntz-Szasz theorem, which concerns approximation on an interval. The fact that L^2 is a Hilbert space is used in the proof of the Radon-Nikodym theorem, which leads to the theorem about differentiation of indefinite integrals (incidentally, differentiation seems to be unduly slighted in most modern texts), which in turn yields the existence of radial limits of bounded harmonic functions. The theorems of Plancherel and Cauchy combined give a theorem of Paley and Wiener which, in turn, is used in the Denjoy-Carleman theorem about infinitely differentiable functions on the real line. The maximum modulus theorem gives information about linear transformations on L^p -spaces.

Since most of the results presented here are quite classical (the novelty lies in the arrangement, and some of the proofs are new), I have not attempted to document the source of every item. References are gathered at the end, in Notes and Comments. They are not always to the original sources, but more often to more recent works where further references can be found. In no case does the absence of a reference imply any claim to originality on my part.

The prerequisite for this book is a good course in advanced calculus (set-theoretic manipulations, metric spaces, uniform continuity, and uniform convergence). The first seven chapters of my earlier book "Principles of Mathematical Analysis" furnish sufficient preparation.

Experience with the first edition shows that first-year graduate students can study the first 15 chapters in two semesters, plus some topics from 1 or 2 of the remaining 5. These latter are quite independent of each other. The first 15 should be taken up in the order in which they are presented, except for Chapter 9, which can be postponed.

Some new exercises have been added in this second edition, and many of the old ones have been regrouped so that they now appear in more or less the same order in which the corresponding topics occur in the text.

The text contains two substantial changes. The first of these was suggested by Jim Serrin, who showed me how to modify my earlier treatment of the differentiation of measures so as to obtain stronger results with no extra effort.

The second one is the inclusion of John Dixon's recently discovered, beautifully simple proof of the global (homology) version of Cauchy's theorem. This can now be proved and used as soon as some basic local properties of holomorphic functions are known. The order of several topics has accordingly been changed.

I have also made many smaller changes in order to improve some details and clarify some obscure points. Almost all of these were suggested by students, colleagues, and other friends. Their constructive comments and criticisms were greatly appreciated. I take this opportunity to thank them.

WALTER RUDIN

CONTENTS

Preface	xi
Prologue: The Exponential Function	1
1 Abstract Integration	5
Set-theoretic notations and terminology	6
The concept of measurability	8
Simple functions	16
Elementary properties of measures	17
Arithmetic in $[0, \infty]$	19
Integration of positive functions	20
Integration of complex functions	25
The role played by sets of measure zero	28
Exercises	32
2 Positive Borel Measures	34
Vector spaces	34
Topological preliminaries	36
The Riesz representation theorem	42
Regularity properties of Borel measures	49
Lebesgue measure	51
Continuity properties of measurable functions	56
Exercises	58

CONTENTS

3	L^p-Spaces	62
	Convex functions and inequalities	62
	The L^p -spaces	66
	Approximation by continuous functions	71
	Exercises	73
4	Elementary Hilbert Space Theory	79
	Inner products and linear functionals	79
	Orthonormal sets	86
	Trigonometric series	92
	Exercises	97
5	Examples of Banach Space Techniques	100
	Banach spaces	100
	Consequences of Baire's theorem	102
	Fourier series of continuous functions	106
	Fourier coefficients of L^1 -functions	109
	The Hahn-Banach theorem	111
	An abstract approach to the Poisson integral	115
	Exercises	119
6	Complex Measures	124
	Total variation	124
	Absolute continuity	128
	Consequences of the Radon-Nikodym theorem	133
	Bounded linear functionals on L^p	135
	The Riesz representation theorem	138
	Exercises	142
7	Integration on Product Spaces	145
	Measurability on cartesian products	145
	Product measures	148
	The Fubini theorem	150
	Completion of product measures	153
	Convolutions	155
	Exercises	157
8	Differentiation	162
	Derivatives of measures	162
	Functions of bounded variation	171
	Differentiation of point functions	175

Differentiable transformations	181
Exercises	188
9 Fourier Transforms	192
Formal properties	192
The inversion theorem	195
The Plancherel theorem	200
The Banach algebra L^1	205
Exercises	208
10 Elementary Properties of Holomorphic Functions	212
Complex differentiation	212
Integration over paths	217
The local Cauchy theorem	221
The power series representation	225
The open mapping theorem	231
The global Cauchy theorem	233
The calculus of residues	241
Exercises	244
11 Harmonic Functions	250
The Cauchy-Riemann equations	250
The Poisson integral	252
The mean value property	259
Positive harmonic functions	261
Exercises	266
12 The Maximum Modulus Principle	270
Introduction	270
The Schwarz lemma	271
The Phragmen-Lindelöf method	273
An interpolation theorem	277
A converse of the maximum modulus theorem	279
Exercises	281
13 Approximation by Rational Functions	284
Preparation	284
Runge's theorem	288
The Mittag-Leffler theorem	291
Simply connected regions	292
Exercises	294

14	Conformal Mapping	296
	Preservation of angles	296
	Linear fractional transformations	298
	Normal families	300
	The Riemann mapping theorem	302
	The class \mathfrak{S}	304
	Continuity at the boundary	308
	Conformal mapping of an annulus	311
	Exercises	313
15	Zeros of Holomorphic Functions	320
	Infinite products	320
	The Weierstrass factorization theorem	323
	An interpolation problem	327
	Jensen's formula	329
	Blaschke products	333
	The Müntz-Szasz theorem	336
	Exercises	339
16	Analytic Continuation	343
	Regular points and singular points	343
	Continuation along curves	347
	The monodromy theorem	351
	Construction of a modular function	352
	The Picard theorem	356
	Exercises	357
17	H^p-Spaces	361
	Subharmonic functions	361
	The spaces H^p and N	363
	The space H^2	365
	The theorem of F. and M. Riesz	369
	Factorization theorems	370
	The shift operator	375
	Conjugate functions	379
	Exercises	382
18	Elementary Theory of Banach Algebras	386
	Introduction	386
	The invertible elements	387
	Ideals and homomorphisms	392

Applications	396
Exercises	400
19 Holomorphic Fourier Transforms	403
Introduction	403
Two theorems of Paley and Wiener	405
Quasi-analytic classes	409
The Denjoy-Carleman theorem	412
Exercises	416
20 Uniform Approximation by Polynomials	419
Introduction	419
Some lemmas	420
Mergelyan's theorem	423
Exercises	427
Appendix: Hausdorff's Maximality Theorem	429
Notes and Comments	432
Bibliography	440
List of Special Symbols	443
Index	445

PROLOGUE

THE EXPONENTIAL FUNCTION

This is the most important function in mathematics. It is defined, for every complex number z , by the formula

$$(1) \quad \exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

The series (1) converges absolutely for every z and converges uniformly on every bounded subset of the complex plane. Thus \exp is a continuous function. The absolute convergence of (1) shows that the computation

$$\sum_{k=0}^{\infty} \frac{a^k}{k!} \sum_{m=0}^{\infty} \frac{b^m}{m!} = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{k=0}^n \frac{n!}{k!(n-k)!} a^k b^{n-k} = \sum_{n=0}^{\infty} \frac{(a+b)^n}{n!}$$

is correct. It gives the important addition formula

$$(2) \quad \exp(a)\exp(b) = \exp(a+b),$$

valid for all complex numbers a and b .

We define the number e to be $\exp(1)$, and shall usually replace $\exp(z)$ by the customary shorter expression e^z . Note that $e^0 = \exp(0) = 1$, by (1).

Theorem

- (a) For every complex z we have $e^z \neq 0$.
 (b) \exp is its own derivative: $\exp'(z) = \exp(z)$.
 (c) The restriction of \exp to the real axis is a monotonically increasing positive function, and

$$e^x \rightarrow \infty \text{ as } x \rightarrow \infty, \quad e^x \rightarrow 0 \text{ as } x \rightarrow -\infty.$$

- (d) There exists a positive number π such that $e^{mi/2} = i$ and such that $e^z = 1$ if and only if $z/(2\pi i)$ is an integer.
 (e) \exp is a periodic function, with period $2\pi i$.
 (f) The mapping $t \rightarrow e^{it}$ maps the real axis onto the unit circle.
 (g) If w is a complex number and $w \neq 0$, then $w = e^z$ for some z .

PROOF By (2), $e^z \cdot e^{-z} = e^{z-z} = e^0 = 1$. This implies (a). Next,

$$\exp'(z) = \lim_{h \rightarrow 0} \frac{\exp(z+h) - \exp(z)}{h} = \exp(z) \lim_{h \rightarrow 0} \frac{\exp(h) - 1}{h} = \exp(z).$$

The first of the above equalities is a matter of definition, the second follows from (2), and the third from (1), and (b) is proved.

That \exp is monotonically increasing on the positive real axis, and that $e^x \rightarrow \infty$ as $x \rightarrow \infty$, is clear from (1). The other assertions of (c) are consequences of $e^x \cdot e^{-x} = 1$.

For any real number t , (1) shows that e^{-it} is the complex conjugate of e^{it} . Thus

$$|e^{it}|^2 = e^{it} \cdot \overline{e^{it}} = e^{it} \cdot e^{-it} = e^{it-it} = e^0 = 1,$$

or

$$(3) \quad |e^{it}| = 1 \quad (t \text{ real}).$$

In other words, if t is real, e^{it} lies on the unit circle. We define $\cos t$, $\sin t$ to be the real and imaginary parts of e^{it} :

$$(4) \quad \cos t = \operatorname{Re}[e^{it}], \quad \sin t = \operatorname{Im}[e^{it}] \quad (t \text{ real}).$$

If we differentiate both sides of Euler's identity

$$(5) \quad e^{it} = \cos t + i \sin t,$$

which is equivalent to (4), and if we apply (b), we obtain

$$\cos' t + i \sin' t = ie^{it} = -\sin t + i \cos t,$$

so that

$$(6) \quad \cos' = -\sin, \quad \sin' = \cos.$$

The power series (1) yields the representation

$$(7) \quad \cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \frac{t^6}{6!} + \dots$$

Take $t = 2$. The terms of the series (7) then decrease in absolute value (except for the first one) and their signs alternate. Hence $\cos 2$ is less than the sum of the first three terms of (7), with $t = 2$; thus $\cos 2 < -\frac{1}{3}$. Since $\cos 0 = 1$ and \cos is a continuous real function on the real axis, we conclude that there is a smallest positive number t_0 for which $\cos t_0 = 0$. We define

$$(8) \quad \pi = 2t_0.$$

It follows from (3) and (5) that $\sin t_0 = \pm 1$. Since

$$\sin'(t) = \cos t > 0$$

on the segment $(0, t_0)$ and since $\sin 0 = 0$, we have $\sin t_0 > 0$, hence $\sin t_0 = 1$, and therefore

$$(9) \quad e^{\pi i/2} = i.$$

It follows that $e^{\pi i} = i^2 = -1$, $e^{2\pi i} = (-1)^2 = 1$, and then $e^{2n\pi i} = 1$ for every integer n . Also, (e) follows immediately:

$$(10) \quad e^{z+2\pi i} = e^z e^{2\pi i} = e^z.$$

If $z = x + iy$, x and y real, then $e^z = e^x e^{iy}$; hence $|e^z| = e^x$. If $e^z = 1$, we therefore must have $e^x = 1$, so that $x = 0$; to prove that $y/2\pi$ must be an integer, it is enough to show that $e^{iy} \neq 1$ if $0 < y < 2\pi$, by (10).

Suppose $0 < y < 2\pi$, and

$$(11) \quad e^{iy/4} = u + iv \quad (u \text{ and } v \text{ real}).$$

Since $0 < y/4 < \pi/2$, we have $u > 0$ and $v > 0$. Also

$$(12) \quad e^{iy} = (u + iv)^4 = u^4 - 6u^2v^2 + v^4 + 4iuv(u^2 - v^2).$$

The right side of (12) is real only if $u^2 = v^2$; since $u^2 + v^2 = 1$, this happens only when $u^2 = v^2 = \frac{1}{2}$, and then (12) shows that

$$e^{iy} = -1 \neq 1.$$

This completes the proof of (d).

We already know that $t \rightarrow e^{it}$ maps the real axis *into* the unit circle. To prove (f), fix w so that $|w| = 1$; we shall show that $w = e^{it}$ for some real t . Write $w = u + iv$, u and v real, and suppose first that $u \geq 0$ and $v \geq 0$. Since $u \leq 1$, the definition of π shows that there exists a t , $0 \leq t \leq \pi/2$, such that $\cos t = u$; then $\sin^2 t = 1 - u^2 = v^2$, and since $\sin t \geq 0$ if $0 \leq t \leq \pi/2$, we have $\sin t = v$. Thus $w = e^{it}$.

If $u < 0$ and $v \geq 0$, the preceding conditions are satisfied by $-iw$. Hence $-iw = e^{it}$ for some real t , and $w = e^{i(t+\pi/2)}$. Finally, if $v < 0$, the preceding two cases show that $-w = e^{it}$ for some real t , hence $w = e^{i(t+\pi)}$. This completes the proof of (f).

If $w \neq 0$, put $\alpha = w/|w|$. Then $w = |w|\alpha$. By (c), there is a real x such that $|w| = e^x$. Since $|\alpha| = 1$, (f) shows that $\alpha = e^{iy}$ for some real y . Hence $w = e^{x+iy}$. This proves (g) and completes the theorem. ///

We shall encounter the integral of $(1 + x^2)^{-1}$ over the real line. To evaluate it, put $\varphi(t) = \sin t/\cos t$ in $(-\pi/2, \pi/2)$. By (6), $\varphi' = 1 + \varphi^2$. Hence φ is a monotonically increasing mapping of $(-\pi/2, \pi/2)$ onto $(-\infty, \infty)$, and we obtain

$$\int_{-\infty}^{\infty} \frac{dx}{1 + x^2} = \int_{-\pi/2}^{\pi/2} \frac{\varphi'(t) dt}{1 + \varphi^2(t)} = \int_{-\pi/2}^{\pi/2} dt = \pi.$$

ABSTRACT INTEGRATION

Toward the end of the nineteenth century it became clear to many mathematicians that the Riemann integral (about which one learns in calculus courses) should be replaced by some other type of integral, more general and more flexible, better suited for dealing with limit processes. Among the attempts made in this direction, the most notable ones were due to Jordan, Borel, W. H. Young, and Lebesgue. It was Lebesgue's construction which turned out to be the most successful.

In brief outline, here is the main idea: The Riemann integral of a function f over an interval $[a, b]$ can be approximated by sums of the form

$$\sum_{i=1}^n f(t_i)m(E_i)$$

where E_1, \dots, E_n are disjoint intervals whose union is $[a, b]$, $m(E_i)$ denotes the length of E_i , and $t_i \in E_i$ for $i = 1, \dots, n$. Lebesgue discovered that a completely satisfactory theory of integration results if the sets E_i in the above sum are allowed to belong to a larger class of subsets of the line, the so-called "measurable sets," and if the class of functions under consideration is enlarged to what he called "measurable functions." The crucial set-theoretic properties involved are the following: The union and the intersection of any countable family of measurable

sets are measurable; so is the complement of every measurable set; and, most important, the notion of "length" (now called "measure") can be extended to them in such a way that

$$m(E_1 \cup E_2 \cup E_3 \cup \cdots) = m(E_1) + m(E_2) + m(E_3) + \cdots$$

for every countable collection $\{E_i\}$ of pairwise disjoint measurable sets. This property of m is called *countable additivity*.

The passage from Riemann's theory of integration to that of Lebesgue is a process of completion (in a sense which will appear more precisely later). It is of the same fundamental importance in analysis as is the construction of the real number system from the rationals.

The above-mentioned measure m is of course intimately related to the geometry of the real line. In this chapter we shall present an abstract (axiomatic) version of the Lebesgue integral, relative to *any* countably additive measure on *any* set. (The precise definitions follow.) This abstract theory is not in any way more difficult than the special case of the real line; it shows that a large part of integration theory is independent of any geometry (or topology) of the underlying space; and, of course, it gives us a tool of much wider applicability. The existence of a large class of measures, among them that of Lebesgue, will be established in Chap. 2.

Set-Theoretic Notations and Terminology

1.1 Some sets can be described by listing their members. Thus $\{x_1, \dots, x_n\}$ is the set whose members are x_1, \dots, x_n ; and $\{x\}$ is the set whose only member is x . More often, sets are described by properties. We write

$$\{x: P\}$$

for the set of all elements x which have the property P . The symbol \emptyset denotes the empty set. The words *collection*, *family*, and *class* will be used synonymously with *set*.

We write $x \in A$ if x is a member of the set A ; otherwise $x \notin A$. If B is a subset of A , i.e., if $x \in B$ implies $x \in A$, we write $B \subset A$. If $B \subset A$ and $A \subset B$, then $A = B$. If $B \subset A$ and $A \neq B$, B is a *proper* subset of A . Note that $\emptyset \subset A$ for every set A .

$A \cup B$ and $A \cap B$ are the union and intersection of A and B , respectively. If $\{A_\alpha\}$ is a collection of sets, where α runs through some index set I , we write

$$\bigcup_{\alpha \in I} A_\alpha \quad \text{and} \quad \bigcap_{\alpha \in I} A_\alpha$$

for the union and intersection of $\{A_\alpha\}$:

$$\begin{aligned}\bigcup_{\alpha \in I} A_\alpha &= \{x: x \in A_\alpha \text{ for at least one } \alpha \in I\} \\ \bigcap_{\alpha \in I} A_\alpha &= \{x: x \in A_\alpha \text{ for every } \alpha \in I\}.\end{aligned}$$

If I is the set of all positive integers, the customary notations are

$$\bigcup_{n=1}^{\infty} A_n \quad \text{and} \quad \bigcap_{n=1}^{\infty} A_n.$$

If no two members of $\{A_\alpha\}$ have an element in common, then $\{A_\alpha\}$ is a *disjoint collection* of sets.

We write $A - B = \{x: x \in A, x \notin B\}$, and denote the complement of A by A^c whenever it is clear from the context with respect to which larger set the complement is taken.

The *cartesian product* $A_1 \times \cdots \times A_n$ of the sets A_1, \dots, A_n is the set of all ordered n -tuples (a_1, \dots, a_n) where $a_i \in A_i$ for $i = 1, \dots, n$.

The *real line* (or real number system) is R^1 , and

$$R^k = R^1 \times \cdots \times R^1 \quad (k \text{ factors}).$$

The *extended real number system* is R^1 with two symbols, ∞ and $-\infty$, adjoined, and with the obvious ordering. If $-\infty \leq a \leq b \leq \infty$, the *interval* $[a, b]$ and the *segment* (a, b) are defined to be

$$[a, b] = \{x: a \leq x \leq b\}, \quad (a, b) = \{x: a < x < b\}.$$

We also write

$$[a, b) = \{x: a \leq x < b\}, \quad (a, b] = \{x: a < x \leq b\}.$$

If $E \subset [-\infty, \infty]$ and $E \neq \emptyset$, the least upper bound (supremum) and greatest lower bound (infimum) of E exist in $[-\infty, \infty]$ and are denoted by $\sup E$ and $\inf E$.

Sometimes (but only when $\sup E \in E$) we write $\max E$ for $\sup E$.

The symbol

$$f: X \rightarrow Y$$

means that f is a *function* (or *mapping* or *transformation*) of the set X into the set Y ; i.e., f assigns to each $x \in X$ an element $f(x) \in Y$. If $A \subset X$ and $B \subset Y$, the *image* of A and the *inverse image* (or pre-image) of B are

$$\begin{aligned}f(A) &= \{y: y = f(x) \text{ for some } x \in A\}, \\ f^{-1}(B) &= \{x: f(x) \in B\}.\end{aligned}$$