

# Graduate Texts in Mathematics

**John C. Oxtoby**

## **Measure and Category**

**Second Edition**

2

**测度与范畴学 第2版**

**Springer**

世界图书出版公司  
[www.wpcbj.com.cn](http://www.wpcbj.com.cn)

John C. Oxtoby

0154/Y2=2

2009.

# Measure and Category

A Survey of the Analogies between  
Topological and Measure Spaces

Second Edition



Springer-Verlag  
New York Heidelberg Berlin

**John C. Oxtoby**

Department of Mathematics  
Bryn Mawr College  
Bryn Mawr, PA 19010  
USA

### ***Editorial Board***

**P. R. Halmos**

*Managing Editor*  
Department of Mathematics  
University of Santa Clara  
Santa Clara, CA 95053  
USA

**F. W. Gehring**

Department of Mathematics  
University of Michigan  
Ann Arbor, MI 48104  
USA

**C. C. Moore**

Department of Mathematics  
University of California  
Berkeley, CA 94720  
USA

---

**AMS Subject Classification (1980):**

**26 A 21, 28 A 05, 54 C 50, 54 E 50, 54 H 05, 26-01, 28-01, 54-01**

---

### **Library of Congress Cataloging in Publication Data**

**Oxtoby, John C.**

**Measure and category.**

**(Graduate texts in mathematics; 2)**

**Bibliography: p.**

**Includes index.**

**1. Measure theory. 2. Topological spaces. 3. Categories (Mathematics)**

**I. Title. II. Series. QA312.O9 1980 515.4'2 80-15770**

**All rights reserved.**

**No part of this book may be translated or reproduced in any form  
without written permission from Springer-Verlag.**

**© 1971, 1980 by Springer-Verlag New York Inc.**

**9 8 7 6 5 4 3 (third printing, 1987)**

**ISBN 0-387-90508-1 Springer-Verlag New York**

**ISBN 3-540-90508-1 Springer-Verlag Berlin Heidelberg**

**This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale  
in the Mainland China only and not for export therefrom.**

**图书在版编目 (CIP) 数据**

测度与范畴学 = Measure and Category: 第2版: 英文/  
(美) 奥凯斯屠拜 (Oxtoby, J. C.) 著. —北京: 世界图  
书出版公司北京公司, 2009. 4  
ISBN 978-7-5100-0439-1

I. 测… II. 奥… III. ①测度论—英文②范畴论—英文  
IV. 0174.12 0154.1

中国版本图书馆 CIP 数据核字 (2009) 第 048105 号

---

书 号: Measure and Category 2nd ed.

作 者: John C. Oxtoby

---

中 译 名: 测度和范畴学 第2版

责任编辑: 高蓉 刘慧

---

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河国英印务有限公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64021602, 010-64015659

电子信箱: kjb@wpcbj.com.cn

---

开 本: 24 开

印 张: 5

版 次: 2009 年 04 月

版权登记: 图字: 01-2009-0683

---

书 号: 978-7-5100-0439-1/O · 654

定 价: 18.00 元

---

世界图书出版公司北京公司已获得 Springer 授权在中国大陆独家重印发行

# Graduate Texts in Mathematics 2

## *Editorial Board*

F. W. Gehring P. R. Halmos (Managing Editor)

C. C. Moore

# Graduate Texts in Mathematics

- 1 TAKEUTI/ZARING. Introduction to Axiomatic Set Theory. 2nd ed.
- 2 OXToby. Measure and Category. 2nd ed.
- 3 SCHAEFFER. Topological Vector Spaces.
- 4 HILTON/STAMMBACH. A Course in Homological Algebra.
- 5 MACLANE. Categories for the Working Mathematician.
- 6 HUGHES/PIPER. Projective Planes.
- 7 SERRE. A Course in Arithmetic.
- 8 TAKEUTI/ZARING. Axiomatic Set Theory.
- 9 HUMPHREYS. Introduction to Lie Algebras and Representation Theory.
- 10 COHEN. A Course in Simple Homotopy Theory.
- 11 CONWAY. Functions of One Complex Variable. 2nd ed.
- 12 BEALS. Advanced Mathematical Analysis.
- 13 ANDERSON/FULLER. Rings and Categories of Modules.
- 14 GOLUBITSKY/GUILLEMIN. Stable Mappings and Their Singularities.
- 15 BERBERIAN. Lectures in Functional Analysis and Operator Theory.
- 16 WINTER. The Structure of Fields.
- 17 ROSENBLATT. Random Processes. 2nd ed.
- 18 HALMOS. Measure Theory.
- 19 HALMOS. A Hilbert Space Problem Book. 2nd ed., revised.
- 20 HUSEMOLLER. Fibre Bundles. 2nd ed.
- 21 HUMPHREYS. Linear Algebraic Groups.
- 22 BARNES/MACK. An Algebraic Introduction to Mathematical Logic.
- 23 GREUB. Linear Algebra. 4th ed.
- 24 HOLMES. Geometric Functional Analysis and its Applications.
- 25 HEWITT/STROMBERG. Real and Abstract Analysis.
- 26 MANES. Algebraic Theories.
- 27 KELLEY. General Topology.
- 28 ZARISKI/SAMUEL. Commutative Algebra. Vol. I.
- 29 ZARISKI/SAMUEL. Commutative Algebra. Vol. II.
- 30 JACOBSON. Lectures in Abstract Algebra I: Basic Concepts.
- 31 JACOBSON. Lectures in Abstract Algebra II: Linear Algebra.
- 32 JACOBSON. Lectures in Abstract Algebra III: Theory of Fields and Galois Theory.
- 33 HIRSCH. Differential Topology.
- 34 SPITZER. Principles of Random Walk. 2nd ed.
- 35 WERMER. Banach Algebras and Several Complex Variables. 2nd ed.
- 36 KELLEY/NAMIOKA et al. Linear Topological Spaces.
- 37 MONK. Mathematical Logic.
- 38 GRAUERT/FRITZSCHE. Several Complex Variables.
- 39 ARVESON. An Invitation to  $C^*$ -Algebras.
- 40 KEMENY/SNELL/KNAPP. Denumerable Markov Chains. 2nd ed.
- 41 APOSTOL. Modular Functions and Dirichlet Series in Number Theory.
- 42 SERRE. Linear Representations of Finite Groups.
- 43 GILLMAN/JERISON. Rings of Continuous Functions.

*(continued on back page)*

## **Preface to the Second Edition**

In this edition, a set of Supplementary Notes and Remarks has been added at the end, grouped according to chapter. Some of these call attention to subsequent developments, others add further explanation or additional remarks. Most of the remarks are accompanied by a briefly indicated proof, which is sometimes different from the one given in the reference cited. The list of references has been expanded to include many recent contributions, but it is still not intended to be exhaustive.

Bryn Mawr, April 1980

John C. Oxtoby

## **Preface to the First Edition**

This book has two main themes: the Baire category theorem as a method for proving existence, and the “duality” between measure and category. The category method is illustrated by a variety of typical applications, and the analogy between measure and category is explored in all of its ramifications. To this end, the elements of metric topology are reviewed and the principal properties of Lebesgue measure are derived. It turns out that Lebesgue integration is not essential for present purposes—the Riemann integral is sufficient. Concepts of general measure theory and topology are introduced, but not just for the sake of generality. Needless to say, the term “category” refers always to Baire category; it has nothing to do with the term as it is used in homological algebra.

A knowledge of calculus is presupposed, and some familiarity with the algebra of sets. The questions discussed are ones that lend themselves naturally to set-theoretical formulation. The book is intended as an introduction to this kind of analysis. It could be used to supplement a standard course in real analysis, as the basis for a seminar, or for independent study. It is primarily expository, but a few refinements of known results are included, notably Theorem 15.6 and Proposition 20.4. The references are not intended to be complete. Frequently a secondary source is cited where additional references may be found.

The book is a revised and expanded version of notes originally prepared for a course of lectures given at Haverford College during the spring of 1957 under the auspices of the William Pyle Philips Fund. These, in turn, were based on the Earle Raymond Hedrick Lectures presented at the Summer Meeting of the Mathematical Association of America at Seattle, Washington, in August, 1956.

Bryn Mawr, April 1971

John C. Oxtoby



**Graduate Texts in Mathematics** (*continued from page ii*)

- 44 KENDIG. Elementary Algebraic Geometry.
- 45 LOÈVE. Probability Theory I. 4th ed.
- 46 LOÈVE. Probability Theory II. 4th ed.
- 47 MOISE. Geometric Topology in Dimensions 2 and 3.
- 48 SACHS/WU. General Relativity for Mathematicians.
- 49 GRUENBERG/WEIR. Linear Geometry. 2nd ed.
- 50 EDWARDS. Fermat's Last Theorem.
- 51 KLINGENBERG. A Course in Differential Geometry.
- 52 HARTSHORNE. Algebraic Geometry.
- 53 MANIN. A Course in Mathematical Logic.
- 54 GRAVER/WATKINS. Combinatorics with Emphasis on the Theory of Graphs.
- 55 BROWN/PEARCY. Introduction to Operator Theory I: Elements of Functional Analysis.
- 56 MASSEY. Algebraic Topology: An Introduction.
- 57 CROWELL/FOX. Introduction to Knot Theory.
- 58 KOBLITZ.  $p$ -adic Numbers,  $p$ -adic Analysis, and Zeta-Functions.
- 59 LANG. Cyclotomic Fields.
- 60 ARNOLD. Mathematical Methods in Classical Mechanics.
- 61 WHITEHEAD. Elements of Homotopy Theory.
- 62 KARGAPOLOV/MERZLIAKOV. Fundamentals of the Theory of Groups.
- 63 BOLLOBÁS. Graph Theory.
- 64 EDWARDS. Fourier Series. Vol. I. 2nd ed.
- 65 WELLS. Differential Analysis on Complex Manifolds. 2nd ed.
- 66 WATERHOUSE. Introduction to Affine Group Schemes.
- 67 SERRE. Local Fields.
- 68 WEIDMANN. Linear Operators in Hilbert Spaces.
- 69 LANG. Cyclotomic Fields II.
- 70 MASSEY. Singular Homology Theory.
- 71 FARKAS/KRA. Riemann Surfaces.
- 72 STILLWELL. Classical Topology and Combinatorial Group Theory.
- 73 HUNGERFORD. Algebra.
- 74 DAVENPORT. Multiplicative Number Theory. 2nd ed.
- 75 HOCHSCHILD. Basic Theory of Algebraic Groups and Lie Algebras.
- 76 IITAKA. Algebraic Geometry.
- 77 HECKE. Lectures on the Theory of Algebraic Numbers.
- 78 BURRIS/SANKAPPANAVAR. A Course in Universal Algebra.
- 79 WALTERS. An Introduction to Ergodic Theory.
- 80 ROBINSON. A Course in the Theory of Groups.
- 81 FORSTER. Lectures on Riemann Surfaces.
- 82 BOTT/TU. Differential Forms in Algebraic Topology.
- 83 WASHINGTON. Introduction to Cyclotomic Fields.
- 84 IRELAND/ROSEN. A Classical Introduction Modern Number Theory.
- 85 EDWARDS. Fourier Series. Vol. II. 2nd ed.
- 86 VAN LINT. Introduction to Coding Theory.

- 87 BROWN. Cohomology of Groups.
- 88 PIERCE. Associative Algebras.
- 89 LANG. Introduction to Algebraic and Abelian Functions. 2nd ed.
- 90 BRØNDSTED. An Introduction to Convex Polytopes.
- 91 BEARDON. On the Geometry of Discrete Groups.
- 92 DIESTEL. Sequences and Series in Banach Spaces.
- 93 DUBROVIN/FOMENKO/NOVIKOV. Modern Geometry—Methods and Applications  
Vol. I.
- 94 WARNER. Foundations of Differentiable Manifolds and Lie Groups.
- 95 SHIRYAYEV. Probability, Statistics, and Random Processes.
- 96 ZEIDLER. Nonlinear Functional Analysis and Its Applications: Fixed Points  
Theorem.

# Contents

1. Measure and Category on the Line . . . . .	1
Countable sets, sets of first category, nullsets, the theorems of Cantor, Baire, and Borel	
2. Liouville Numbers . . . . .	6
Algebraic and transcendental numbers, measure and category of the set of Liouville numbers	
3. Lebesgue Measure in $r$ -Space. . . . .	10
Definitions and principal properties, measurable sets, the Lebesgue density theorem	
4. The Property of Baire . . . . .	19
Its analogy to measurability, properties of regular open sets	
5. Non-Measurable Sets . . . . .	22
Vitali sets, Bernstein sets, Ulam's theorem, inaccessible cardinals, the continuum hypothesis	
6. The Banach-Mazur Game . . . . .	27
Winning strategies, category and local category, indeterminate games	
7. Functions of First Class . . . . .	31
Oscillation, the limit of a sequence of continuous functions, Riemann integrability	
8. The Theorems of Lusin and Egoroff . . . . .	36
Continuity of measurable functions and of functions having the property of Baire, uniform convergence on subsets	
9. Metric and Topological Spaces . . . . .	39
Definitions, complete and topologically complete spaces, the Baire category theorem	
10. Examples of Metric Spaces. . . . .	42
Uniform and integral metrics in the space of continuous functions, integrable functions, pseudo-metric spaces, the space of measurable sets	
11. Nowhere Differentiable Functions . . . . .	45
Banach's application of the category method	

12. The Theorem of Alexandroff . . . . .	47
Remetrization of a $G_\delta$ subset, topologically complete subspaces	
13. Transforming Linear Sets into Nullsets . . . . .	49
The space of automorphisms of an interval, effect of monotone substitution on Riemann integrability, nullsets equivalent to sets of first category	
14. Fubini's Theorem . . . . .	52
Measurability and measure of sections of plane measurable sets	
15. The Kuratowski-Ulam Theorem . . . . .	56
Sections of plane sets having the property of Baire, product sets, reducibility to Fubini's theorem by means of a product transformation	
16. The Banach Category Theorem . . . . .	62
Open sets of first category or measure zero, Montgomery's lemma, the theorems of Marczewski and Sikorski, cardinals of measure zero, decomposition into a nullset and a set of first category	
17. The Poincaré Recurrence Theorem . . . . .	65
Measure and category of the set of points recurrent under a nondissipative transformation, application to dynamical systems	
18. Transitive Transformations . . . . .	70
Existence of transitive automorphisms of the square, the category method	
19. The Sierpinski-Erdős Duality Theorem . . . . .	74
Similarities between the classes of sets of measure zero and of first category, the principle of duality	
20. Examples of Duality . . . . .	78
Properties of Lusin sets and their duals, sets almost invariant under transformations that preserve nullsets or category	
21. The Extended Principle of Duality . . . . .	82
A counter example, product measures and product spaces, the zero-one law and its category analogue	
22. Category Measure Spaces . . . . .	86
Spaces in which measure and category agree, topologies generated by lower densities, the Lebesgue density topology	
Supplementary Notes and Remarks . . . . .	93
References . . . . .	101
Supplementary References . . . . .	102
Index . . . . .	105

## 1. Measure and Category on the Line

The notions of measure and category are based on that of countability. Cantor's theorem, which says that no interval of real numbers is countable, provides a natural starting point for the study of both measure and category. Let us recall that a set is called *denumerable* if its elements can be put in one-to-one correspondence with the natural numbers  $1, 2, \dots$ . A *countable* set is one that is either finite or denumerable. The set of rational numbers is denumerable, because for each positive integer  $k$  there are only a finite number ( $\leq 2k - 1$ ) of rational numbers  $p/q$  in reduced form ( $q > 0$ ,  $p$  and  $q$  relatively prime) for which  $|p| + q = k$ . By numbering those for which  $k = 1$ , then those for which  $k = 2$ , and so on, we obtain a sequence in which each rational number appears once and only once. Cantor's theorem reads as follows.

**Theorem 1.1 (Cantor).** *For any sequence  $\{a_n\}$  of real numbers and for any interval  $I$  there exists a point  $p$  in  $I$  such that  $p \neq a_n$  for every  $n$ .*

One proof runs as follows. Let  $I_1$  be a closed subinterval of  $I$  such that  $a_1 \notin I_1$ . Let  $I_2$  be a closed subinterval of  $I_1$  such that  $a_2 \notin I_2$ . Proceeding inductively, let  $I_n$  be a closed subinterval of  $I_{n-1}$  such that  $a_n \notin I_n$ . The nested sequence of closed intervals  $I_n$  has a non-empty intersection. If  $p \in \bigcap I_n$ , then  $p \in I$  and  $p \neq a_n$  for every  $n$ .

This proof involves infinitely many unspecified choices. To avoid this objection the intervals must be chosen according to some definite rule. One such rule is this: divide  $I_{n-1}$  into three subintervals of equal length and take for  $I_n$  the first one of these that does not contain  $a_n$ . If we take  $I_0$  to be the closed interval concentric with  $I$  and half as long, say, then all the choices are specified, and we have a well defined function of  $(I, a_1, a_2, \dots)$  whose value is a point of  $I$  different from all the  $a_n$ .

The fact that no interval is countable is an immediate corollary of Cantor's theorem.

With only a few changes, the above proof becomes a proof of the Baire category theorem for the line. Before we can formulate this theorem we need some definitions. A set  $A$  is *dense in the interval  $I$*  if  $A$  has a non-empty intersection with every subinterval of  $I$ ; it is called *dense* if it is

dense in the line  $R$ . A set  $A$  is *nowhere dense* if it is dense in no interval, that is, if every interval has a subinterval contained in the complement of  $A$ . A nowhere dense set may be characterized as one that is "full of holes." The definition can be stated in two other useful ways:  $A$  is nowhere dense if and only if its complement  $A'$  contains a dense open set, and if and only if  $\bar{A}$  (or  $A^-$ , the closure of  $A$ ) has no interior points. The class of nowhere dense sets is closed under certain operations, namely

**Theorem 1.2.** *Any subset of a nowhere dense set is nowhere dense. The union of two (or any finite number) of nowhere dense sets is nowhere dense. The closure of a nowhere dense set is nowhere dense.*

*Proof.* The first statement is obvious. To prove the second, note that if  $A_1$  and  $A_2$  are nowhere dense, then for each interval  $I$  there is an interval  $I_1 \subset I - A_1$  and an interval  $I_2 \subset I_1 - A_2$ . Hence  $I_2 \subset I - (A_1 \cup A_2)$ . This shows that  $A_1 \cup A_2$  is nowhere dense. Finally, any open interval contained in  $A'$  is also contained in  $A'^-$ .  $\square$

A denumerable union of nowhere dense sets is not in general nowhere dense, it may even be dense. For instance, the set of rational numbers is dense, but it is also a denumerable union of singletons (sets having just one element), and singletons are nowhere dense in  $R$ .

A set is said to be of *first category* if it can be represented as a countable union of nowhere dense sets. A subset of  $R$  that cannot be so represented is said to be of *second category*. These definitions were formulated in 1899 by R. Baire [18, p. 48], to whom the following theorem is due.

**Theorem 1.3 (Baire).** *The complement of any set of first category on the line is dense. No interval in  $R$  is of first category. The intersection of any sequence of dense open sets is dense.*

*Proof.* The three statements are essentially equivalent. To prove the first, let  $A = \bigcup A_n$  be a representation of  $A$  as a countable union of nowhere dense sets. For any interval  $I$ , let  $I_1$  be a closed subinterval of  $I - A_1$ . Let  $I_2$  be a closed subinterval of  $I_1 - A_2$ , and so on. Then  $\bigcap I_n$  is a non-empty subset of  $I - A$ , hence  $A'$  is dense. To specify all the choices in advance, it suffices to arrange the (denumerable) class of closed intervals with rational endpoints into a sequence, take  $I_0 = I$ , and for  $n > 0$  take  $I_n$  to be the first term of the sequence that is contained in  $I_{n-1} - A_n$ .

The second statement is an immediate corollary of the first. The third statement follows from the first by complementation.  $\square$

Evidently Baire's theorem implies Cantor's. Its proof is similar, although a different rule for choosing  $I_n$  was needed.

**Theorem 1.4.** *Any subset of a set of first category is of first category. The union of any countable family of first category sets is of first category.*

It is obvious that the class of first category sets has these closure properties. However, the closure of a set of first category is not in general of first category. In fact, the closure of a linear set  $A$  is of first category if and only if  $A$  is nowhere dense.

A class of sets that contains countable unions and arbitrary subsets of its members is called a  $\sigma$ -ideal. The class of sets of first category and the class of countable sets are two examples of  $\sigma$ -ideals of subsets of the line. Another example is the class of nullsets, which we shall now define.

The length of any interval  $I$  is denoted by  $|I|$ . A set  $A \subset R$  is called a *nullset* (or a set of *measure zero*) if for each  $\varepsilon > 0$  there exists a sequence of intervals  $I_n$  such that  $A \subset \bigcup I_n$  and  $\sum |I_n| < \varepsilon$ .

It is obvious that singletons are nullsets and that any subset of a nullset is a nullset. Any countable union of nullsets is also a nullset. For suppose  $A_i$  is a nullset for  $i = 1, 2, \dots$ . Then for each  $i$  there is a sequence of intervals  $I_{ij}$  ( $j = 1, 2, \dots$ ) such that  $A_i \subset \bigcup_j I_{ij}$  and  $\sum_j |I_{ij}| < \varepsilon/2^i$ . The set of all the intervals  $I_{ij}$  covers  $A$ , and  $\sum_{i,j} |I_{ij}| < \varepsilon$ , hence  $A$  is a nullset. This shows that the class of nullsets is a  $\sigma$ -ideal. Like the class of sets of first category, it includes all countable sets.

**Theorem 1.5 (Borel).** *If a finite or infinite sequence of intervals  $I_n$  covers an interval  $I$ , then  $\sum |I_n| \geq |I|$ .*

*Proof.* Assume first that  $I = [a, b]$  is closed and that all of the intervals  $I_n$  are open. Let  $(a_1, b_1)$  be the first interval that contains  $a$ . If  $b_1 \leq b$ , let  $(a_2, b_2)$  be the first interval of the sequence that contains  $b_1$ . If  $b_{n-1} \leq b$ , let  $(a_n, b_n)$  be the first interval that contains  $b_{n-1}$ . This procedure must terminate with some  $b_N > b$ . Otherwise the increasing sequence  $\{b_n\}$  would converge to a limit  $x \leq b$ , and  $x$  would belong to  $I_k$  for some  $k$ . All but a finite number of the intervals  $(a_n, b_n)$  would have to precede  $I_k$  in the given sequence, namely, all those for which  $b_{n-1} \in I_k$ . This is impossible, since no two of these intervals are equal. (Incidentally, this reasoning reproduces Borel's own proof of the "Heine-Borel theorem" [5, p. 228].) We have

$$b - a < b_N - a_1 = \sum_{i=2}^N (b_i - b_{i-1}) + b_1 - a_1 \leq \sum_{i=1}^N (b_i - a_i),$$

and so the theorem is true in this case.

In the general case, for any  $\alpha > 1$  let  $J$  be a closed subinterval of  $I$  with  $|J| = |I|/\alpha$ , and let  $J_n$  be an open interval containing  $I_n$  with  $|J_n| = \alpha |I_n|$ . Then  $J$  is covered by the sequence  $\{J_n\}$ . We have already shown that

$\sum |J_n| \geq |J|$ . Hence  $\alpha \sum |I_n| = \sum |J_n| \geq |J| = |I|/\alpha$ . Letting  $\alpha \rightarrow 1$  we obtain the desired conclusion.  $\square$

This theorem implies that no interval is a nullset; it therefore provides still another proof of Cantor's theorem.

Every countable set is of first category and of measure zero. Some uncountable sets also belong to both classes. The simplest example is the *Cantor set*  $C$ , which consists of all numbers in the interval  $[0, 1]$  that admit a ternary development in which the digit 1 does not appear. It can be constructed by deleting the open middle third of the interval  $[0, 1]$ , then deleting the open middle thirds of each of the intervals  $[0, 1/3]$  and  $[2/3, 1]$ , and so on. If  $F_n$  denotes the union of the  $2^n$  closed intervals of length  $1/3^n$  which remain at the  $n$ -th stage, then  $C = \bigcap F_n$ .  $C$  is closed, since it is an intersection of closed sets.  $C$  is nowhere dense, since  $F_n$  (and therefore  $C$ ) contains no interval of length greater than  $1/3^n$ . The sum of the lengths of the intervals that compose  $F_n$  is  $(2/3)^n$ , which is less than  $\varepsilon$  if  $n$  is taken sufficiently large. Hence  $C$  is a nullset. Finally, each number  $x$  in  $(0, 1]$  has a unique non-terminating binary development  $x = .x_1x_2x_3\ldots$ . If  $y_i = 2x_i$ , then  $.y_1y_2y_3\ldots$  is the ternary development with  $y_i \neq 1$  of some point  $y$  of  $C$ . This correspondence between  $x$  and  $y$ , extended by mapping 0 onto itself, defines a one-to-one map of  $[0, 1]$  onto a (proper) subset of  $C$ . It follows that  $C$  is uncountable; it has cardinality  $c$  (the power of the continuum).

The sets of measure zero and the sets of first category constitute two  $\sigma$ -ideals, each of which properly contains the class of countable sets. Their properties suggest that a set belonging to either class is "small" in one sense or another. A nowhere dense set is small in the intuitive geometric sense of being perforated with holes, and a set of first category can be "approximated" by such a set. A set of first category may or may not have any holes, but it always has a dense set of gaps. No interval can be represented as the union of a sequence of such sets. On the other hand, a nullset is small in the metric sense that it can be covered by a sequence of intervals of arbitrarily small total length. If a point is chosen at random in an interval in such a way that the probability of its belonging to any subinterval  $J$  is proportional to  $|J|$ , then the probability of its belonging to any given nullset is zero. It is natural to ask whether these notions of smallness are related. Does either class include the other? That neither class does, and that in some cases the two notions may be diametrically opposed, is shown by the following

**Theorem 1.6.** *The line can be decomposed into two complementary sets  $A$  and  $B$  such that  $A$  is of first category and  $B$  is of measure zero.*

*Proof.* Let  $a_1, a_2, \ldots$  be an enumeration of the set of rational numbers (or of any countable dense subset of the line). Let  $I_{ij}$  be the open interval



with center  $a_i$  and length  $1/2^{i+j}$ . Let  $G_j = \bigcup_{i=1}^{\infty} I_{ij}$  ( $j = 1, 2, \dots$ ) and  $B = \bigcap_{j=1}^{\infty} G_j$ . For any  $\varepsilon > 0$  we can choose  $j$  so that  $1/2^j < \varepsilon$ . Then  $B \subset \bigcup_i I_{ij}$  and  $\sum_i |I_{ij}| = \sum_i 1/2^{i+j} = 1/2^j < \varepsilon$ . Hence  $B$  is a nullset. On the other hand,  $G_j$  is a dense open subset of  $R$ , since it is the union of a sequence of open intervals and it includes all rational points. Therefore its complement  $G_j'$  is nowhere dense, and  $A = B' = \bigcup_j G_j'$  is of first category.  $\square$

**Corollary 1.7.** *Every subset of the line can be represented as the union of a nullset and a set of first category.*

There is of course nothing paradoxical in the fact that a set that is small in one sense may be large in some other sense.