

Oleg T. Izhboldin    Bruno Kahn  
Nikita A. Karpenko    Alexander Vishik

# Geometric Methods in the Algebraic Theory of Quadratic Forms

1835

Summer School, Lens, 2000

Editor: Jean-Pierre Tignol



Springer

Oleg T. Izhboldin   Bruno Kahn  
Nikita A. Karpenko   Alexander Vishik

# Geometric Methods in the Algebraic Theory of Quadratic Forms

Summer School, Lens, 2000

Editor:

Jean-Pierre Tignol



Springer

## Authors

Oleg T. Izhboldin  
(Deceased April 17, 2000)

Bruno Kahn  
Institut de Mathématiques de Jussieu  
175-179 rue du Chevaleret  
75013 Paris, France  
*e-mail: kahn@math.jussieu.fr*

Alexander Vishik  
Institute for Information  
Transmission Problems  
Russian Academy of Sciences  
Bolshoj Karetnyj Pereulok, Dom 19  
101447 Moscow, Russia  
*e-mail: vishik@mccme.ru*

Nikita A. Karpenko  
Université d'Artois  
Rue Jean Souvraz SP 18  
62307 Lens, France  
*e-mail: karpenko@euler.univ-artois.fr*

## Editor

Jean-Pierre Tignol  
Institut de Mathématique Pure et Appliquée  
Université catholique de Louvain  
Chemin du Cyclotron 2  
1348 Louvain-la-Neuve, Belgium  
*e-mail: tignol@math.ucl.ac.be*

## Cataloging-in-Publication Data applied for

Bibliographic information published by Die Deutsche Bibliothek

Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie;  
detailed bibliographic data is available in the Internet at <http://dnb.ddb.de>

Mathematics Subject Classification (2000): 11E81, 14C15, 14F42

ISSN 0075-8434

ISBN 3-540-20728-7 Springer-Verlag Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

Springer-Verlag is a part of Springer Science+Business Media  
[springeronline.com](http://springeronline.com)

© Springer-Verlag Berlin Heidelberg 2004  
Printed in Germany

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Typesetting: Camera-ready T<sub>E</sub>X output by the author

SPIN: 10976173      41/3142/du - 543210 - Printed on acid-free paper

In memory of Oleg Tomovich Izhboldin (1963–2000)

---

## Preface

The geometric approach to the algebraic theory of quadratic forms is the study of projective quadrics over arbitrary fields. Function fields of quadrics were a basic ingredient already in the proof of the Arason–Pfister Hauptsatz of 1971 (or even in Pfister’s 1965 construction of fields with prescribed level); they are central in the investigation of deep properties of quadratic forms, such as their splitting pattern, but also in the construction of fields which exhibit particular properties, such as a given  $u$ -invariant. Recently, finer geometric tools have been brought to bear on problems from the algebraic theory of quadratic forms: results on Chow groups of quadrics led to an efficient use of motives, and ultimately to Voevodsky’s proof of the Milnor conjecture.

The goal of the June 2000 summer school at Université d’Artois in Lens (France), organized locally by J. Burési, N. Karpenko and P. Mammone, was to survey three aspects of the algebraic theory of quadratic forms where geometric methods had led to spectacular advances. Bruno Kahn was invited to talk on the unramified cohomology of quadrics, Alexander Vishik on motives of quadrics, and Oleg Izhboldin on his construction of fields whose  $u$ -invariant is 9. However, Izhboldin passed away unexpectedly on April 17, 2000. His work was surveyed by Karpenko, who had collaborated with Izhboldin on several papers.

The closely related texts collected in this volume were written from somewhat different perspectives. The reader will find below:

1. The notes from the lectures of B. Kahn [K], A. Vishik [V] and N. Karpenko [K1] prepared and updated by the authors. Additional material has been included, in particular in Vishik’s notes.
2. Two papers left unfinished by O. Izhboldin, and edited by N. Karpenko. The first paper [I1] was essentially complete and formed the basis for the first part of Karpenko’s lectures. The second [I2] is only a sketch, listing properties and examples that Izhboldin intended to develop in subsequent work.

3. A paper by N. Karpenko [K2] which provides complete proofs for the statements that Izhboldin listed in [I2].

To give a more precise overview, we introduce some notation. Let  $F$  be an arbitrary field of characteristic different from 2. To every quadratic form<sup>1</sup>  $q$  in at least two variables over  $F$  corresponds the projective quadric  $Q$  with equation  $q = 0$  (which has an  $F$ -rational point if and only if  $q$  is isotropic). The quadric  $Q$  is a smooth variety if  $q$  is nonsingular (which we always assume in the sequel); its dimension is  $\dim Q = \dim q - 2$ , and it is irreducible if  $q$  is not the hyperbolic plane  $\mathbb{H}$ . We may then consider its function field  $F(Q)$ , which is also referred to as the *function field of  $q$*  and denoted  $F(q)$ .

The field extension  $F(Q)/F$  is of particular interest. Much insight into quadratic forms could be obtained if we knew which quadratic forms over  $F$  become isotropic over  $F(Q)$ . This question can be readily rephrased into geometric terms: a quadratic form  $q'$  over  $F$  becomes isotropic over  $F(q)$  if and only if there is a rational map  $Q \dashrightarrow Q'$  between the corresponding quadrics. If there are rational maps in both directions  $Q \dashrightarrow Q'$  and  $Q' \dashrightarrow Q$ , the quadrics are stably birationally equivalent, and the quadratic forms  $q$  and  $q'$  are also called *stably birationally equivalent*. By the preceding observation, this relation, denoted  $q \stackrel{st}{\sim} q'$ , holds if and only if the forms  $q_{F(q')}$  and  $q'_{F(q)}$  are both isotropic.

A very useful geometric construction is to view the quadric  $Q$  as an object in a category where the maps are given by Chow correspondences. We thus get the (Chow-) motive  $M(Q)$  of the quadric, whose structure carries a lot of information on the form  $q$ . For example, Vishik has shown<sup>2</sup> that the motives  $M(Q)$ ,  $M(Q')$  associated with quadratic forms  $q$ ,  $q'$  are isomorphic if and only if every field extension  $E$  of  $F$  produces the same amount of splitting in  $q$  and  $q'$ , i.e., the quadratic forms  $q_E$  and  $q'_E$  have the same *Witt index*, a notion which is spelled out next.

Recall from [Sch, Corollary 5.11 of Chap. 1] that every quadratic form  $q$  has a (Witt) decomposition into an orthogonal sum of an anisotropic quadratic form  $q_{\text{an}}$ , called an *anisotropic kernel* of  $q$ , and a certain number of hyperbolic planes  $\mathbb{H}$ ,

$$q \simeq q_{\text{an}} \perp \underbrace{\mathbb{H} \perp \dots \perp \mathbb{H}}_{i_W(q)}.$$

The number  $i_W(q)$  of hyperbolic planes in this decomposition (which is unique up to isomorphism) is called the *Witt index* of  $q$ . Even if  $q$  is anisotropic (i.e.,  $i_W(q) = 0$ ), it obviously becomes isotropic over  $F(q)$ , and we have a Witt decomposition over  $F(q)$ ,

$$q_{F(q)} \simeq q_1 \perp \mathbb{H} \perp \dots \perp \mathbb{H}$$

<sup>1</sup> With the usual abuse of terminology, a quadratic form is sometimes viewed as a quadratic polynomial, sometimes as a quadratic map on a vector space or a quadratic space.

<sup>2</sup> See [I2, Sect. 1].

where  $q_1$  is an anisotropic form over  $F(q)$ . Letting  $F_1 = F(q)$ , we may iterate this construction. The process terminates in a finite number of steps since  $\dim q > \dim q_1 > \dots$ . We thus obtain the *generic splitting tower* of  $q$ , first constructed by M. Knebusch [Kn],

$$F \subset F_1 \subset \dots \subset F_h.$$

Clearly,  $0 < i_W(q_{F_1}) < i_W(q_{F_2}) < \dots < i_W(q_{F_h})$ . It turns out that for any field extension  $E/F$ , the Witt index  $i_W(q_E)$  is equal to one of the indices  $i_W(q_{F_j})$ . The *splitting pattern* of  $q$  is the set

$$\{i_W(q_E) \mid E \text{ a field extension of } F\} = \{i_W(q_{F_1}), \dots, i_W(q_{F_h})\}.$$

Variants of this notion appear in [V] and [I1]: Vishik calls (*incremental*) *splitting pattern*<sup>3</sup> of  $q$  the sequence

$$\mathbf{i}(q) = (i_1(q), \dots, i_h(q))$$

defined by  $i_1(q) = i_W(q_{F_1})$  and  $i_j(q) = i_W(q_{F_j}) - i_W(q_{F_{j-1}})$  for  $j > 1$ . The integer  $i_j(q)$  indeed measures the Witt index increment resulting from the field extension  $F_j/F_{j-1}$ ; it is called a *higher Witt index* of  $q$ . On the other hand, Izhboldin concentrates on the dimension of the anisotropic kernels and sets

$$\text{Dim}(q) = \{\dim(q_E)_{\text{an}} \mid E \text{ a field extension of } F\}.$$

By counting dimensions in the Witt decomposition of  $q_E$ , we obtain

$$\dim q = \dim q_E = \dim(q_E)_{\text{an}} + 2i_W(q_E),$$

hence the set  $\text{Dim}(q)$  and the splitting pattern of  $q$  carry the same information.

Vishik's contribution [V] to this volume is intended as a general introduction to the state-of-the-art in the theory of motives of quadrics. After setting up the basic principles, he proves the main structure theorems on motives of quadrics. The study of direct sum decompositions of these motives is a powerful tool for investigating the dimensions of anisotropic forms in the powers of the fundamental ideal of the Witt ring, the stable equivalence of quadrics and splitting patterns of quadratic forms. This last application is particularly developed in the last section of [V], where all the possible splitting patterns of odd-dimensional forms of dimension at most 21 and of even-dimensional forms of dimension at most 12 are determined.

The papers of Karpenko [K1, K2] and Izhboldin [I1, I2] are closely intertwined. They also rely less on motives and more on elementary arguments. As mentioned above, [K1] is an exposition of Izhboldin's results in [I1] and on

<sup>3</sup> No confusion should arise since Vishik's splitting patterns are sequences, whereas the "usual" splitting patterns are sets.

the  $u$ -invariant. Recall from [Sch, Sect. 16 of Chap. 2] that the  $u$ -invariant of a field  $F$  is

$$u(F) = \sup\{\dim q \mid q \text{ anisotropic quadratic form over } F\}.$$

Quadratically closed fields have  $u$ -invariant 1, but no other field with odd  $u$ -invariant was known before Izhboldin's construction of a field with  $u$ -invariant 9. In the second part of [K1], Karpenko discusses the strategy of this construction and provides alternative proofs for the main results on which it is based. In the first part, he gives a simple proof of a theorem of Izhboldin on the first Witt index  $i_1(q)$  of quadratic forms of dimension  $2^n + 3$ . Izhboldin's original proof is given in [I1], while [I2] classifies the pairs of quadratic forms of dimension at most 9 which are stably equivalent and lists without proofs assorted isotropy criteria for quadratic forms over function fields of quadrics. The proofs of Izhboldin's claims are given in [K2], which also contains an extensive discussion of correspondences on odd-dimensional quadrics.

In [K], Kahn studies the field extension  $F(Q)/F$  from a different angle. The induced scalar extension map in Galois cohomology with coefficients  $\mu_2 = \{\pm 1\}$ , called the *restriction map*

$$\text{Res}: H^n(F, \mu_2) \rightarrow H^n(F(Q), \mu_2)$$

is a typical case of the maps he considers. For every closed point  $x$  of  $Q$  of codimension 1, the image of this map lies in the kernel of the residue map

$$\partial_x: H^n(F(Q), \mu_2) \rightarrow H^{n-1}(F(x), \mu_2).$$

Therefore, we may restrict the target of Res to the *unramified cohomology group*

$$H_{\text{nr}}^n(F(Q), \mu_2) = \bigcap_{x \in Q^{(1)}} \text{Ker } \partial_x.$$

The kernel and cokernel of the restriction map  $H^n(F, \mu_2) \rightarrow H_{\text{nr}}^n(F(Q), \mu_2)$  were studied by Kahn–Rost–Sujatha [KRS] and Kahn–Sujatha [KS1, KS2] for  $n \leq 4$ . In his contribution to this volume, Kahn develops a vast generalization which applies to various cohomology theories besides Galois cohomology with  $\mu_2$  coefficients, and to arbitrary smooth projective varieties besides quadrics. If  $X$  is a smooth projective variety which is also geometrically cellular, there are two spectral sequences converging to the motivic cohomology of  $X$ . Results on the restriction map are obtained by comparing these two sequences, since one of them contains the unramified cohomology of  $X$  in its  $E_2$ -term. The unramified cohomology of quadrics occurs as a crucial ingredient in the other papers collected here, see [K1, Sect. 2.3], [K2, Lemma 7.5], [V, Lemmas 6.14 and 7.12].

The untimely death of Oleg Izhboldin was felt as a great loss by all the contributors to this volume, who decided to dedicate it to his memory. A



tribute to his work, written by his former thesis advisor Alexandr Merkurjev, and posted on the web site [www-math.univ-mlv.fr/~abakumov/oleg/](http://www-math.univ-mlv.fr/~abakumov/oleg/), is included as an appendix. We are grateful to A. Merkurjev and E. Abakumov for the permission to reproduce it.

## References

- [K] Kahn, B.: Cohomologie non ramifiée des quadriques. This volume.
- [KRS] Kahn, B., Rost, M., Sujatha, R.: Unramified cohomology of quadrics, I. Amer. J. Math., **120**, 841–891 (1998)
- [KS1] Kahn, B., Sujatha, R.: Unramified cohomology of quadrics, II. Duke Math. J. **106**, 449–484 (2001)
- [KS2] Kahn, B., Sujatha, R.: Motivic cohomology and unramified cohomology of quadrics. J. Eur. Math. Soc. **2**, 145–177 (2000)
- [K1] Karpenko, N.A.: Motives and Chow groups of quadrics with application to the  $u$ -invariant (after Oleg Izhboldin). This volume.
- [K2] Karpenko, N.A.: Izhboldin’s results on stably birational equivalence of quadrics. This volume.
- [Kn] Knebusch, M: Generic splitting of quadratic forms, I. Proc. London Math. Soc., **33**, 65–93 (1976)
- [I1] Izhboldin, O.T.: Virtual Pfister neighbors and first Witt index. This volume.
- [I2] Izhboldin, O.T.: Some new results concerning isotropy of low-dimensional forms. (List of examples and results (without proofs)). This volume.
- [Sch] Scharlau, W.: Quadratic and Hermitian Forms. Springer, Berlin Heidelberg New York Tokyo (1985)
- [V] Vishik, A.: Motives of quadrics with applications to the theory of quadratic forms. This volume.

---

# Contents

**Cohomologie non ramifiée des quadriques**

*Bruno Kahn*..... 1

1   Partie non ramifiée d’une théorie cohomologique..... 2

2   Pureté ; complexes de Cousin et complexes de Gersten ..... 4

3   Conjecture de Gersten ..... 5

4   Exemples de bonnes théories cohomologiques ..... 7

5   Cohomologie non ramifiée finie et divisible..... 8

6   Suite spectrale des poids ..... 9

7   Poids 0, 1, 2 ..... 11

8   Poids 3 ..... 15

9   Exemple : norme réduite ..... 17

10   Exemple : quadriques ..... 19

Références ..... 21

**Motives of Quadrics with Applications to the Theory of Quadratic Forms**

*Alexander Vishik*..... 25

1   Grothendieck Category of Chow Motives ..... 26

2   The Motive and the Chow Groups of a Hyperbolic Quadric ..... 28

3   General Theorems ..... 31

4   Indecomposable Direct Summands in the Motives of Quadrics..... 36

5   Proofs ..... 43

6   Some Applications..... 61

7   Splitting Patterns of Small-dimensional Forms ..... 71

References ..... 100

**Motives and Chow Groups of Quadrics with Application to the  $u$ -invariant (after Oleg Izhboldin)**

*Nikita A. Karpenko* ..... 103

1   Virtual Pfister Neighbors and First Witt Index ..... 104

2    $u$ -invariant 9..... 111

References ..... 127

**Virtual Pfister Neighbors and First Witt Index**

*Oleg T. Izhboldin*..... 131

1 Generic Principles ..... 132

2 Maximal Splitting ..... 133

3 Basic Construction ..... 133

4 Stable Equivalence of Quadratic Forms ..... 135

5 The Invariant  $d(\phi)$  ..... 137

References ..... 142

**Some New Results Concerning Isotropy of Low-dimensional Forms**

*Oleg T. Izhboldin*..... 143

1 Stable Equivalence ..... 145

2 Stable Equivalence of 7- and 8-dimensional Forms ..... 145

3 Isotropy of 9-dimensional Forms over Function Fields of Quadrics .. 146

4 Isotropy of Some 10- and 12-dimensional Forms ..... 149

References ..... 149

**Izhboldin’s Results on Stably Birational Equivalence of Quadrics**

*Nikita A. Karpenko* ..... 151

1 Notation and Results We Are Using ..... 153

2 Correspondences on Odd-dimensional Quadrics..... 157

3 Forms of Dimension 7 ..... 163

4 Forms of Dimension 8 ..... 167

5 Forms of Dimension 9 ..... 168

6 Examples of Non-similar Stably Equivalent Forms of Dimension 9 .. 175

7 Other Related Results ..... 177

References ..... 182

**Appendix: My Recollections About Oleg Izhboldin**

*Alexander S. Merkurjev*..... 185

**Index** ..... 189

---

# Cohomologie non ramifiée des quadriques

Bruno Kahn

Institut de Mathématiques de Jussieu  
175–179 rue du Chevaleret  
75013 Paris, France  
kahn@math.jussieu.fr

## Introduction

Le but de ce texte est de donner un survol de techniques permettant le calcul de la cohomologie non ramifiée de certaines variétés projectives homogènes en poids  $\leq 3$ . Bien que la cohomologie non ramifiée soit un invariant birationnel des variétés propres et lisses (cf. théorème 3.3), ces techniques exigent la donnée d'un modèle projectif lisse explicite.

Dans les §§1, 2 et 3, on rappelle les bases de la théorie : suite spectrale de coniveau, complexes de Cousin, complexes de Gersten, conjecture de Gersten. Ces rappels, essentiellement fondés sur l'article [6], sont formulés pour une « théorie cohomologique à supports » quelconque qui satisfait à certains axiomes convenables. Des exemples de telles théories sont donnés au §4.

À partir du §6, on choisit comme théorie cohomologique la cohomologie motivique étale à coefficients entiers et on suppose que les variétés considérées sont lisses et géométriquement cellulaires (c'est-à-dire admettent une décomposition cellulaire sur la clôture algébrique) : c'est le cas par exemple des variétés projectives homogènes. On introduit le complément indispensable aux suites spectrales de coniveau : les suites spectrales dites « des poids », cf. [13]. La construction de ces suites spectrales repose sur la théorie des motifs triangulés de Voevodsky [44], ce qui oblige pour l'instant à supposer que le corps de base  $k$  est de caractéristique zéro.

Si  $X$  est une  $k$ -variété projective homogène, on souhaite calculer le noyau et le conoyau des homomorphismes

$$H^{n+2}(k, \mathbb{Z}(n)) \rightarrow H_{\text{nr}}^{n+2}(X, \mathbb{Z}(n)), n \geq 0. \quad (*)$$

La méthode est de considérer ensemble la suite spectrale de coniveau et la suite spectrale des poids, chacune en poids  $n$  : elles convergent toutes les deux vers la cohomologie motivique de poids  $n$  de  $X$ . La cohomologie non ramifiée faisant partie du terme  $E_2$  de la première suite spectrale et le terme  $E_2$  de la seconde étant en grande partie calculable, on peut espérer étudier (\*) de cette

manière. Des exemples sont donnés dans les §§6 à 10 : la plupart concernent l'étude de  $(*)$  pour les quadriques et pour  $n \leq 3$ , faite en collaboration avec Rost et Sujatha. Un bref aperçu de l'application de ces techniques aux groupes  $SK_1$  et  $SK_2$  des algèbres centrales simples est également donné au §9.

## 1 Partie non ramifiée d'une théorie cohomologique

**Définition 1.1 (pour ce mini-cours).** a) Soit  $k$  un anneau de base (noëthérien régulier). Nous utiliserons la catégorie  $P/k$  suivante :

- Les objets de  $P/k$  sont les couples  $(X, Z)$ , où  $X$  est un schéma régulier de type fini sur  $k$  et  $Z$  est un fermé (réduit) de  $X$ .
- Un morphisme  $f: (X', Z') \rightarrow (X, Z)$  est un morphisme  $f: X' \rightarrow X$  tel que  $f^{-1}(Z) \subset Z'$ .

b) Une *théorie cohomologique* (à supports) sur  $P/k$  est une famille de foncteurs

$$(h^q: (P/k)^o \rightarrow Ab)_{q \in \mathbf{Z}}$$

$$(X, Z) \mapsto h_Z^q(X)$$

vérifiant la condition suivante : pour tout triplet  $(Z \subset Y \subset X)$  avec  $(X, Y)$ ,  $(X, Z) \in P/k$ , on a une longue suite exacte

$$\cdots \rightarrow h_Z^q(X) \rightarrow h_Y^q(X) \rightarrow h_{Y-Z}^q(X - Z) \rightarrow h_Z^{q+1}(X) \rightarrow \cdots$$

fonctorielle en  $(X, Y, Z)$  en un sens évident.

On note  $h^q(X) = h_X^q(X)$  et on remarque que  $h_\emptyset^q(X) = 0$  pour tout  $(q, X)$ .

**Définition 1.2.** La théorie  $h^q$  vérifie l'*excision Zariski* (resp. *Nisnevich*) si elle est additive :

$$h_{Z \amalg Z'}^q(X \amalg X') = h_Z^q(X) \oplus h_{Z'}^q(X')$$

et si  $f^*: h_Z^q(X) \xrightarrow{\sim} h_{Z'}^q(X')$  lorsque  $f: (X', Z') \rightarrow (X, Z)$  est donnée par une immersion ouverte (resp. par un morphisme étale) tel que  $Z' = f^{-1}(Z)$  et  $f: Z' \xrightarrow{\sim} Z$ .

Si  $h^*$  vérifie l'excision Zariski, pour tout recouvrement ouvert  $X = U \cup V$  on a une longue suite exacte de Mayer–Vietoris :

$$\cdots \rightarrow h^q(X) \rightarrow h^q(U) \oplus h^q(V) \rightarrow h^q(U \cap V) \rightarrow h^{q+1}(X) \rightarrow \cdots$$

Si  $h^*$  vérifie l'excision Zariski, on peut construire des *complexes de Cousin* et une *suite spectrale de coniveau* (Grothendieck) :

A) Soit  $\vec{Z} = (\emptyset \subset Z_d \subset Z_{d-1} \subset \cdots \subset Z_0 = X)$  une chaîne de fermés. Les suites exactes

$$\begin{aligned} \cdots \rightarrow h_{Z_{p+1}}^{p+q}(X) &\xrightarrow{i^{p+1, q-1}} h_{Z_p}^{p+q}(X) \xrightarrow{j^{p, q}} h_{Z_p - Z_{p+1}}^{p+q}(X - Z_{p+1}) \\ &\xrightarrow{k^{p, q}} h_{Z_{p+1}}^{p+q+1}(X) \rightarrow \cdots \end{aligned}$$

définissent un couple exact

$$\begin{array}{ccc} D^{p+1,q-1} & \xrightarrow{i^{p+1,q-1}} & D^{p,q} \\ & \swarrow k^{p,q} \quad \searrow j^{p,q} & \\ & E^{p,q} & \end{array}$$

(où  $k^{p,q}$  est de degré  $(0, +1)$ ), avec  $D^{p,q} = h_{Z_p}^{p+q}(X)$ ,  $E^{p,q} = h_{Z_p - Z_{p+1}}^{p+q}(X - Z_{p+1})$ . Cela donne une suite spectrale de type cohomologique qui converge vers  $D^{0,n} = h^n(X)$ , la filtration associée étant

$$F^p h^n(X) = \text{Im} \left( h_{Z_p}^n(X) \rightarrow h^n(X) \right)$$

avec

$$E_1^{p,q} = E^{p,q}, \quad d_1^{p,q} = k j.$$

B) On suppose  $X$  équidimensionnel de dimension  $d$  et on ne s'intéresse qu'aux  $\vec{Z}$  tels que  $\text{codim}_X Z_p \geq p$ . On passe à la limite sur ces  $\vec{Z}$  : on obtient un nouveau couple exact, avec

$$D^{p,q} = \varinjlim_{\vec{Z}} h_{Z_p}^{p+q}(X) =: h_{\geq p}^{p+q}(X)$$

$$E^{p,q} = \varinjlim_{\vec{Z}} h_{Z_p - Z_{p+1}}^{p+q}(X - Z_{p+1}).$$

En utilisant l'excision Zariski, on trouve un isomorphisme

$$\varinjlim_{\vec{Z}} h_{Z_p - Z_{p+1}}^{p+q}(X - Z_{p+1}) \simeq \bigoplus_{x \in X^{(p)}} h_x^{p+q}(X)$$

où  $X^{(p)} = \{x \in X \mid \text{codim}_X \overline{\{x\}} = p\}$  et

$$h_x^{p+q}(X) := \varinjlim_{\substack{U \ni x \\ U \text{ ouvert}}} h_{\{x\} \cap U}^{p+q}(U)$$

(groupe de cohomologie locale), ce qui donne la forme classique du terme  $E_1$  de la *suite spectrale de coniveau* :

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)}} h_x^{p+q}(X) \Rightarrow h^{p+q}(X). \quad (1)$$

La filtration à laquelle elle aboutit est la filtration par la codimension du support

$$\begin{aligned} N^p h^n(X) &= \bigcup_{\text{codim}_X Z \geq p} \text{Im} \left( h_Z^n(X) \rightarrow h^n(X) \right) \\ &= \bigcup_{\text{codim}_X Z \geq p} \text{Ker} \left( h^n(X) \rightarrow h^n(X - Z) \right). \end{aligned}$$

**Définition 1.3.** a) Le *complexe de Cousin en degré  $q$  de  $h$  sur  $X$*  est le complexe des termes  $E_1$  de la suite spectrale :

$$0 \rightarrow \bigoplus_{x \in X^{(0)}} h_x^q(X) \xrightarrow{d_1^{0,q}} \bigoplus_{x \in X^{(1)}} h_x^{1+q}(X) \xrightarrow{d_1^{1,q}} \dots$$

$$\xrightarrow{d_1^{p-1,q}} \bigoplus_{x \in X^{(p)}} h_x^{p+q}(X) \xrightarrow{d_1^{p,q}} \dots$$

b) La *cohomologie non ramifiée de  $h$  sur  $X$*  (en degré  $q$ ) est le groupe

$$E_2^{0,q} = \text{Ker} \left( \bigoplus_{x \in X^{(0)}} h_x^q(X) \xrightarrow{d_1^{0,q}} \bigoplus_{x \in X^{(1)}} h_x^{1+q}(X) \right) =: h_{\text{nr}}^q(X).$$

Si  $X = X_1 \amalg \dots \amalg X_r$ , on a  $h_{\eta_i}^q(X) = h_{\eta_i}^q(X_i) = \varinjlim_{U \subset X_i} h^q(U)$ , où  $\eta_i$  est le point générique de  $X_i$  : ce groupe ne dépend que de  $\eta_i$  et nous le noterons habituellement  $h^q(\eta_i)$  ou  $h^q(K_i)$  si  $\eta_i = \text{Spec } K_i$ . On a  $h_{\text{nr}}^q(X) = \bigoplus_i h_{\text{nr}}^q(X_i)$ . Pour  $X$  connexe, on a donc

$$h_{\text{nr}}^q(X) = \text{Ker} \left( h^q(\eta) \rightarrow \bigoplus_{x \in X^{(1)}} h_x^{1+q}(X) \right).$$

## 2 Pureté ; complexes de Cousin et complexes de Gersten

On se donne une théorie cohomologique graduée

$$h^* : (X, Z) \mapsto h_Z^q(X, n), \quad q, n \in \mathbf{Z}.$$

(L'entier  $n$  s'appelle le *poids*.)

**Définition 2.1.**  $h^*$  est *pure* si, pour tout  $(X, Z) \in P/k$  avec  $X$  régulier et  $Z$  régulier purement de codimension  $c$  dans  $X$ , on s'est donné des isomorphismes

$$\pi_{X,Z} : h^{q-2c}(Z, n-c) \xrightarrow{\sim} h_Z^q(X, n)$$

contravariants en les  $(X, Z)$  comme au-dessus (à  $c$  fixé).

(On dit que  $h^*$  est *faiblement pure* si la pureté n'est exigée que pour  $X$  et  $Z$  lisses sur  $k$  : si  $k$  est un corps parfait, cela revient au même.) Si  $k$  est raisonnable (par exemple un corps ou  $\text{Spec } \mathbf{Z}$ ), cette condition entraîne l'excision Nisnevich : c'est évident pour des couples comme dans la définition, et en général on s'y ramène par récurrence noéthérienne en considérant le lieu non régulier de  $Z$ , qui est fermé et différent de  $Z$ .

Si  $h^*$  est pure, la suite spectrale (1) prend la forme peut-être plus familière

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)}} h^{q-p}(\kappa(x), n-p) \Rightarrow h^{p+q}(X). \quad (2)$$

En particulier, les complexes de Cousin deviennent des *complexes de Gersten* (on suppose  $X$  connexe pour simplifier) :

$$0 \rightarrow h^q(\kappa(X), n) \rightarrow \bigoplus_{x \in X^{(1)}} h^{q-1}(\kappa(x), n-1) \rightarrow \bigoplus_{x \in X^{(2)}} h^{q-2}(\kappa(x), n-2) \dots$$

et on retrouve une définition plus familière de  $h_{\text{nr}}$  :

$$h_{\text{nr}}^q(X, n) = \text{Ker} \left( h^q(\kappa(X), n) \rightarrow \bigoplus_{x \in X^{(1)}} h^{q-1}(\kappa(x), n-1) \right).$$

*Remarque 2.2.* Dans certains cas, on n'a la pureté qu'à *isomorphisme près* ; pour obtenir des isomorphismes de pureté canoniques, on doit introduire des variantes de la théorie  $h$ , à coefficients dans des fibrés en droites. C'est le cas notamment pour les groupes de Witt triangulaires de Barge–Sansuc–Vogel, Pardon, Ranicki et Balmer–Walter ([3], voir aussi [39]).

### 3 Conjecture de Gersten

**Définition 3.1.** Pour tout  $(p, q)$ , on note  $\mathcal{E}_1^{p,q}$  le faisceau associé au préfaisceau Zariski

$$U \mapsto E_1^{p,q}(U) = \bigoplus_{x \in U^{(p)}} h_x^{p+q}(U).$$

On a ainsi pour tout  $q$  un complexe de faisceaux

$$0 \rightarrow \mathcal{H}^q \rightarrow \mathcal{E}_1^{0,q} \rightarrow \mathcal{E}_1^{1,q} \rightarrow \dots \rightarrow \mathcal{E}_1^{p,q} \rightarrow \dots$$

avec les  $\mathcal{E}_1^{p,q}$  *flasques* pour la topologie de Zariski, où  $\mathcal{H}^q$  est le faisceau associé au préfaisceau  $U \mapsto h^q(U)$ .

**Définition 3.2.** On dit que  $h$  vérifie la *conjecture de Gersten* sur  $X$  si ce complexe est *exact* pour tout  $q$ .

Si c'est le cas, le complexe

$$0 \rightarrow \mathcal{E}_1^{0,q} \rightarrow \mathcal{E}_1^{1,q} \rightarrow \dots \rightarrow \mathcal{E}_1^{p,q} \rightarrow \dots$$

définit une résolution flasque de  $\mathcal{H}^q$ , et on peut écrire le terme  $E_2$  de la suite spectrale de coniveau

$$E_2^{p,q} = H_{\text{Zar}}^p(X, \mathcal{H}^q).$$

**Théorème 3.3 (Gabber [8], essentiellement).** *Supposons que  $k$  soit un corps infini. Alors, pour que  $h$  vérifie la conjecture de Gersten sur tout  $X$  lisse sur  $k$ , il suffit que les deux conditions suivantes soient vérifiées :*

(1)  *$h$  vérifie l'excision Nisnevich.*



(2) Lemme clé. Pour tout  $n$ , pour tout ouvert  $V$  de  $\mathbb{A}_k^n$ , pour tout fermé  $F \subset V$  et pour tout  $q \in \mathbf{Z}$ , le diagramme de gauche est commutatif :

$$\begin{array}{ccc}
 h_{\mathbb{A}_F}^q(\mathbb{A}_V^1) & \xleftarrow{j^*} & h_{\mathbb{P}_F^1}^q(\mathbb{P}_V^1) \\
 \searrow \pi^* & & \downarrow s_\infty^* \\
 & & h_F^q(V)
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{A}_V^1 & \xrightarrow{j} & \mathbb{P}_V^1 \\
 \searrow \pi & & \downarrow \tilde{\pi} \\
 & & V
 \end{array}
 \quad \begin{array}{c} \uparrow s_\infty \end{array}$$

où  $s_\infty$  est la section à l'infini.

La condition (2) est vérifiée dans chacun des cas suivants :

- (3)  $h$  est invariante par homotopie : pour tout  $V$  lisse,  $h^*(V) \xrightarrow{\sim} h^*(\mathbb{A}_V^1)$  (il suffit que ce soit vrai pour  $V$  comme en (2)).
- (4)  $h$  est « orientable » : il existe une théorie cohomologique  $e$  et, pour tout  $(X, Z) \in P_k$ , une application

$$\mathrm{Pic}(X) \rightarrow \mathrm{Hom}(e_Z^*(X), h_Z^*(X))$$

naturelle en  $(X, Z)$ , d'où (pour  $(X, Z) = (\mathbb{P}_V^1, \mathbb{P}_Z^1)$ ) un homomorphisme  $\alpha_{V,F}$

$$\begin{array}{ccc}
 e_{\mathbb{P}_F^1}^*(\mathbb{P}_V^1) & \xrightarrow{[\mathcal{O}(1)] - [\mathcal{O}]} & h_{\mathbb{P}_F^1}^*(\mathbb{P}_V^1) \\
 \uparrow \tilde{\pi} & \nearrow \alpha_{V,F} & \\
 e_F^*(V) & & 
 \end{array}$$

et, pour  $(V, F)$  comme en (2), l'application

$$h_F^q(V) \oplus e_F^q(V) \xrightarrow{(\pi^*, \alpha_{V,F})} h_{\mathbb{P}_F^1}^q(\mathbb{P}_V^1)$$

est un isomorphisme.

*Preuve.* Voir [6]. Pour  $k$  fini, on s'en tire en supposant l'existence de transferts sur  $h$  (pour des revêtements étales provenant d'extensions du corps de base).  $\square$

## Conséquences pour la cohomologie non ramifiée

**Théorème 3.4.** Sous les hypothèses (1) et (2) du théorème 3.3, pour toute variété  $X$  lisse sur  $k$  :

a)  $h_{\mathrm{nr}}^q(X) \simeq H_{\mathrm{Zar}}^0(X, \mathcal{H}^q) \simeq H_{\mathrm{Nis}}^0(X, \mathcal{H}^q)$ , où  $H_{\mathrm{Nis}}^*$  désigne la cohomologie de Nisnevich (ceci s'étend à tous les termes  $E_2$  de la suite spectrale de coniveau, et ne sera pas utilisé ici).

b) Si  $X$  est de plus propre,  $h_{\mathrm{nr}}^q(X)$  est un invariant birationnel.

c) Soient  $X, Y$  lisses et intègres et  $p: X \rightarrow Y$  un morphisme propre. Supposons que la fibre générique de  $p$  soit  $k(Y)$ -birationnelle à l'espace projectif  $\mathbb{P}_{k(Y)}^d$ . Alors,  $h_{\mathrm{nr}}^q(X) \xrightarrow{p^*} h_{\mathrm{nr}}^q(Y)$  est un isomorphisme.