

**J. Gmehling
U. Onken
W. Arlt**

VAPOR-LIQUID EQUILIBRIUM DATA COLLECTION

Halogen, Nitrogen, Sulfur and other Compounds



Chemistry Data Series

Vol. I, Part 8

Vapor-Liquid Equilibrium Data Collection

8

**Halogen, Nitrogen, Sulfur and other
Compounds**

Tables and diagrams of data for binary and
multicomponent mixtures up to moderate pressures.
Constants of correlation equations for computer use.

J. Gmehling, U. Onken, W. Arlt

Lehrstuhl Technische Chemie B
(Prof. Dr. U. Onken)
Universität Dortmund

© 1984 DECHEMA Deutsche Gesellschaft für Chemisches Apparatewesen
6000 Frankfurt/Main, Federal Republic of Germany

ISBN 3-921 567-24-6

All rights reserved

No part of this work may be reproduced or used in any form or by any means — graphic, electronic, or mechanical, including photocopying, recording, or taping, or information storage and retrieval systems — without written permission of the publishers.

Printed by Schön & Wetzel GmbH, Frankfurt/Main, F. R. Germany

J. Gmehling
U. Onken
W. Arlt

VAPOR-LIQUID EQUILIBRIUM DATA COLLECTION

Halogen, Nitrogen, Sulfur and other Compounds



Chemistry Data Series

Vol. I, Part 8

Published by DECHEMA
Deutsche Gesellschaft für Chemisches Apparatewesen
Editors: Dieter Behrens, Reiner Eckermann

8

Halogen, Nitrogen, Sulfur and other Compounds

Systems with:

Halogenated Hydrocarbons

Silanes, Siloxanes

Nitro Compounds, Amines, Imines, Oximes, Nitriles,
Isocyanates, Amides, Pyrrolidines, Pyridines,
Quinolines

Mercaptans, Sulfides, Sulfoxides, Carbon Disulfide,
Thiophenes, Thiazoles

SUBJECTS OF VOLUME I

The table lists all parts of Volume I with year of publication.

Subtitle	Vol. I, Part
Aqueous-Organic Systems	1 (1980)
Supplement 1	1a (1981)
Organic Hydroxy Compounds	
Alcohols	2a (1977)
Alcohols and Phenols	2b (1978)
Supplement 1	2c (1982)
Supplement 2	2d (1982)
Aldehydes, Ketones, Ethers	3/4 (1979)
Carboxylic Acids, Anhydrides, Esters	5 (1982)
Aliphatic Hydrocarbons C ₄ -C ₆	6a (1980)
Aliphatic Hydrocarbons C ₇ -C ₁₈	6b (1980)
Supplement 1	6c (1984)
Aromatic Hydrocarbons	7 (1980)
Halogen, Nitrogen, Sulfur and other Compounds	8 (1984)

AUTHORS' PREFACE

The appearance of this volume marks the completion of our VLE data collection. When we began publishing this series in 1977, we were not aware of the amount of work ahead. Perhaps if we really had known then what we were embarking upon, we might not have started. And we were able to continue during these years because we have been lucky enough to obtain collaboration and to find encouragement from all the members of our team and from many colleagues.

In preparing this part we were helped by Dipl.-Chem. J. Berg, Dipl.-Ing. P. Grenzheuser, Mrs. Dr. B. Kolbe, Mrs. L. Kunzner, Dipl.-Chem. J. Menke, Mrs. Dipl.-Chem. U. Schwaitzer, Dipl.-Chem. D. Tiegs, Dipl.-Chem. U. Weidlich and by the students R. Rarey and J. Schomschor. For translating some sources of VLE data we were able to draw on the help of two of our foreign guests during their research stay with us: Ing. Yang Zhendong from the People's Republic of China, who translated several publications from Chinese, and Ing. C. Osorio Bastiades, who assisted us in understanding some Spanish nomenclature. They and all co-workers mentioned in the other parts of our data collection deserve our wholehearted thanks. Likewise we should again like to express our gratitude to the computer center of the University of Dortmund, especially to Mr. T. Blaszyk for his co-operation, and to Dr. R. Eckermann and Dipl.-Ing. C. Hammer (both DECHEMA, Frankfurt/M.) for their efforts in the editing of our data collection.

Finally, we must not forget the numerous suggestions and the stimulation coming from colleagues from all over the world. We shall continue our work on the DDB (Dortmund VLE Data Bank), adding newly published results and publishing supplements to the data collection from time to time. On this occasion we should again like to ask all workers in the field of vapour-liquid equilibrium to send us reprints containing new results. This will help to ensure that this truly international partnership can grow and continue.

Dortmund, May 1984

Ulfert Onken

Jürgen Gmehling

Wolfgang Arlt

PREFACE OF EDITORS

Subjects of the Dechema Chemistry Data Series are the physical and thermodynamic property data of chemical compounds and mixtures essentially for the fluid state covering PVT data, heat capacity, enthalpy, and entropy data, phase equilibrium data, transport and interfacial tension data.

The main purpose is to provide chemists and engineers with data for process design and development. For computer based calculations in process design appropriate correlation methods and accurate data must be used. These are only in some cases available in the open literature. For that reason the most urgent requirement regarding the publication of data is to offer classified and critically evaluated data, thus giving an impression which of them are reliable or not. This will be the goal of the series.

DECHEMA gives the opportunity to authors especially from universities to publish not only their theoretical results, but also their measured or compiled data, most often a large amount, that would otherwise never have been published.

The research work of Dr. Gmehling, Prof. Onken and co-workers on vapor-liquid equilibria which was partly supported by the Federal Ministry of Research and Technology and DECHEMA has been very fruitful; in particular, it led to an extension of the UNIFAC method. The authors have produced what is probably the largest collection of vapor-liquid equilibrium data that is today available with evaluation programs and experimental data.

We present the evaluation of this material in several parts of the first volume of the series. We hope that this gives particularly the users an instrument that will allow them to solve their problems considerably more easily and quickly than before.

Frankfurt/Main, May 1984

Dieter Behrens
Reiner Eckermann

GUIDE TO TABLES

1. Order of Succession of Systems of Data Sets.

- 1.1 In this part binary and ternary systems with halogen, nitrogen, sulfur and other compounds are given.
- 1.2 Within each class of mixtures all binary systems are given first, followed by ternary and quaternary systems.
- 1.3 The sequence of systems is based on the empirical formula convention used in the index of Chemical Abstracts, which means arranging the compounds according to increasing C and H, with the remaining elements in alphabetical order: Br, Cl, F, I, N, O, S. Compounds with identical empirical formula are arranged alphabetically according to their names.
- 1.4 Order of succession for different data sets for a given system (a set of data points, taken from one original paper, which have been measured varying the composition at constant temperature or pressure, is referred to as a data set):
 1. Name of first author in alphabetical order,
 2. isothermal data sets, with increasing temperature,
 3. isobaric data sets, with increasing pressure,
 4. recommended values (see section 7).

2. Antoine Vapor Pressure Equation

The Antoine vapor pressure equation is used in the following form:

$$\log[p_i^0] = A - \frac{B}{t + C} \quad (70)$$

with $[p_i^0]$ vapor pressure of pure component i in mm Hg
 t temperature in degrees Celsius ($^{\circ}$ C)

The Antoine constants A, B, and C are given with respective temperature regions (in $^{\circ}$ C).

If pure component vapor pressures are given together with the VLE data in the original paper, the first Antoine constant A is fitted to these data with the two other Antoine constants B and C being taken from our data files for pure compounds, as explained in the General Remarks and Explanations (part 1, p. XXXIX). If the Antoine constant A has been fitted in this way, it is given at the bottom of the respective table, using the symbol A'.

3. Consistency Tests

Method 1: Point test of Van Ness et al. [5] in the version of Fredenslund et al. [1].

Method 2: Integral or area test of Redlich - Kister [4] and Herington [2, 3].

Meaning of symbols (see also **Table 1**):

+	consistent
-	not consistent
blank	no result

For more detailed information see sections 2.3 (p. XXII to XXIX) and 4.2 (p. XXXVIII to XLIII) of General Remarks and Explanations in Part 1 of this volume.

4. Parameters of Activity Coefficient Equations

Parameters for the following equations are given:

Margules, eq. (28)
van Laar, eq. (29)
Wilson, eqs. (30) and (36)
NRTL, eqs. (31) and (37)
UNIQUAC, eqs. (32) to (35) and (38) to (40)

Equations and parameters are given in **Tables 2 to 4** and in Tables 1 to 4 of General Remarks and Explanations, (p. XVI to XXI), Part 1.

For ternary systems, only Wilson, NRTL, and UNIQUAC parameters are given; for systems with more than three components no parameters of activity coefficient equations are given.

For notation of constants of activity coefficient equations see **Table 2**.

Wilson, NRTL, and UNIQUAC parameters are given in cal/mol with the gas constant $R = 1.98721$ cal/mol K and the temperature T in K.

Note: In Volume V of the DECHEMA Chemistry Data Series (Liquid-Liquid Equilibria [11]), parameters are given in K.

Data for the pure liquid components, which are required for the Wilson equation (molar volume V_1^L) and UNIQUAC equation (volume parameter r_i and area parameter q_i), are given in Appendix A.

5. Vapor – Liquid Equilibrium Data

5.1 Experimental data

<i>Symbol</i>	<i>Meaning</i>
P MM HG	pressure in mm Hg
T DEG C	temperature in °C (degrees Celsius)
X1 (X2 . . .)	liquid mole fraction x_1 ($x_1 . . .$)
Y1 (Y2 . . .)	vapor mole fraction y_1 ($y_1 . . .$)

Table 1 Thermodynamic Consistency Tests

Character in the table	Method 1 Van Ness et al. [5] and Fredenslund et al. [1]	Method 2 Redlich-Kister [4] and Herington [2,3]
+ (consistent)	$\overline{\Delta Y} \leq 0,01$ ¹⁾	isothermal data: $D \leq 10\%$ ²⁾ isobaric data: $(D-J) \leq 10\%$ ³⁾ exception: for nearly ideal systems with $0.95 < \gamma < 1.10$ (for all γ_i) method 2 is not employed, but character + is given.
- (not consistent)	$\overline{\Delta Y} > 0.01$ ¹⁾	either a) isothermal data: $D > 10\%$ ²⁾ isobaric data: $(D-J) > 10\%$ ³⁾ or b) maximum deviation in $\ln \frac{\gamma_1}{\gamma_2}$ between 3 rd order interpolation polynomial and experimental data point > 1.17 fold mean deviation
blank (no result)	calculation does not converge	a) less than 5 data points in the set b) interpolation polynomial does not intersect x-axis ⁴⁾
1) $\overline{\Delta Y}$ $\Delta Y = Y_{\text{exp}} - Y_{\text{calc}}$	mean deviation in vapor mole fraction y	3) $J = 150 \cdot \frac{ \Delta T_{\text{max}} }{T_{\text{min}}} [\%]$
2) D	$D = \frac{ A' - B' }{A' + B'} \cdot 100 [\%]$	4) this may happen, when all data points are within a small range of composition near one of the pure compounds.

Table 2 Activity Coefficient Equations for Binary Systems and Notation of Parameters*

Type of Equation	Parameters	$\ln \gamma_1 =$ $\ln \gamma_2 =$	Notation of Parameters in Data Sheet
Margules [6]	A ₁₂	$[A_{12} + 2(A_{21} - A_{12}) x_1] x_2^2$	A 12
	A ₂₁	$[A_{21} + 2(A_{12} - A_{21}) x_2] x_1^2$	A 21
van Laar [7]	A ₁₂	$A_{12} \left(\frac{A_{21} x_2}{A_{12} x_1 + A_{21} x_2} \right)^2$	A 12
	A ₂₁	$A_{21} \left(\frac{A_{12} x_1}{A_{12} x_1 + A_{21} x_2} \right)^2$	A 21
Wilson [8]	$\lambda_{12} - \lambda_{11}^1$	$-\ln(x_1 + \Lambda_{12} x_2) + x_2 \left(\frac{\Lambda_{12}}{x_1 + \Lambda_{12} x_2} - \frac{\Lambda_{21}}{\Lambda_{21} x_1 + x_2} \right)$	A 12
	$\lambda_{21} - \lambda_{22}$	$-\ln(x_2 + \Lambda_{21} x_1) - x_1 \left(\frac{\Lambda_{12}}{x_1 + \Lambda_{12} x_2} - \frac{\Lambda_{21}}{\Lambda_{21} x_1 + x_2} \right)$	A 21
NRTL [9] (continued)	$g_{12} - g_{22}^2$	$x_2^2 \left[\tau_{21} \left(\frac{G_{21}}{x_1 + x_2 G_{21}} \right)^2 + \left(\frac{\tau_{12} G_{12}}{(x_2 + x_1 G_{12})^2} \right) \right]$	A 12
	$g_{21} - g_{11}$ α_{12}	$x_1^2 \left[\tau_{12} \left(\frac{G_{12}}{x_2 + x_1 G_{12}} \right)^2 + \left(\frac{\tau_{21} G_{21}}{(x_1 + x_2 G_{21})^2} \right) \right]$	A 21 ALPHA 12

Table 2 (continued)

Type of Equation	Parameters	$\ln \gamma_1 =$ $\ln \gamma_2 =$	Notation of Parameters in Data Sheet
UNIQUAC [10]	$u_{12} - u_{22}$ ³⁾ $u_{21} - u_{11}$	$\ln \gamma_1^C + \ln \gamma_1^R$ ³⁾ $\ln \gamma_2^C + \ln \gamma_2^R$	A 12 A 21
1)	$\Lambda_{12} = \frac{V_2^L}{V_1^L} \exp - \frac{\lambda_{12} - \lambda_{11}}{RT}$	$\Lambda_{21} = \frac{V_1^L}{V_2^L} \exp - \frac{\lambda_{21} - \lambda_{22}}{RT}$	
<p>V_i^L molar volume of pure liquid component i. For values of V_i^L see Appendix A.</p> <p>λ_{ij} interaction energy between components i and j $\lambda_{ij} = \lambda_{ji}$</p>			
2)	$\tau_{12} = \frac{g_{12} - g_{22}}{RT}$	$\tau_{21} = \frac{g_{21} - g_{11}}{RT}$	
	$G_{12} = \exp(-\alpha_{12}\tau_{12})$	$G_{21} = \exp(-\alpha_{21}\tau_{21})$	
<p>g_{ij} parameter for interaction between components i and j; $g_{ij} = g_{ji}$</p> <p>α_{ij} nonrandomness parameter; $\alpha_{ij} = \alpha_{ji}$</p>			
3)	for detailed equations see table 4		
<p>*) Wilson, NRTL, and UNIQUAC parameters are given in cal/mol with the gas constant R = 1.98721 cal/mol K and the temperature T in K.</p>			

Table 3 Activity Coefficient Equations for Multicomponent Systems

Type of Equation	Parameters	$\ln \gamma_i =$
Wilson [8] *)	1) $\Lambda_{ij} = \frac{V_j^L}{V_i^L} \exp - \left[\frac{(\lambda_{ij} - \lambda_{ii})}{RT} \right]$ $\Lambda_{ii} = \Lambda_{jj} = 1$	1) $-\ln \left(\sum_{j=1}^m x_j \Lambda_{ij} \right) + 1 - \sum_{k=1}^m \frac{x_k \Lambda_{ki}}{\sum_{j=1}^m x_j \Lambda_{kj}}$ (36)
NRTL [9]	2) $\tau_{ji} = \frac{(g_{ji} - g_{ii})}{RT}$ $G_{ji} = \exp (-\alpha_{ji} \tau_{ji})$ $\tau_{ii} = \tau_{jj} = 0$ $G_{ii} = G_{jj} = 1$	2) $\sum_{j=1}^m \frac{\tau_{ji} G_{ji} x_j}{\sum_{i=1}^m G_{ii} x_i} + \sum_{j=1}^m \frac{x_j G_{jj}}{\sum_{i=1}^m G_{ij} x_i} - \left(\tau_{ij} - \frac{\sum_{n=1}^m x_n \tau_{nj} G_{nj}}{\sum_{i=1}^m G_{ij} x_i} \right)$ (37)
UNIQUAC [10] *) (continued)	3) $\tau_{ji} = \exp - \left(\frac{u_{ji} - u_{ii}}{RT} \right)$	3) $\ln \gamma_i^C + \ln \gamma_i^R$ (38)

Table 3 (continued)

	$\tau_{ii} = \tau_{jj} = 1$	$\ln \gamma_i^C = \ln \frac{\varphi_i}{x_i} + \frac{z}{2} q_i \ln \frac{\vartheta_i}{\varphi_i} + l_i - \frac{\varphi_i}{x_i} \sum_j x_j l_j \quad (39)$ $\ln \gamma_i^R = q_i \left[1 - \ln \left(\sum_{j=1}^m \vartheta_j \tau_{ji} \right) - \sum_{j=1}^m \sum_{k=1}^m \frac{\vartheta_j \tau_{ij}}{\sum_{k=1}^m \vartheta_k \tau_{kj}} \right] \quad (40)$ $l_i = \frac{z}{2} (r_i - q_i) - (r_i - 1) \quad z = 10 \quad (35)$
<p>1) explanation of symbols see footnote 1 of table 2 2) explanation of symbols see footnote 2 of table 2 3) explanation of symbols see table 4</p> <p>*) For values of V_i^L, r_i and q_i see Appendix A</p>		

Table 4 UNIQUAC Activity Coefficient Equations [10] for Binary Systems

$\ln \gamma_1 = \ln \gamma_1^C + \ln \gamma_1^R \quad (32a)$	
$\ln \gamma_1^C = \ln \frac{\varphi_1}{x_1} + \frac{z}{2} q_1 \ln \frac{\vartheta_1}{\varphi_1} + \varphi_2 \left(l_1 - \frac{r_1}{r_2} l_2 \right) \quad (33a)$	
$\ln \gamma_1^R = -q_1 \ln (\vartheta_1 + \vartheta_2 \tau_{21}) + \vartheta_2 q_1 \left(\frac{\tau_{21}}{\vartheta_1 + \vartheta_2 \tau_{21}} - \frac{\tau_{12}}{\vartheta_1 \tau_{12} + \vartheta_2} \right) \quad (34a)$	
$\ln \gamma_2 = \ln \gamma_2^C + \ln \gamma_2^R \quad (32b)$	
$\ln \gamma_2^C = \ln \frac{\varphi_2}{x_2} + \frac{z}{2} q_2 \ln \frac{\vartheta_2}{\varphi_2} + \varphi_1 \left(l_2 - \frac{r_2}{r_1} l_1 \right) \quad (33b)$	
$\ln \gamma_2^R = -q_2 \ln (\vartheta_1 \tau_{12} + \vartheta_2) + \vartheta_1 q_2 \left(\frac{\tau_{12}}{\vartheta_1 \tau_{12} + \vartheta_2} - \frac{\tau_{21}}{\vartheta_1 + \vartheta_2 \tau_{21}} \right) \quad (34b)$	
$l_i = \frac{z}{2} (r_i - q_i) - (r_i - 1) \quad z = 10^{\sigma} \quad (35)$	
Symbols	
l_i	see equation (35)
q_i	area parameter of component i *)
r_i	volume parameter of component i *)
u_{ij}	Parameter of interaction between components i and j; $u_{ij} = u_{ji}$
z	coordination number
γ_i^C	combinatorial part of activity coefficient of component i
γ_i^R	residual part of activity coefficient of component i
$\vartheta_i = \frac{q_i x_i}{\sum_j q_j x_j}$	area fraction of component i
$\varphi_i = \frac{r_i x_i}{\sum_j r_j x_j}$	volume fraction of component i
τ_{ij}	see Table 3
*) For values of r_i and q_i see Appendix A	

5.2 Calculated data

<i>Symbol</i>	<i>Meaning</i>
DIFF P	$P = P_{\text{exp}} - P_{\text{calc}}$
DIFF T	$T = T_{\text{exp}} - T_{\text{calc}}$
DIFF Y1	$Y_1 = Y_{1,\text{exp}} - Y_{1,\text{calc}}$
DIFF Y2	$Y_2 = Y_{2,\text{exp}} - Y_{2,\text{calc}}$

$$\text{MEAN DEVIATION e.g. for T: } \Delta T = \frac{1}{n} \sum_n |\Delta T|$$

MAX. DEVIATION maximum absolute deviation

Results of calculations with activity coefficient equations are usually tabulated as the difference of experimental value minus calculated value. Calculated vapor mole fractions are given directly only for isothermal P-x data and isobaric T-x data. In all other cases the calculated values are obtained by subtracting the tabulated difference from the experimental value, e.g. $Y_{1,\text{calc}} = Y_{1,\text{exp}} - \Delta Y_1$.

6. Equilibrium Diagrams

6.1 Binary Systems

For complete data sets and for isothermal x - y data, x - y diagrams show data points and equilibrium line calculated by activity coefficient equation yielding the lowest mean deviation Δy_1 . For sets of data without experimental vapor compositions y (isothermal P-x data and isobaric T-x data), the equilibrium line from the equation showing the best fit (lowest mean ΔP or ΔT , for isothermal and isobaric data respectively) is plotted. The type of this equation is printed on the diagram, together with the values for the activity coefficients at infinite dilution γ_1^∞ and γ_2^∞ (for equations see Table 3 of General Remarks and Explanations, p. XIX of Part 1).

In the case of limited liquid miscibility, calculated solubilities (which may be different from experimental values!), are inserted into the equilibrium line as small circles connected by a horizontal line.

6.2 Ternary Systems

Triangular diagrams show experimental data points with equilibrium composition of vapor (sign \ast) and liquid (sign \circ) connected by a straight line.