

科技资料

# Current Topics in Operator Algebras

Proceedings of the Satellite Conference of ICM - 90

# CURRENT TOPICS IN OPERATOR ALGEBRAS

Nara-ken New Public Hall, Japan 16 - 19 August 1990

Edited by

H. Araki

(Kyoto University)

H. Choda

(Osaka Kyoiku University)

Y. Nakagami

(Yokohama City University)

K. Saitô

(Tohoku University)

J. Tomiyama

(Tokyo Metropolitan University)



**World Scientific**

Singapore • New Jersey • London • Hong Kong

*Published by*

World Scientific Publishing Co. Pte. Ltd.

P O Box 128, Farrer Road, Singapore 9128

USA office: 687 Hartwell Street, Teaneck, NJ 07666

UK office: 73 Lynton Mead, Totteridge, London N20 8DH

## CURRENT TOPICS IN OPERATOR ALGEBRAS

Copyright © 1991 by World Scientific Publishing Co. Pte. Ltd.

*All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the Publisher.*

ISBN 981-02-0651-8

Printed in Singapore by JBW Printers & Binders Pte. Ltd.

World Scientific  
Singapore - New Jersey - London - Hong Kong

## Preface

It is our greatest pleasure to present here the Proceedings of a satellite conference of ICM-90 "Current Topics in Operator Algebras", held on August 16-19 at the Nara-ken New Public Hall, Japan. The topics covered include the  $C^*$ -dynamical systems, the index theory for subfactors, the noncommutative differential geometry, the quantum groups and so on.

We hope that the present volume serves the purpose of presenting an overview of the present status of the theory of operator algebras as well as an outlook for its future development.

We greatly acknowledge the cooperation of the research group of operator theory and operator algebras in Japan and the sponsorship of the Mathematical Society of Japan and Nara-ken for the conference. The faculties of the Department of Mathematics of the Yokohama City University, especially Professor Shigeo Ichiraku, were helpful throughout the preparation of the conference. The transportation from the Osaka International Airport to Nara was offered by Professor Mitsutoshi Hayashi of Osaka Institute of Technology. Professor Satoshi Kawakami of the Nara University of Education was helpful in providing us with the services during the meeting despite the summer staff shortage.

We appreciate the contributions from the following donors, Nara-ken, the Japan Society for Promotion of Science, Inoue Foundation for Science, the Grant-in-Aid of the Ministry of Education and many others listed in the last page of the Proceedings, especially the President Yukitsugu Shimokawa of Nippon Information Industry Co. Ltd.

Finally we thank speakers, contributors, chairpersons and participants of the conference for making this conference possible.

### Organizers

H. Araki, H. Choda, Y. Nakagami,  
K. Saitô, J. Tomiyama

January, 1991

## Contents

1	Quantum symmetry and subfactors <i>Adrian Ocneanu</i>	1
2	Group actions on injective factors <i>Yasuyuki Kawahigashi</i>	2
3	On the classification of inductive limits of inner actions of a compact group <i>Ola Bratteli, George A. Elliott, David Evans and Akitaka Kishimoto</i>	13
4	Real rank of $C^*$ -algebras <i>Gert K. Pedersen</i>	25
5	Asymptotic Galois correspondence on AFD factors <i>Yoshikazu Katayama</i>	33
6	Subfactors and ergodic theory <i>Colin E. Sutherland</i>	38
7	The form of invariant subspaces of analytic crossed products <i>Kichi-Suke Saito</i>	43
8	Operator algebras and non-standard analysis <i>Masanao Ozawa</i>	52
9	Minimum index for subfactors and entropy for canonical shifts <i>Fumio Hiai</i>	59
10	Classification and values of the index for subfactors <i>Sorin Popa</i>	71
11	Angles and entropies between two subfactors <i>Takashi Sano and Yasuo Watatani</i>	72
12	Commuting squares and subfactors <i>V.S. Sunder</i>	83

13	Invariants for subfactors arising from Coxeter graphs	84
	<i>Shingo Okamoto</i>	
14	An example of a subfactor of the hyperfinite $II_1$ factor	104
	whose principal graph invariant is the Coxeter graph $E_6$	
	<i>Jocelyne Bion-Nadal</i>	
15	Structural characterization of compact operators	114
	<i>José M. Isidro</i>	
16	The $C^*$ -algebra of a homogeneous ideal in two variables is	130
	type $I$	
	<i>Raul E. Curto, Paul S. Muhly and Keren Yan</i>	
17	$C^*$ -systems without norm-continuity properties	137
	<i>Anton Amann</i>	
18	Recent developments in operator spaces	146
	<i>Edward G. Effros and Zhong-Jin Ruan</i>	
19	An order of tensor products in $C^*$ -algebras	165
	<i>Takashi Itoh</i>	
20	The tensor product of dual spaces	172
	<i>Sa-Ge Lee</i>	
21	$C^*$ -algebraic isomorphism determined by completely positive	178
	maps and completely bounded maps	
	<i>Liang-Sen Wu</i>	
22	A functional calculus for pairs of non-commuting selfadjoint	185
	operators	
	<i>V. Peller</i>	
23	Partial $*$ -algebras of closable operators	186
	<i>J.-P. Antoine</i>	
24	Partial $W^*$ -dynamical systems	202
	<i>G.O.S. Ekhaguere</i>	

- 25 An application of the Tomita-Takesaki theory of 2D foliations 218  
with harmonic measures  
**Benjamin Enriquez**
- 26 Cyclic cohomology, higher indices and non-local invariants 222  
**Henri Moscovici**
- 27 Subfactors arising from representations of compact and infinite-  
dimensional groups 223  
**Antony Wassermann**
- 28 On commuting squares 224  
**Vaughan F. R. Jones**
- 29 Elliptic renormalization 225  
**Ronald G. Douglas**
- 30 Non commutative 3-spheres 234  
**Kengo Matsumoto**
- 31 Analytic invariants for algebraic  $K$ -theory 246  
**Jerome Kaminker**
- 32 On the curvature of the determinant bundle 255  
**Michel Hilsum**
- 33 Minimal dynamical systems, ordered Bratteli diagrams 264  
and associated  $C^*$ -crossed products  
**Christian Skau**
- 34  $C^*$ -bundles and twisted crossed product  $C^*$ -algebras 281  
**Judith A. Packer**
- 35 Dynamical systems based on monotone complete  $C^*$ -algebras 282  
**Masamichi Hamana**
- 36 Abelian partial  $*$ -algebras 297  
— The case of normal commuting operators  
**W. Krieger**

37	Operator algebras and group duality	304
	<i>Sergio Doplicher</i>	
38	Problems related with $C^*$ -algebra approach to quantum groups	313
	<i>Stanislaw L. Woronowicz</i>	
39	Quantum deformation of the Heisenberg group	314
	<i>A. Van Daele</i>	
40	Some examples of singular foliation $C^*$ -algebras	326
	<i>Albert Jeu-Liang Sheu</i>	
41	Toeplitz algebras of Reinhardt domains	341
	— An introduction between analysis and geometry	
	<i>Albert Jeu-Liang Sheu</i>	
42	Quantum groups as objects of non-commutative geometry	359
	<i>Tetsuya Masuda and Junsei Watanabe</i>	
43	Stochastic Ising model as a $C^*$ -dynamical system	371
	<i>Taku Matsui</i>	
44	Intertwining operators, derivations of the CAR algebra, and representations of $U(p, q)$	381
	<i>Palle E. T. Jørgensen</i>	
45	Exponential rank in operator algebras	395
	<i>N. Christopher Phillips and John R. Ringrose</i>	
46	Duality for finite bipartite graphs	414
	<i>Marie Choda</i>	
47	Classification of antiautomorphisms of injective factors of type $III_\lambda$ , $\lambda \neq 0$	425
	<i>Thierry Giordano</i>	
	List of Participants	432
	List of Donors	436



# QUANTUM SYMMETRY AND SUBFACTORS

Adrian Ocneanu

Department of Mathematics

Pennsylvania State University

University Park, PA 16802, U.S.A.

## ABSTRACT

We discussed a geometric viewpoint of the invariant for inclusions of subfactors.

A family of graphs together with a biunitary connection plays the role of geometric manifold. Under very general conditions, one can construct with it an inclusion of von Neumann algebras, and the initial system is *ergodic*, i.e. the towers of commutants are obtained as finite intersections of path algebras on the graph. This makes the computation of the complete inclusion invariants algorithmic.

We defined a general *Fourier transform* on the invariant and showed that intermediate subfactors correspond precisely to projections for which the Fourier transform is, up to a scalar, also a projection.

We showed that the invariant for the central sequence inclusion associated to an inclusion  $N \subset M$  yields exactly the *fusion graph* of the bimodules for the initial factor.

# Group Actions on Injective Factors

YASUYUKI KAWAHIGASHI

*Department of Mathematics, Faculty of Science  
University of Tokyo, Hongo, Tokyo, 113, JAPAN*

**Abstract.** We survey recent progress in classification of group actions on injective factors. We deal with three kinds of groups; discrete groups with Kazhdan's property T, the real number group, and compact abelian groups. For the first and second groups, actions on the injective factor of type  $II_1$  are considered, and for the last one, we give a complete classification on injective factors of type III.

## 1. Introduction

Since Connes' breakthrough<sup>9,11</sup> for classification of integer actions on the injective factor of type  $II_1$ , several authors have studied classification problem of group actions up to (cocycle/stable) conjugacy on injective factors. As Connes' original research was tightly connected to structure of injective factors of type  $III_\lambda$ ,  $0 < \lambda < 1$ , automorphism approach for studying von Neumann algebras has been one of the most powerful and fruitful ones. Here we present recent progress in this area. We deal with three topics; actions of discrete groups with Kazhdan's property T on the injective factor  $\mathcal{R}$  of type  $II_1$ ,<sup>25</sup> one-parameter automorphism groups of  $\mathcal{R}$ ,<sup>21,22,23,24</sup> and compact abelian group actions on injective factors of type III.<sup>26,27</sup>

Jones<sup>16</sup> and Ocneanu<sup>29</sup> followed Connes and extended classification to finite and discrete amenable group actions on the injective factor  $\mathcal{R}$  of type  $II_1$ , respectively. The most important result in their work is that an outer action of a discrete amenable group  $G$  on  $\mathcal{R}$  is unique up to cocycle conjugacy. Here cocycle conjugacy means that in addition to conjugacy, we allow a perturbation of an action  $\alpha$  by a unitary cocycle  $u_g$  with  $u_{gh} = u_g \alpha_g(u_h)$ ,  $g, h \in G$ . Later, Jones showed that this uniqueness characterizes discrete amenable groups<sup>17,18</sup>; if a discrete group  $G$  is non-amenable, then it has two mutually non-cocycle conjugate outer actions on  $\mathcal{R}$ . He used ergodicity on the central sequence algebra to distinguish two actions.

Because discrete groups with Kazhdan's property  $T^{20,43}$  are known to be opposite extremes of amenable groups, we expect that the contrast also holds in group actions. We show that this is the case by showing that 1-cohomology for actions of groups with Kazhdan's property  $T$  "almost" vanishes under a certain condition. This approach also characterizes Kazhdan's property  $T$  in terms of cohomology of actions on factors of type  $II_1$ . These will be explained in Section 1.

Section 2 is devoted to one-parameter automorphism groups. Compact abelian group actions on injective factors of type  $II$  are classified by Jones-Takesaki<sup>19</sup> by reducing the problem to discrete abelian group actions on semifinite injective von Neumann algebras, which are not necessarily factors, via Takesaki duality.<sup>39</sup> Now the most important group, for still unsolved classification problem of actions on the injective factor of type  $II_1$ , is the real number group  $\mathbb{R}$ . Study of one-parameter automorphism groups of injective factors of type  $II$  is related to that of injective factor of type  $III_\lambda$ , as in the relation of automorphisms of injective factors of type  $II$  and injective factors of type  $III_\lambda$ ,  $0 < \lambda < 1$ . Though uniqueness of the injective factor of type  $III_1$  was finally established by Connes and Haagerup,<sup>13,15</sup> classification of one-parameter automorphism groups of injective factors of type  $II$  would deepen the understanding of its structure. For example, as Connes' method<sup>9</sup> is useful for Loi's study of subfactors of injective factors of type  $III_\lambda$ ,  $0 < \lambda < 1$ ,<sup>28</sup> it would give much information for study of subfactors of the injective factor of type  $III_1$ . Though complete classification has not been obtained, certain progress for several interesting cases has been made. This will be presented in Section 2.

We show a complete classification of compact abelian group actions on injective factors of type  $III$  in Section 3. This is a joint work with M. Takesaki. Sutherland and Takesaki classified discrete amenable group actions on injective factors of type  $III_\lambda$ ,  $0 < \lambda < 1$ ,<sup>38</sup> and these two and the author classified discrete abelian group actions on injective factors of type  $III_1$ .<sup>26</sup> Since abelian groups are amenable, we should be able to use these results to obtain a classification of compact abelian group actions on injective factors of type  $III$  via Takesaki duality as in work of Jones-Takesaki for semifinite cases.<sup>19</sup> However two kinds of new difficulty arise in type  $III$  situation. One is caused by the fact that each fibre in the central decomposition of an injective von Neumann algebra of type  $III$  is not necessarily mutually isomorphic even when we have a centrally ergodic action on it. The other arises from the difference of centrally trivial automorphisms and inner automorphisms. We show in Section 3 that the difficulties can be overcome. Because the classification of Sutherland, Takesaki and the author for discrete groups is up to cocycle conjugacy, we get a classification up to stable conjugacy at first. (Two actions are said to be stably conjugate if their second dual actions are conjugate.) Stable conjugacy is weaker than cocycle conjugacy, but the two notions coincide when the factor is properly infinite, hence we get a cocycle conjugacy classification for injective factors of type  $III$ , and after it, we get a complete classification (up to conjugacy) by means of inner invariant as in Jones-Takesaki.<sup>19</sup>

## 2. Actions of groups with Kazhdan's property T

Significance of Kazhdan's property  $T^{20}$  in theory of operator algebras was first pointed out by Connes<sup>12</sup>. He showed a group von Neumann algebra of a discrete ICC group with Kazhdan's property T has a countable fundamental group by rigidity argument. Since then, several authors have studied property T in operator algebra situations and in ergodic theory.<sup>35,42,43</sup> Roughly speaking, rigidity argument shows two mathematical objects are "equivalent" if they are "close" in a certain sense.

We would like to apply this type of argument to unitary cocycles. Our basic observation is that we have the following correspondence in the study of automorphism groups and unitary cocycles:

Automorphism	$\longleftrightarrow$	Cocycle
Inner automorphism	$\longleftrightarrow$	Coboundary
Factor	$\longleftrightarrow$	Ergodic action

Thus for an ergodic action  $\alpha$  of a discrete group with Kazhdan's property T on a factor of type  $II_1$ , we consider  $\alpha$ -unitary cocycles  $u_g$  with  $\text{Ad}(u_g) \cdot \alpha_g$  is also ergodic, then the number of cohomology classes of these can be shown to be at most countable. This means the number of unitary cocycles connecting ergodic actions is "small". Thus if we have a continuous family of mutually non-conjugate outer ergodic actions on the injective factor  $\mathcal{R}$  of type  $II_1$ , then their cocycle conjugacy classes have the cardinality continuum, which is the opposite extreme of uniqueness of outer actions for discrete amenable groups. Here note that the possible maximal cardinality of cocycle conjugacy classes is always continuum. (See Popa's articles<sup>31,32</sup> for related results and problems.) Groups  $SL(n, \mathbb{Z})$ ,  $n \geq 3$ , and  $Sp(n, \mathbb{Z})$ ,  $n \geq 2$ , are typical ones with Kazhdan's property T, and M. Choda<sup>5,6</sup> showed that these have a continuous family of outer ergodic actions on  $\mathcal{R}$ . (Her construction uses the irrational rotation algebra  $A_\theta$  for different  $\theta$ .) Thus we get the following.<sup>25</sup>

**THEOREM.** *Each of  $SL(n, \mathbb{Z})$ ,  $n \geq 3$ , and  $Sp(n, \mathbb{Z})$ ,  $n \geq 2$ , has a continuous family of mutually non-cocycle conjugate ergodic outer actions on the injective factor of type  $II_1$ .*

Each of these actions has ergodicity at infinity in the sense of Jones<sup>18</sup> and the crossed product algebra by each action has property T as shown by M. Choda.<sup>6</sup> Thus these actions cannot distinguished by these properties.

It may seem artificial that we used only cocycles  $u_g$  with  $\text{Ad}(u_g) \cdot \alpha_g$  ergodic in the above argument. But without this restriction, the conclusion is invalid, and these cocycles make a closed subset in the set of all the cocycles. This trouble comes from non-commutativity of the algebra, which prevents unitary cocycles from making a group. These would justify the above restriction. We can also show the fixed point algebra of  $\text{Ad}(u_g) \cdot \alpha_g$  continuously depends on  $u_g$  as follows.<sup>25</sup>

**PROPOSITION.** Let  $\alpha$  be an action of a discrete group  $G$  with Kazhdan's property T on a factor  $\mathcal{M}$  of type  $II_1$ . Then the correspondence from a unitary  $\alpha$ -cocycle  $u_g$  to the fixed point algebra  $\mathcal{M}^{Ad(u_g) \cdot \alpha_g}$  is uniformly continuous in the following sense: For any  $\varepsilon_0 > 0$ , there exists  $\delta > 0$  such that if  $d(u_g, v_g) < \delta$ , then  $\|\mathcal{M}^{Ad(u_g) \cdot \alpha_g} - \mathcal{M}^{Ad(v_g) \cdot \alpha_g}\|_2 < \varepsilon_0$ .

See Christensen<sup>7</sup> for a notation. Here the metric  $d$  is defined by  $d(u_g, v_g) = \sum_{n=1}^{\infty} \frac{1}{2^n} \|u_{g_n} - v_{g_n}\|_2$ . This shows another opposite extreme of amenable group actions. (Ocneanu's theorem<sup>29</sup> shows that an ergodic outer action of discrete amenable groups on  $\mathcal{R}$  can be perturbed to an outer action with a large fixed point algebra with an arbitrarily small cocycle.)

This kind of cohomological consideration also characterizes Kazhdan's property T, which extends a result of Araki and M. Choda<sup>2</sup> as follows.<sup>25</sup>

**THEOREM.** Let  $G$  be a discrete group. Then the following conditions are equivalent.

- (1)  $G$  has Kazhdan's property T.
- (2) Any action of  $G$  on a factor of type  $II_1$  is strong in the sense of Araki-Choda.<sup>2</sup>
- (3) For any action  $\alpha$  of  $G$  on a factor of type  $II_1$ , each cohomology class is closed.
- (4) For any action  $\alpha$  of  $G$  on a factor of type  $II_1$ , the 1-cohomology space with the quotient topology is Hausdorff.

Amenability can be also characterized in terms of cohomological terms of group actions in a similar way to Schmidt's result<sup>34</sup> as follows.<sup>25</sup>

**THEOREM.** Let  $G$  be a countable group. Then the following conditions are equivalent.

- (1)  $G$  is amenable.
- (2) No free action of  $G$  on the injective factor  $\mathcal{R}$  of type  $II_1$  is strong in the sense of Araki-Choda.<sup>2</sup>
- (3) For all outer actions of  $G$  on the injective factor  $\mathcal{R}$  of type  $II_1$ , coboundaries make a dense subset of the set of all the cocycles, but these two sets do not coincide.

### 3. One-parameter automorphism groups

First recall that classification of one-parameter automorphism groups of injective factors of type  $II$  is related to classification of injective factor of type  $III_1$  in the sense that uniqueness of the injective factor of type  $III_1$  is equivalent to uniqueness of a one-parameter automorphism group  $\alpha$  of the injective factor of type  $II_{\infty}$  with  $\text{tr} \cdot \alpha_t = e^{-t} \text{tr}$ . (See Takesaki.<sup>39</sup>) Thus classification of trace-preserving cases is still open. We present a recent progress in classification of one-parameter automorphism groups of the injective factor of type  $II_1$  here, though the complete classification has not been obtained yet.

All the work of Connes, Jones and Ocneanu<sup>9,16,29</sup> showed that a free action of a certain group on the injective factor  $\mathcal{R}$  of type  $II_1$  is unique up to cocycle conjugacy. In this context, a free action means an action  $\alpha$  for which  $\alpha_g$  is outer for all  $g \neq 1$ . Then what does a "free" action mean for the real number group? It is easy to see that for a one-parameter automorphism  $\alpha_t$ , the condition that  $\alpha_t$  is outer for all  $t \neq 0$  does not imply that the crossed product algebra is a factor. This is an analogue of the fact that  $T(\mathcal{M}) = \{0\}$  does not imply  $S(\mathcal{M}) = [0, \infty)$  for a factor  $\mathcal{M}$  of type  $III$ . (See Connes<sup>7</sup> for  $T$ -sets and  $S$ -sets.) Hence we use Connes spectrum  $\Gamma(\alpha)$  as an analogue of  $S$ -sets. But it turns out that unlike type  $III$  factor cases,  $\Gamma(\alpha) = \mathbb{R}$  does not imply that  $\alpha_t$  is outer for all  $t \neq 0$ ,<sup>21</sup> thus in order to get a uniqueness result we would like to assume both  $\Gamma(\alpha) = \mathbb{R}$ , which is equivalent to saying that the crossed product algebra is a factor, and  $\alpha_t \notin \text{Int}(\mathcal{R})$  for all  $t \neq 0$ . We remark that these two are implied by  $(\mathcal{R} \rtimes_{\alpha} \mathbb{R}) \cap \mathcal{R}' = \mathbb{C}$ . (Note that if  $\Gamma(\alpha) \neq \mathbb{R}$ , a one-parameter automorphism group  $\alpha$  can be classified more easily.<sup>21,22</sup>)

A splitting of a model action from a given free action was a key step in work of all of Connes, Jones, and Ocneanu. We next discuss what the model action is for the real number group. Ocneanu's model<sup>29</sup> is obtained by an infinite tensor product of an almost left regular representations on subsets of the group. But we use an analogue of modular automorphism groups for one-parameter automorphism groups of  $\mathcal{R}$ , and choose the following as the model action.

$$\bigotimes_{n=1}^{\infty} \text{Ad exp } it \begin{pmatrix} -\lambda/2 & 0 \\ 0 & \lambda/2 \end{pmatrix} \otimes \text{Ad exp } it \begin{pmatrix} -\mu/2 & 0 \\ 0 & \mu/2 \end{pmatrix}.$$

Here  $\lambda$  and  $\mu$  are non-zero real numbers with  $\lambda/\mu \notin \mathbb{Q}$ . This is an analogue of the ITPFI factor of type  $III_1$  of Araki-Woods,<sup>3</sup> and we can show the cocycle conjugacy class of this action does not depend on the choice of  $\lambda$  and  $\mu$ , and moreover an infinite tensor product type actions of any size of matrix gives the same cocycle conjugacy class if it has full Connes spectrum  $\mathbb{R}$  as follows.<sup>22</sup>

**THEOREM.** *If a one-parameter automorphism group  $\alpha$  of the injective factor  $\mathcal{R}$  of type  $II_1$  fixes a Cartan subalgebra of  $\mathcal{R}$  elementwise and  $\Gamma(\alpha) = \mathbb{R}$ , then its cocycle conjugacy class is unique.*

This is an analogue of uniqueness of the ITPFI (or Krieger) factor of type  $III_1$ .<sup>3</sup> This theorem is also valid for any locally compact abelian separable group action.

Our approach to the splitting problem of a model action from a "free" one-parameter automorphism  $\alpha$  of the injective factor  $\mathcal{R}$  of type  $II_1$  is as follows.

**APPROACH.** Find a weakly dense  $C^*$ -algebra  $A$  of  $\mathcal{R}$  on which  $\alpha$  is a norm continuous action, and a norm-dense  $*$ -subalgebra  $A^{\infty}$  of  $C^{\infty}$  elements in  $A$  with respect to  $\alpha$  and  $C^*$ -norm with the following properties:

(1) For any  $\varepsilon > 0$ , there exists a  $2 \times 2$  matrix unit  $(e_{jk})$  of  $A_{e_{11}+e_{22}}$  in  $A^{\infty}$  with  $\text{tr}(1 - e_{11} - e_{22}) < \varepsilon$ .



(2) For a given  $\nu \in \mathbf{R}$  and  $x_1, \dots, x_n \in A^\infty$  and a matrix unit  $(e_{jk})$  in (1), there exists an automorphism  $\sigma$  of  $A^\infty$  such that  $\|\delta(\sigma(e_{jk})) - i|j - k|\nu e_{jk}\| < \varepsilon$  and  $\|\sigma(e_{jk}), x_l\| < \varepsilon$ , where  $\delta$  is the derivation generating  $\alpha$ .

In general, it is too much that we expect existence of a matrix unit in  $A$ , because of  $K$ -theoretic restrictions. Thus we adopt a weaker version (1) as above. The second estimate in (2) is for making a central sequence, and related to asymptotical abelian systems.<sup>35</sup> The first estimate in (2) is for making a piece of the model action. Condition (2) is similar to Araki's property  $L'_\lambda$ ,<sup>1</sup> and a key tool for producing the model action as Ocneanu's non-commutative Rohlin theorem.<sup>29</sup> By working on several estimates using (1), (2), we can show a splitting of the model action in terms of central sequences.<sup>23</sup>

But unfortunately, it is unknown how to produce  $A^\infty$  and  $\sigma$  in a general setting, and we are lead to working on a concrete example in detail. A key to the uniqueness result cited above was abundant existence of elements fixed by  $\alpha$ . From this point, the opposite extreme is an ergodic action. Now we deal with the following construction of an action.

In the irrational rotation  $C^*$ -algebra  $A_\theta$  with  $uv = e^{2\pi i\theta}vu$ , consider the following one-parameter automorphism group  $\alpha_t$ :  $\alpha_t(u) = e^{i\lambda t}u$ ,  $\alpha_t(v) = e^{i\mu t}v$ . Here  $\lambda$  and  $\mu$  are non-zero real numbers with  $\lambda/\mu \notin \mathbf{Q}$ . We extend this one-parameter automorphism group to the weak closure  $\mathcal{R}$  of  $A_\theta$  with respect to the trace  $\tau$ , which is the injective  $\text{II}_1$  factor. It is easy to see this action has full Connes spectrum  $\mathbf{R}$  but this is outer if and only if  $\lambda/\mu$  is not in the  $GL(2, \mathbf{Q})$  orbit of  $\theta$ , where a  $GL(2, \mathbf{Q})$ -action is given by a fractional transformation. This subtlety is another interesting point of this action. We would like to show that this action is cocycle conjugate to the model action. Though this action is almost periodic (i.e., this action can be extended to a  $\mathbf{T}^2$  action), it is impossible to perturb this  $\mathbf{T}^2$  action to a product type one, by Olesen-Pedersen-Takesaki.<sup>30</sup> It makes this action the best one for trying our approach. For this action, we use a Rieffel projection construction<sup>33</sup> for (1) in the above approach and an  $SL(2, \mathbf{Z})$  action on  $A_\theta$  considered by Brenken<sup>4</sup> and Watatani<sup>41</sup> for (2) in the approach. Then a delicate estimate shows the condition  $\lambda/\mu \notin GL(2, \mathbf{Q})\theta$  exactly implies (2). Hence we get a splitting of the model action. Then we can use the following theorem to show the model action "absorbs" our action  $\alpha$ .<sup>23</sup>

**THEOREM.** Let  $\alpha$  be an almost periodic one-parameter automorphism group of  $\mathcal{R}$ . Let  $\beta_t$  be an infinite tensor product type one-parameter automorphism group of  $\mathcal{R}$  with  $\Gamma(\beta) = \mathbf{R}$ . Then  $\alpha_t \otimes \beta_t$  is cocycle conjugate to the  $\beta_t$  on  $\mathcal{R}$ .

Hence we get the following.<sup>23</sup>

**THEOREM.** Let  $\alpha$  be the one-parameter automorphism as above. We assume  $\lambda/\mu \notin \mathbf{Q}$ , and  $\lambda/\mu$  is not in the  $GL(2, \mathbf{Q})$  orbit of  $\theta$ . Then  $\alpha_t$  is cocycle conjugate to an infinite tensor product type action  $\beta$  of  $\mathbf{R}$  on  $\mathcal{R}$  with  $\Gamma(\beta) = \mathbf{R}$ , which is unique up to cocycle conjugacy.

Certainly it is an open interesting problem how far our approach can go in general.

#### 4. Compact abelian group actions

After Sutherland and Takesaki classified discrete amenable group actions on injective factors of type  $\text{III}_\lambda$ ,  $0 \leq \lambda < 1$ , up to cocycle conjugacy,<sup>38</sup> they and the author extended the same type of classification to discrete abelian group actions on the injective factor of type  $\text{III}_1$ .<sup>26</sup> In the both work, we have to invoke a result of Ocneanu<sup>29</sup> at a certain stage. Because the classes of approximately inner automorphisms and centrally trivial automorphisms play an important role in Ocneanu's theorem as in Connes' work,<sup>9</sup> we need characterizations for these two classes of automorphisms on injective factors of type  $\text{III}$ . Connes announced these characterizations in 1976<sup>10</sup> as follows, but the proof has not been available for a long time, though this theorem has been used in Connes<sup>13</sup> and Sutherland-Takesaki.<sup>38</sup>

**THEOREM.** *For injective factors  $\mathcal{M}$  of type  $\text{III}$ , we have:*

- (i)  $\text{Ker}(\text{mod}) = \overline{\text{Int}}(\mathcal{M})$ ;
- (ii) *An automorphism  $\alpha$  of  $\mathcal{M}$  is centrally trivial if and only if  $\alpha$  is of the form  $\alpha = \text{Ad}(u) \cdot \bar{\sigma}_c^\varphi$ , where  $\bar{\sigma}_c^\varphi$  is an extended modular automorphism for a dominant weight  $\varphi$  on  $\mathcal{M}$ ,  $c$  is a  $\theta$ -cocycle on  $\mathcal{U}(\mathcal{C}_\varphi)$ , and  $u \in \mathcal{U}(\mathcal{M})$ .*

(See Connes-Takesaki<sup>14</sup> for related definitions and notations.)

Sutherland, Takesaki and the author supplied a proof of the above theorem for the first time,<sup>26</sup> and made use of the characterizations. After Ocneanu's work,<sup>29</sup> Jones and Takesaki<sup>19</sup> classified compact abelian group actions on injective factors of type  $\text{II}$  via Takesaki duality. They work on dual actions, which no more act on factors in general, and regard the actions as groupoid actions on factors. It is now natural to expect the same approach extends in type  $\text{III}$  settings. We show that this is the case, but several kinds of technical difficulty arise from the fact that the algebra is of type  $\text{III}$ .<sup>27</sup>

For a given action  $\alpha$  of a separable compact abelian group  $A$  on an injective factor  $\mathcal{M}$  of type  $\text{III}$ , we work on its dual action on  $\mathcal{M} \rtimes_\alpha A$ . We would like to represent the crossed product algebra as  $\mathcal{N} \bar{\otimes} L^\infty(X, \mu)$  for a certain injective factor  $\mathcal{N}$  and work on a groupoid  $X \rtimes \hat{A}$  to use tools like Krieger's cohomology lemma as in Jones-Takesaki.<sup>19</sup> But this type of central decomposition is impossible in general because classification of type  $\text{III}_0$  factors is not smooth. That is, each fibre factor in the central decomposition is not mutually isomorphic in general. This phenomenon actually occurs in this crossed product algebra setting as shown in our work.<sup>27</sup> Hence we have to take a more technical version of groupoid approach, and will use Sutherland's generalized cohomology lemma<sup>36</sup> instead of Krieger's. Then we can apply the groupoid approach of Sutherland-Takesaki.<sup>37</sup> The next difficulty arises from the fact that the class of centrally trivial automorphisms is now strictly larger than the class of inner automorphisms. For example, we have to prove that in the central decomposition  $\mathcal{M} \rtimes_\alpha A = \int_X^\oplus \mathcal{N}(x) d\mu(x)$ , the correspondence  $x \mapsto$



$\text{Cnt}(\mathcal{N}(x))$  is Borel in a certain sense defined by Sutherland.<sup>36</sup> This can be done since we have a complete characterization of  $\text{Cnt}(\mathcal{N}(x))$ .<sup>26</sup>

An outline of the arguments is as follows. First, consider two extreme cases. In one case, we have a factor crossed product algebra  $\mathcal{M} \rtimes_{\alpha} A$ . This type of action is called prime, and we have a full Connes spectrum,  $\Gamma(\alpha) = \hat{A}$ , in this case. Then we have invariants as in Sutherland-Takesaki<sup>38</sup> for the dual action on the factor  $\mathcal{M} \rtimes_{\alpha} A$ . In the other extreme case, we have a zero Connes spectrum,  $\Gamma(\alpha) = \{0\}$ . Then we have an (ergodic) faithful action of  $\hat{A}$  on the center  $\mathcal{Z}(\mathcal{M} \rtimes_{\alpha} A)$  and the module given by the dual action, as invariants. (The action of  $\hat{A}$  is like a flow of weights, and the module is also a part of invariants in the first case.)

In general, the situation is a mixture of these two, but no new invariants appear, and these are all we need. The first family of invariants are handles by a classification of Sutherland-Takesaki,<sup>38</sup> and these two and the author.<sup>26</sup> We use Sutherland's generalized cohomology lemma<sup>36</sup> to deal with the second family of invariants. For the cohomology lemma, we need a certain density argument. For example, in the easiest situation, we need  $\text{Ker}(\text{mod}) = \bar{\text{Int}}$ , and here we need more delicate result. The following is a key to it.

**THEOREM.** *Let  $\mathcal{M}$  be an injective factor of type III, and  $\varphi$  be a dominant weight on  $\mathcal{M}$ . If an automorphism  $\alpha$  of  $\mathcal{M}$  is approximately inner, then there exist a unitary  $u \in \mathcal{M}$  and a sequence of unitaries  $\{v_n\}$  in  $\mathcal{M}_{\varphi}$  such that  $\alpha = \text{Ad}(u) \cdot \lim_{n \rightarrow \infty} \text{Ad}(v_n)$ .*

Here inner perturbation by  $u$  does not matter very much, thus this essentially means that approximately inner automorphisms are approximated by unitaries in the centralizer. For the proof of the above theorem, we need some technical lemmas, but we have a complete list of automorphisms of injective factors of type III,<sup>26,38</sup> hence it is enough to check the conclusion for all the members of the list.

Then we have to combine results for  $\Gamma(\alpha)$  and  $\hat{A}/\Gamma(\alpha)$  to supply a complete cocycle. This can be done by constructing a groupoid model action on the injective factor  $\mathcal{R}$  of type II<sub>1</sub> which can kill any bicharacter on the groupoid times a subgroup of  $\hat{A}$ . Because all the fibre factors in the central decomposition split  $\mathcal{R}$  as a tensor product factor, this is meaningful. Then the error from being a cocycle is given by a certain bicharacter, and it can be killed by a good property of the model action. The final form is as follows.<sup>27</sup>

**THEOREM.** *Let  $\alpha$  and  $\beta$  be actions of a compact abelian group  $A$  on an injective factor  $\mathcal{M}$  of type III. Then we conclude:*

- i)  $\alpha$  and  $\beta$  are cocycle conjugate if and only if
  - a)  $\mathcal{M} \rtimes_{\alpha} A \cong \mathcal{M} \rtimes_{\beta} A$ ,
  - b)  $N(\hat{\alpha}) = N(\hat{\beta})$ ,
  - c) there exists an isomorphism  $\theta$  of  $\mathcal{F}(\mathcal{M} \rtimes_{\alpha} A)$  onto  $\mathcal{F}(\mathcal{M} \rtimes_{\beta} A)$  which conjugates the restriction of  $\hat{\alpha}$  and  $\hat{\beta}$  to the center  $\mathcal{Z}(\mathcal{M} \rtimes_{\alpha} A)$  and  $\mathcal{Z}(\mathcal{M} \rtimes_{\beta} A)$  and  $\text{mod}(\hat{\alpha}, \chi_{\hat{\alpha}}, \nu_{\alpha}) = (\text{mod}(\hat{\beta}), \chi_{\hat{\beta}}, \nu_{\beta})$ .