

美国数学会经典影印系列



Fourier Analysis

傅里叶分析

Javier Duoandikoetxea

Translated and revised by
David Cruz-Uribe, SFO



高等教育出版社

美国数学会经典影印系列



Fourier Analysis

傅里叶分析

Javier Duoandikoetxea

Translated and revised by

David Cruz-Urbe, SFO



高等教育出版社·北京

图字: 01-2016-2507 号

Fourier Analysis, by Javier Duoandikoetxea, Translated and revised by David Cruz-Urbe, SFO,
first published by the American Mathematical Society.

Copyright © 2001 by the American Mathematical Society. All rights reserved.

This present reprint edition is published by Higher Education Press Limited Company under authority
of the American Mathematical Society and is published under license.

Special Edition for People's Republic of China Distribution Only. This edition has been authorized by
the American Mathematical Society for sale in People's Republic of China only, and is not for export therefrom.

本书最初由美国数学会于 2001 年出版, 原书名为 *Fourier Analysis*, 作者为 Javier Duoandikoetxea,
由 David Cruz-Urbe, SFO 翻译和修订。美国数学会保留原书所有版权。

原书版权声明: Copyright © 2001 by the American Mathematical Society.

本影印版由高等教育出版社有限公司经美国数学会独家授权出版。

本版只限于中华人民共和国境内发行。本版经由美国数学会授权仅在中华人民共和国境内销售, 不得出口。

傅里叶分析

Fuliye Fenxi

图书在版编目 (CIP) 数据

傅里叶分析 = *Fourier Analysis*: 英文 / (西) 贾维尔著.
—影印本. —北京: 高等教育出版社, 2017.1

ISBN 978-7-04-046901-1

I. ①傅… II. ①贾… III. ①傅里叶分析—英文 IV.
①O174.2

中国版本图书馆 CIP 数据核字 (2016) 第 281767 号

策划编辑 赵天夫 责任编辑 赵天夫
封面设计 张申申 责任印制 毛斯璐

出版发行 高等教育出版社
社址 北京市西城区德外大街4号
邮政编码 100120
购书热线 010-58581118
咨询电话 400-810-0598
网址 <http://www.hep.edu.cn>

<http://www.hep.com.cn>
网上订购 <http://www.hepmall.com.cn>
<http://www.hepmall.com>
<http://www.hepmall.cn>
印刷: 北京新华印刷有限公司

开本 787mm×1092mm 1/16
印张 15.25
字数 310千字
版次 2017年1月第1版
印次 2017年1月第1次印刷
定价 99.00元

本书如有缺页、倒页、脱页等质量问题,
请到所购图书销售部门联系调换
版权所有 侵权必究
[物料号 46901-00]



美国数学会经典影印系列

出版者的话

近年来,我国的科学技术取得了长足进步,特别是在数学等自然科学基础领域不断涌现出一流的研究成果。与此同时,国内的科研队伍与国外的交流合作也越来越密切,越来越多的科研工作者可以熟练地阅读英文文献,并在国际顶级期刊发表英文学术文章,在国外出版社出版英文学术著作。

然而,在国内阅读海外原版英文图书仍不是非常便捷。一方面,这些原版图书主要集中在科技、教育比较发达的大中城市的大型综合图书馆以及科研院所的资料室中,普通读者借阅不甚容易;另一方面,原版书价格昂贵,动辄上百美元,购买也很不方便。这极大地限制了科技工作者对于国外先进科学技术知识的获取,间接阻碍了我国科技的发展。

高等教育出版社本着植根教育、弘扬学术的宗旨服务我国广大科技和教育工作者,同美国数学会(American Mathematical Society)合作,在征求海内外众多专家学者意见的基础上,精选该学会近年出版的数十种专业著作,组织出版了“美国数学会经典影印系列”丛书。美国数学会创建于1888年,是国际上极具影响力的专业学术组织,目前拥有近30000会员和580余个机构成员,出版图书3500多种,冯·诺依曼、莱夫谢茨、陶哲轩等世界级数学大家都是其作者。本影印系列涵盖了代数、几何、分析、方程、拓扑、概率、动力系统所有主要数学分支以及新近发展的数学主题。

我们希望这套书的出版,能够对国内的科研工作者、教育工作者以及青年学生起到重要的学术引领作用,也希望今后能有更多的海外优秀英文著作被介绍到中国。

高等教育出版社

2016年12月

*Dedicated to the memory of
José Luis Rubio de Francia, my teacher and friend,
who would have written a much better book than I have*

Preface

Fourier Analysis is a large branch of mathematics whose point of departure is the study of Fourier series and integrals. However, it encompasses a variety of perspectives and techniques, and so many different introductions with that title are possible. The goal of this book is to study the real variable methods introduced into Fourier analysis by A. P. Calderón and A. Zygmund in the 1950's.

We begin in Chapter 1 with a review of Fourier series and integrals, and then in Chapters 2 and 3 we introduce two operators which are basic to the field: the Hardy-Littlewood maximal function and the Hilbert transform. Even though they appeared before the techniques of Calderón and Zygmund, we treat these operators from their point of view. The goal of these techniques is to enable the study of analogs of the Hilbert transform in higher dimensions; these are of great interest in applications. Such operators are known as singular integrals and are discussed in Chapters 4 and 5 along with their modern generalizations. We next consider two of the many contributions to the field which appeared in the 1970's. In Chapter 6 we study the relationship between H^1 , BMO and singular integrals, and in Chapter 7 we present the elementary theory of weighted norm inequalities. In Chapter 8 we discuss Littlewood-Paley theory; its origins date back to the 1930's, but it has had extensive later development which includes a number of applications. Those presented in this chapter are useful in the study of Fourier multipliers, which also uses the theory of weighted inequalities. We end the book with an important result of the 80's, the so-called $T1$ theorem, which has been of crucial importance to the field.

At the end of each chapter there is a section in which we try to give some idea of further results which are not discussed in the text, and give

references for the interested reader. A number of books and all the articles cited appear only in these notes; the bibliography at the end of the text is reserved for books which treat in depth the ideas we have presented.

The material in this book comes from a graduate course taught at the Universidad Autónoma de Madrid during the academic year 1988-89. Part of it is based on notes I took as a student in a course taught by José Luis Rubio de Francia at the same university in the fall of 1985. It seemed to have been his intention to write up his course, but he was prevented from doing so by his untimely death. Therefore, I have taken the liberty of using his ideas, which I learned both in his class and in many pleasant conversations in the hallway and at the blackboard, to write this book. Although it is dedicated to his memory, I almost regard it as a joint work. Also, I would like to thank my friends at the Universidad Autónoma de Madrid who encouraged me to teach this course and to write this book.

The book was first published in Spanish in the *Colección de Estudios* of the Universidad Autónoma de Madrid (1991), and then was republished with only some minor typographical corrections in a joint edition of Addison-Wesley/Universidad Autónoma de Madrid (1995). From the very beginning some colleagues suggested that there would be interest in an English translation which I never did. But when Professor David Cruz-Urbe offered to translate the book I immediately accepted. I realized at once that the text could not remain the same because some of the many developments of the last decade had to be included in the informative sections closing each chapter together with a few topics omitted from the first edition. As a consequence, although only minor changes have been introduced to the core of the book, the sections named "Notes and further results" have been considerably expanded to incorporate new topics, results and references.

The task of updating the book would have not been accomplished as it has been without the invaluable contribution of Professor Cruz-Urbe. Apart from reading the text, suggesting changes and clarifying obscure points, he did a great work on expanding the above mentioned notes, finding references and proposing new results to be included. The improvements of this book with respect to the original have certainly been the fruit of our joint work, and I am very grateful to him for sharing with me his knowledge of the subject much beyond the duties of a mere translator.

Javier Duoandikoetxea
Bilbao, June 2000

Acknowledgment: The translator would like to thank the Ford Foundation and the Dean of Faculty at Trinity College for their generous support during the academic year 1998–99. It was during this year-long sabbatical that this project was conceived and the first draft of the translation produced.

Preliminaries

Here we review some notation and basic results, but we assume that they are mostly well known to the reader. For more information, see, for example, Rudin [14].

In general we will work in $\mathbb{R}^n$. The Euclidean norm will be denoted by $|\cdot|$. If $x \in \mathbb{R}^n$ and $r > 0$,

$$B(x, r) = \{y \in \mathbb{R}^n : |x - y| < r\}$$

is the ball with center x and radius r . Lebesgue measure in $\mathbb{R}^n$ is denoted by dx and on the unit sphere S^{n-1} in $\mathbb{R}^n$ by $d\sigma$. If E is a subset of $\mathbb{R}^n$, $|E|$ denotes its Lebesgue measure and χ_E its characteristic function: $\chi_E(x) = 1$ if $x \in E$ and 0 if $x \notin E$. The expressions *almost everywhere* or *for almost every* x refer to properties which hold except on a set of measure 0; they are abbreviated by “a.e.” and “a.e. x .”

If $a = (a_1, \dots, a_n) \in \mathbb{N}^n$ is a multi-index and $f : \mathbb{R}^n \rightarrow \mathbb{C}$, then

$$D^a f = \frac{\partial^{|a|} f}{\partial x_1^{a_1} \cdots \partial x_n^{a_n}},$$

where $|a| = a_1 + \cdots + a_n$ and $x^a = x_1^{a_1} \cdots x_n^{a_n}$.

Let (X, μ) be a measure space. $L^p(X, \mu)$, $1 \leq p < \infty$, denotes the Banach space of functions from X to $\mathbb{C}$ whose p -th powers are integrable; the norm of $f \in L^p(X, \mu)$ is

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{1/p}.$$

$L^\infty(X, \mu)$ denotes the Banach space of essentially bounded functions from X to $\mathbb{C}$; more precisely, functions f such that for some $C > 0$,

$$\mu(\{x \in X : |f(x)| > C\}) = 0.$$

The norm of f , $\|f\|_\infty$, is the infimum of the constants with this property. In general X will be $\mathbb{R}^n$ (or a subset of $\mathbb{R}^n$) and $d\mu = dx$; in this case we often do not give the measure or the space but instead simply write L^p . For general measure spaces we will frequently write $L^p(X)$ instead of $L^p(X, \mu)$; if μ is absolutely continuous and $d\mu = w dx$ we will write $L^p(w)$. The conjugate exponent of p is always denoted by p' :

$$\frac{1}{p} + \frac{1}{p'} = 1.$$

The triangle inequality on L^p has an integral version which we refer to as Minkowski's integral inequality and which we will use repeatedly. Given measure spaces (X, μ) and (Y, ν) with σ -finite measures, the inequality is

$$\left(\int_X \left| \int_Y f(x, y) d\nu(y) \right|^p d\mu(x) \right)^{1/p} \leq \int_Y \left(\int_X |f(x, y)|^p d\mu(x) \right)^{1/p} d\nu(y).$$

The convolution of two functions f and g defined on $\mathbb{R}^n$ is given by

$$f * g(x) = \int_{\mathbb{R}^n} f(y)g(x - y) dy = \int_{\mathbb{R}^n} f(x - y)g(y) dy$$

whenever this expression makes sense.

The spaces of test functions are $C_c^\infty(\mathbb{R}^n)$, the space of infinitely differentiable functions of compact support, and $\mathcal{S}(\mathbb{R}^n)$, the so-called Schwartz functions. A Schwartz function is an infinitely differentiable function which decreases rapidly at infinity (more precisely, the function and all its derivatives decrease more rapidly than any polynomial increases). Given the appropriate topologies, their duals are the spaces of distributions and tempered distributions. It makes sense to define the convolution of a distribution and a test function as follows: if $T \in C_c^\infty(\mathbb{R}^n)'$ and $f \in C_c^\infty(\mathbb{R}^n)$, then

$$T * f(x) = \langle T, \tau_x \tilde{f} \rangle,$$

where $\tilde{f}(y) = f(-y)$ and $\tau_x f(y) = f(x + y)$. Note that this definition coincides with the previous one if T is a locally integrable function. Similarly, we can take $T \in \mathcal{S}(\mathbb{R}^n)'$ and $f \in \mathcal{S}(\mathbb{R}^n)$. We denote the duality by either $\langle T, f \rangle$ or $T(f)$ without distinction.

References in square brackets are to items in the bibliography at the end of the book.

Finally, we remark that C will denote a positive constant which may be different even in a single chain of inequalities.

Contents

Preface	xiii
Preliminaries	xvii
Chapter 1. Fourier Series and Integrals	1
§1. Fourier coefficients and series	1
§2. Criteria for pointwise convergence	2
§3. Fourier series of continuous functions	6
§4. Convergence in norm	8
§5. Summability methods	9
§6. The Fourier transform of L^1 functions	11
§7. The Schwartz class and tempered distributions	12
§8. The Fourier transform on L^p , $1 < p \leq 2$	15
§9. The convergence and summability of Fourier integrals	17
§10. Notes and further results	19
Chapter 2. The Hardy-Littlewood Maximal Function	25
§1. Approximations of the identity	25
§2. Weak-type inequalities and almost everywhere convergence	26
§3. The Marcinkiewicz interpolation theorem	28
§4. The Hardy-Littlewood maximal function	30

§5. The dyadic maximal function	32
§6. The weak $(1, 1)$ inequality for the maximal function	35
§7. A weighted norm inequality	37
§8. Notes and further results	38
Chapter 3. The Hilbert Transform	49
§1. The conjugate Poisson kernel	49
§2. The principal value of $1/x$	50
§3. The theorems of M. Riesz and Kolmogorov	51
§4. Truncated integrals and pointwise convergence	55
§5. Multipliers	58
§6. Notes and further results	61
Chapter 4. Singular Integrals (I)	69
§1. Definition and examples	69
§2. The Fourier transform of the kernel	70
§3. The method of rotations	73
§4. Singular integrals with even kernel	77
§5. An operator algebra	80
§6. Singular integrals with variable kernel	83
§7. Notes and further results	85
Chapter 5. Singular Integrals (II)	91
§1. The Calderón-Zygmund theorem	91
§2. Truncated integrals and the principal value	94
§3. Generalized Calderón-Zygmund operators	98
§4. Calderón-Zygmund singular integrals	101
§5. A vector-valued extension	105
§6. Notes and further results	107
Chapter 6. H^1 and BMO	115
§1. The space atomic H^1	115
§2. The space BMO	117

§3. An interpolation result	121
§4. The John-Nirenberg inequality	123
§5. Notes and further results	126
Chapter 7. Weighted Inequalities	133
§1. The A_p condition	133
§2. Strong-type inequalities with weights	137
§3. A_1 weights and an extrapolation theorem	140
§4. Weighted inequalities for singular integrals	143
§5. Notes and further results	147
Chapter 8. Littlewood-Paley Theory and Multipliers	157
§1. Some vector-valued inequalities	157
§2. Littlewood-Paley theory	159
§3. The Hörmander multiplier theorem	163
§4. The Marcinkiewicz multiplier theorem	166
§5. Bochner-Riesz multipliers	168
§6. Return to singular integrals	172
§7. The maximal function and the Hilbert transform along a parabola	178
§8. Notes and further results	184
Chapter 9. The $T1$ Theorem	195
§1. Cotlar's lemma	195
§2. Carleson measures	197
§3. Statement and applications of the $T1$ theorem	201
§4. Proof of the $T1$ theorem	205
§5. Notes and further results	212
Bibliography	217
Index	219

Fourier Series and Integrals

1. Fourier coefficients and series

The problem of representing a function f , defined on (an interval of) $\mathbb{R}$, by a trigonometric series of the form

$$(1.1) \quad f(x) = \sum_{k=0}^{\infty} a_k \cos(kx) + b_k \sin(kx)$$

arises naturally when using the method of separation of variables to solve partial differential equations. This is how J. Fourier arrived at the problem, and he devoted the better part of his *Théorie Analytique de la Chaleur* (1822, results first presented to the Institute de France in 1807) to it. Even earlier, in the middle of the 18th century, Daniel Bernoulli had stated it while trying to solve the problem of a vibrating string, and the formula for the coefficients appeared in an article by L. Euler in 1777.

The right-hand side of (1.1) is a periodic function with period 2π , so f must also have this property. Therefore it will suffice to consider f on an interval of length 2π . Using Euler's identity, $e^{ikx} = \cos(kx) + i \sin(kx)$, we can replace the functions $\sin(kx)$ and $\cos(kx)$ in (1.1) by $\{e^{ikx} : k \in \mathbb{Z}\}$; we will do so from now on. Moreover, we will consider functions with period 1 instead of 2π , so we will modify the system of functions to $\{e^{2\pi i k x} : k \in \mathbb{Z}\}$. Our problem is thus transformed into studying the representation of f by

$$(1.2) \quad f(x) = \sum_{k=-\infty}^{\infty} c_k e^{2\pi i k x}.$$

If we assume, for example, that the series converges uniformly, then by multiplying by $e^{-2\pi imx}$ and integrating term-by-term on $(0, 1)$ we get

$$c_m = \int_0^1 f(x) e^{-2\pi imx} dx$$

because of the orthogonality relationship

$$(1.3) \quad \int_0^1 e^{2\pi ikx} e^{-2\pi imx} dx = \begin{cases} 0 & \text{if } k \neq m \\ 1 & \text{if } k = m. \end{cases}$$

Denote the additive group of the reals modulo 1 (that is $\mathbb{R}/\mathbb{Z}$) by $\mathbb{T}$, the one-dimensional torus. This can also be identified with the unit circle, S^1 . Saying that a function is defined on $\mathbb{T}$ is equivalent to saying that it is defined on $\mathbb{R}$ and has period 1. To each function $f \in L^1(\mathbb{T})$ we associate the sequence $\{\hat{f}(k)\}$ of Fourier coefficients of f , defined by

$$(1.4) \quad \hat{f}(k) = \int_0^1 f(x) e^{-2\pi ikx} dx.$$

The trigonometric series with these coefficients,

$$(1.5) \quad \sum_{k=-\infty}^{\infty} \hat{f}(k) e^{2\pi ikx},$$

is called the Fourier series of f .

Our problem now consists in determining when and in what sense the series (1.5) represents the function f .

2. Criteria for pointwise convergence

Denote the N -th symmetric partial sum of the series (1.5) by $S_N f(x)$; that is,

$$S_N f(x) = \sum_{k=-N}^N \hat{f}(k) e^{2\pi ikx}.$$

Note that this is also the N -th partial sum of the series when it is written in the form of (1.1).

Our first approach to the problem of representing f by its Fourier series is to determine whether $\lim S_N f(x)$ exists for each x , and if so, whether it is equal to $f(x)$. The first positive result is due to P. G. L. Dirichlet (1829), who proved the following convergence criterion: if f is bounded, piecewise continuous, and has a finite number of maxima and minima, then $\lim S_N f(x)$ exists and is equal to $\frac{1}{2}[f(x+) + f(x-)]$. Jordan's criterion, which we prove below, includes this result as a special case.

In order to study $S_N f(x)$ we need a more manageable expression. Dirichlet wrote the partial sums as follows:

$$\begin{aligned} S_N f(x) &= \sum_{k=-N}^N \int_0^1 f(t) e^{-2\pi i k t} dt \cdot e^{2\pi i k x} \\ &= \int_0^1 f(t) D_N(x-t) dt \\ &= \int_0^1 f(x-t) D_N(t) dt, \end{aligned}$$

where D_N is the Dirichlet kernel,

$$D_N(t) = \sum_{k=-N}^N e^{2\pi i k t}.$$

If we sum this geometric series we get

$$(1.6) \quad D_N(t) = \frac{\sin(\pi(2N+1)t)}{\sin(\pi t)}.$$

This satisfies

$$\int_0^1 D_N(t) dt = 1 \quad \text{and} \quad |D_N(t)| \leq \frac{1}{\sin(\pi\delta)}, \quad \delta \leq |t| \leq 1/2.$$

We will prove two criteria for pointwise convergence.

Theorem 1.1 (Dini's Criterion). *If for some x there exists $\delta > 0$ such that*

$$\int_{|t| < \delta} \left| \frac{f(x+t) - f(x)}{t} \right| dt < \infty,$$

then

$$\lim_{N \rightarrow \infty} S_N f(x) = f(x).$$

Theorem 1.2 (Jordan's Criterion). *If f is a function of bounded variation in a neighborhood of x , then*

$$\lim_{N \rightarrow \infty} S_N f(x) = \frac{1}{2} [f(x+) + f(x-)].$$

At first it may seem surprising that these results are local, since if we modify the function slightly, the Fourier coefficients of f change. Nevertheless, the convergence of a Fourier series is effectively a local property, and if the modifications are made outside of a neighborhood of x , then the behavior of the series at x does not change. This is made precise by the following result.