

Gordon James & Martin Liebeck

Representations and Characters of Groups

Second Edition

群的表示与群的特征 第2版

CAMBRIDGE

世界图书出版公司
www.wpcbj.com.cn

0152.6/Y9

2009.

REPRESENTATIONS AND CHARACTERS OF GROUPS

GORDON JAMES AND MARTIN LIEBECK

*Department of Mathematics,
Imperial College, London*

Second Edition



CAMBRIDGE
UNIVERSITY PRESS

图书在版编目 (CIP) 数据

表示与群的特征: 第2版 = Representations and Characters of Groups 2nd ed. : 英文/ (英) 詹姆斯 (James, G.) 著.
—2 版. —北京: 世界图书出版公司北京公司, 2009. 5
ISBN 978-7-5100-0458-2

I. 表… II. 詹… III. 群表示—英文 IV. 0152.6

中国版本图书馆 CIP 数据核字 (2009) 第 055596 号

书 名:	Representations and Characters of Groups 2nd ed.
作 者:	Gordon James & Martin Liebeck
中 译 名:	群的表示与群的特征 第2版
责任编辑:	高蓉 刘慧
出 版 者:	世界图书出版公司北京公司
印 刷 者:	三河国英印务有限公司
发 行:	世界图书出版公司北京公司 (北京朝内大街 137 号 100010)
联系电话:	010-64021602, 010-64015659
电子信箱:	kjb@wpcbj.com.cn
开 本:	24 开
印 张:	19.5
版 次:	2009 年 05 月
版权登记:	图字: 01-2009-1667
书 号:	978-7-5100-0458-2/O · 673
定 价:	49.00 元

Representations and Characters of Groups, 2nd ed. (978-0-521-00392-6) by Gordon James & Martin Liebeck first published by Cambridge University Press 2001

All rights reserved.

This reprint edition for the People's Republic of China is published by arrangement with the Press Syndicate of the University of Cambridge, Cambridge, United Kingdom.

© Cambridge University Press & Beijing World Publishing Corporation 2009

This book is in copyright. No reproduction of any part may take place without the written permission of Cambridge University Press or Beijing World Publishing Corporation.

This edition is for sale in the mainland of China only, excluding Hong Kong SAR, Macao SAR and Taiwan, and may not be bought for export therefrom.

此版本仅限中华人民共和国境内销售，不包括香港、澳门特别行政区及中国台湾。不得出口。

Preface

We have attempted in this book to provide a leisurely introduction to the representation theory of groups. But why should this subject interest you?

Representation theory is concerned with the ways of writing a group as a group of matrices. Not only is the theory beautiful in its own right, but it also provides one of the keys to a proper understanding of finite groups. For example, it is often vital to have a concrete description of a particular group; this is achieved by finding a representation of the group as a group of matrices. Moreover, by studying the different representations of the group, it is possible to prove results which lie outside the framework of representation theory. One simple example: all groups of order p^2 (where p is a prime number) are abelian; this can be shown quickly using only group theory, but it is also a consequence of basic results about representations. More generally, all groups of order $p^a q^b$ (p and q primes) are soluble; this again is a statement purely about groups, but the best proof, due to Burnside, is an outstanding example of the use of representation theory. In fact, the range of applications of the theory extends far beyond the boundaries of pure mathematics, and includes theoretical physics and chemistry – we describe one such application in the last chapter.

The book is suitable for students who have taken first undergraduate courses involving group theory and linear algebra. We have included two preliminary chapters which cover the necessary background material. The basic theory of representations is developed in Chapters 3–23, and our methods concentrate upon the use of modules; although this accords with the more modern style of algebra, in several instances our proofs differ from those found in other textbooks. The main results are elegant and surprising, but at first sight they sometimes have an air of mystery

about them; we have chosen the approach which we believe to be the most transparent.

We also emphasize the practical aspects of the subject, and the text is illustrated with a wealth of examples. A feature of the book is the wide variety of groups which we investigate in detail. By the end of Chapter 28, we have presented the character tables of all groups of order less than 32, of all p -groups of order at most p^4 , and of all the simple groups of order less than 1000.

Every chapter is accompanied by a set of Exercises, and the solutions to all of these are provided at the end of the book.

We would like to thank Dr Hans Liebeck for his careful reading of our manuscript and the many helpful suggestions which he made.

Preface to Second Edition

In this second edition, we have included two new chapters; one (Chapter 28) deals with the character tables of an infinite series of groups, and the other (Chapter 29) covers aspects of the representation theory of permutation groups. We have also added a considerable amount of new material to Chapters 20, 23 and 30, and made minor amendments elsewhere.

Contents

<i>Preface</i>	<i>page</i> vii
1 Groups and homomorphisms	1
2 Vector spaces and linear transformations	14
3 Group representations	30
4 <i>FG</i> -modules	38
5 <i>FG</i> -submodules and reducibility	49
6 Group algebras	53
7 <i>FG</i> -homomorphisms	61
8 Maschke's Theorem	70
9 Schur's Lemma	78
10 Irreducible modules and the group algebra	89
11 More on the group algebra	95
12 Conjugacy classes	104
13 Characters	117
14 Inner products of characters	133
15 The number of irreducible characters	152
16 Character tables and orthogonality relations	159
17 Normal subgroups and lifted characters	168
18 Some elementary character tables	179
19 Tensor products	188
20 Restriction to a subgroup	210
21 Induced modules and characters	224
22 Algebraic integers	244
23 Real representations	263
24 Summary of properties of character tables	283
25 Characters of groups of order pq	288
26 Characters of some p -groups	298
27 Character table of the simple group of order 168	311

28	Character table of $GL(2, q)$	322
29	Permutations and characters	337
30	Applications to group theory	348
31	Burnside's Theorem	361
32	An application of representation theory to molecular vibration	367
	<i>Solutions to exercises</i>	397
	<i>Bibliography</i>	454
	<i>Index</i>	455

1

Groups and homomorphisms

This book is devoted to the study of an aspect of group theory, so we begin with a résumé of facts about groups, most of which you should know already. In addition, we introduce several examples, such as dihedral groups and symmetric groups, which we shall use extensively to illustrate the later theory. An elementary course on abstract algebra would normally cover all the material in the chapter, and any book on basic group theory will supply you with further details. One or two results which we shall use only infrequently are demoted to the exercises at the end of the chapter – you can refer to the solutions if necessary.

Groups

A *group* consists of a set G , together with a rule for combining any two elements g, h of G to form another element of G , written gh ; this rule must satisfy the following axioms:

- (1) for all g, h, k in G ,

$$(gh)k = g(hk);$$

- (2) there exists an element e in G such that for all g in G ,

$$eg = ge = g;$$

- (3) for all g in G , there exists an element g^{-1} in G such that

$$gg^{-1} = g^{-1}g = e.$$

We refer to the rule for combining elements of G as the *product operation* on G .

Axiom (1) states that the product operation is *associative*; the element e in axiom (2) is an *identity* element of G ; and g^{-1} is an *inverse* of g in axiom (3).

It is elementary to see that G has just one identity element, and that every g in G has just one inverse. Usually we write 1, rather than e , for the identity element of G .

The product of an element g with itself, gg , is written g^2 ; similarly $g^3 = g^2g$, $g^{-2} = (g^{-1})^2$, and so on. Also, $g^0 = 1$.

If the number of elements in G is finite, then we call G a *finite group*; the number of elements in G is called the *order* of G , and is written $|G|$.

1.1 Examples


(1) Let n be a positive integer, and denote by \mathbb{C} the set of all complex numbers. The set of n th roots of unity in \mathbb{C} , with the usual multiplication of complex numbers, is a group of order n . It is written as C_n and is called the *cyclic group* of order n . If $a = e^{2\pi i/n}$, then

$$C_n = \{1, a, a^2, \dots, a^{n-1}\},$$

and $a^n = 1$.

(2) The set \mathbb{Z} of all integers, under addition, is a group.

(3) Let n be an integer with $n \geq 3$, and consider the rotation and reflection symmetries of a regular n -sided polygon.



There are n rotation symmetries: these are $\rho_0, \rho_1, \dots, \rho_{n-1}$ where ρ_k is the (clockwise) rotation about the centre O through an angle $2\pi k/n$. There are also n reflection symmetries: these are reflections in the n lines passing through O and a corner or the mid-point of a side of the polygon.

These $2n$ rotations and reflections form a group under the product operation of *composition* (that is, for two symmetries f and g , the product fg means 'first do f , then do g '). This group is called the *dihedral group* of order $2n$, and is written D_{2n} .

Let A be a corner of the polygon. Write b for the reflection in the

line through O and A , and write a for the rotation ρ_1 . Then the n rotations are

$$1, a, a^2, \dots, a^{n-1}$$

(where 1 denotes the identity, which leaves the polygon fixed); and the n reflections are

$$b, ab, a^2b, \dots, a^{n-1}b.$$

Thus all elements of D_{2n} are products of powers of a and b – that is, D_{2n} is *generated* by a and b .

Check that

$$a^n = 1, b^2 = 1 \text{ and } b^{-1}ab = a^{-1}.$$

These relations determine the product of any two elements of the group. For example, we have $ba^j = a^{-j}b$ (using the relation $ba = a^{-1}b$), and hence

$$(a^i b)(a^j b) = a^i b a^j b = a^i a^{-j} b b = a^{i-j}.$$

We summarize all this in the presentation

$$D_{2n} = \langle a, b : a^n = 1, b^2 = 1, b^{-1}ab = a^{-1} \rangle.$$

(4) For n a positive integer, the set of all permutations of $\{1, 2, \dots, n\}$, under the product operation of composition, is a group. It is called the *symmetric group* of degree n , and is written S_n . The order of S_n is $n!$.

(5) Let F be either \mathbb{R} (the set of real numbers) or \mathbb{C} (the set of complex numbers). The set of all invertible $n \times n$ matrices with entries in F , under matrix multiplication, forms a group. This group is called the *general linear group* of degree n over F , and is denoted by $GL(n, F)$. It is an infinite group. The identity of $GL(n, F)$ is of course the identity matrix, which we denote by I_n or just I .

A group G is said to be *abelian* if $gh = hg$ for all g and h in G . While C_n and \mathbb{Z} are abelian, most of the other examples given above are non-abelian groups.

Subgroups

Let G be a group. A subset H of G is said to be a *subgroup* if H is itself a group under the product operation inherited from G . We use the notation $H \leq G$ to indicate that H is a subgroup of G .

It is easy to see that a subset H of a group G is a subgroup if and only if the following two conditions hold:

- (1) $1 \in H$, and
- (2) if $h, k \in H$ then $hk^{-1} \in H$.

1.2 Examples

- (1) For every group G , both $\{1\}$ and G are subgroups of G .
- (2) Let G be a group and $g \in G$. The subset

$$\langle g \rangle = \{g^n : n \in \mathbb{Z}\}$$

is a subgroup of G , called the *cyclic subgroup generated by g* . If $g^n = 1$ for some $n \geq 1$, then $\langle g \rangle$ is finite. In this case, let r be the least positive integer such that $g^r = 1$; then r is equal to the number of elements in $\langle g \rangle$ – indeed,

$$\langle g \rangle = \{1, g, g^2, \dots, g^{r-1}\}.$$

We call r the *order* of the element g .

If $G = \langle g \rangle$ for some $g \in G$ then we call G a *cyclic group*. The groups C_n and \mathbb{Z} in Examples 1.1 are cyclic.

- (3) Let G be a group and let $a, b \in G$. Define H to be the subset of G consisting of all elements which are products of powers of a and b – that is, all elements of the form

$$a^{i_1} b^{j_1} a^{i_2} b^{j_2} \dots a^{i_n} b^{j_n}$$

for some n , where $i_k, j_k \in \mathbb{Z}$ for $1 \leq k \leq n$. Then H is a subgroup of G ; we call H the subgroup *generated by a and b* , and write

$$H = \langle a, b \rangle.$$

Given any finite set S of elements of G , we can similarly define $\langle S \rangle$, the subgroup of G generated by S .

This construction gives a powerful method of finding new groups as subgroups of given groups, such as general linear or symmetric groups. We illustrate the construction in the next example, and again in Example 1.5 below.

- (4) Let $G = \text{GL}(2, \mathbb{C})$, the group of invertible 2×2 matrices with entries in \mathbb{C} , and let

$$A = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Put $H = \langle A, B \rangle$, the subgroup of G generated by A and B . Check that

$$A^4 = I, A^2 = B^2, B^{-1}AB = A^{-1}.$$

Using the third relation, we see that every element of H has the form $A^i B^j$ for some integers i, j ; and using the first two relations, we can take $0 \leq i \leq 3$ and $0 \leq j \leq 1$. Hence H has at most eight elements. Since the matrices

$$A^i B^j \quad (0 \leq i \leq 3, 0 \leq j \leq 1)$$

are all distinct, in fact $|H| = 8$.

The group H is called the *quaternion group* of order 8, and is written Q_8 . The above three relations determine the product of any two elements of Q_8 , so we have the presentation

$$Q_8 = \langle A, B: A^4 = I, A^2 = B^2, B^{-1}AB = A^{-1} \rangle.$$

(5) A *transposition* in the symmetric group S_n is a permutation which interchanges two of the numbers $1, 2, \dots, n$ and fixes the other $n - 2$ numbers. Every permutation g in S_n can be expressed as a product of transpositions. It can be shown that either all such expressions for g have an even number of transpositions, or they all have an odd number of transpositions; we call g an *even* or an *odd* permutation, accordingly. The subset

$$A_n = \{g \in S_n: g \text{ is an even permutation}\}$$

is a subgroup of S_n , called the *alternating group* of degree n .

Direct products

We describe a construction which produces a new group from given ones.

Let G and H be groups, and consider

$$G \times H = \{(g, h): g \in G \text{ and } h \in H\}.$$

Define a product operation on $G \times H$ by

$$(g, h)(g', h') = (gg', hh')$$

for all $g, g' \in G$ and all $h, h' \in H$. With this product operation, $G \times H$ is a group, called the *direct product* of G and H .

More generally, if G_1, \dots, G_r are groups, then the direct product $G_1 \times \dots \times G_r$ is

$$\{(g_1, \dots, g_r): g_i \in G_i \text{ for } 1 \leq i \leq r\},$$

with product operation defined by

$$(g_1, \dots, g_r)(g'_1, \dots, g'_r) = (g_1 g'_1, \dots, g_r g'_r).$$

If all the groups G_i are finite, then $G_1 \times \dots \times G_r$ is also finite, of order $|G_1| \dots |G_r|$.

1.3 Example

The group $C_2 \times \dots \times C_2$ (r factors) has order 2^r and all its non-identity elements have order 2.

Functions

A *function* from one set G to another set H is a rule which assigns a unique element of H to each element of G . In this book, we generally apply functions on the *right* – that is, the image of g under a function ϑ is written as $g\vartheta$, not as ϑg . We often indicate that ϑ is a function from G to H by the notation $\vartheta: G \rightarrow H$. By an expression $\vartheta: g \rightarrow h$, where $g \in G$ and $h \in H$, we mean that $h = g\vartheta$.

A function $\vartheta: G \rightarrow H$ is *invertible* if there is a function $\phi: H \rightarrow G$ such that for all $g \in G$, $h \in H$,

$$(g\vartheta)\phi = g \text{ and } (h\phi)\vartheta = h.$$

Then ϕ is called the *inverse* of ϑ , and is written as ϑ^{-1} . A function ϑ from G to H is invertible if and only if it is both *injective* (that is, $g_1\vartheta = g_2\vartheta$ for $g_1, g_2 \in G$ implies that $g_1 = g_2$) and *surjective* (that is, for every $h \in H$ there exists $g \in G$ such that $g\vartheta = h$). An invertible function is also called a *bijection*.

Homomorphisms

Given groups G and H , those functions from G to H which ‘preserve the group structure’ – the so-called homomorphisms – are of particular importance.

If G and H are groups, then a *homomorphism* from G to H is a function $\vartheta: G \rightarrow H$ which satisfies

$$(g_1 g_2)\vartheta = (g_1\vartheta)(g_2\vartheta) \text{ for all } g_1, g_2 \in G.$$

An invertible homomorphism is called an *isomorphism*. If there is an isomorphism ϑ from G to H , then G and H are said to be *isomorphic*, and we write $G \cong H$; also, ϑ^{-1} is an isomorphism from H to G , so $H \cong G$.

The following example displays a technique which can often be used to prove that certain functions are homomorphisms.

1.4 Example

Let $G = D_{2n} = \langle a, b: a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle$, and write the $2n$ elements of G in the form $a^i b^j$ with $0 \leq i \leq n-1$, $0 \leq j \leq 1$. Let H be any group, and suppose that H contains elements x and y which satisfy

$$x^n = y^2 = 1, y^{-1}xy = x^{-1}.$$

We shall prove that the function $\vartheta: G \rightarrow H$ defined by

$$\vartheta: a^i b^j \rightarrow x^i y^j \quad (0 \leq i \leq n-1, 0 \leq j \leq 1)$$

is a homomorphism.

Suppose that $0 \leq r \leq n-1$, $0 \leq s \leq 1$, $0 \leq t \leq n-1$, $0 \leq u \leq 1$. Then

$$a^r b^s a^t b^u = a^i b^j$$

for some i, j with $0 \leq i \leq n-1$, $0 \leq j \leq 1$. Moreover, i and j are determined by repeatedly using the relations

$$a^n = b^2 = 1, b^{-1}ab = a^{-1}.$$

Since we have $x^n = y^2 = 1$, $y^{-1}xy = x^{-1}$, we can also deduce that

$$x^r y^s x^t y^u = x^i y^j.$$

Therefore,

$$\begin{aligned} (a^r b^s a^t b^u)\vartheta &= (a^i b^j)\vartheta = x^i y^j = x^r y^s x^t y^u \\ &= (a^r b^s)\vartheta \cdot (a^t b^u)\vartheta, \end{aligned}$$

and so ϑ is a homomorphism.

We now demonstrate the technique of Example 1.4 in action.

1.5 Example

Let $G = S_5$ and let x, y be the following permutations in G :

$$x = (1\ 2\ 3\ 4\ 5), y = (2\ 5)(3\ 4).$$

(Here we adopt the usual cycle notation – thus, $(1\ 2\ 3\ 4\ 5)$ denotes the permutation $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 5 \rightarrow 1$, and so on.) Check that

$$x^5 = y^2 = 1, y^{-1}xy = x^{-1}.$$

Let H be the subgroup $\langle x, y \rangle$ of G . Using the above relations, we see that

$$H = \{x^i y^j : 0 \leq i \leq 4, 0 \leq j \leq 1\},$$

a group of order 10.

Now recall that

$$D_{10} = \langle a, b : a^5 = b^2 = 1, b^{-1}ab = a^{-1} \rangle.$$

By Example 1.4, the function $\vartheta: D_{10} \rightarrow H$ defined by

$$\vartheta: a^i b^j \rightarrow x^i y^j \quad (0 \leq i \leq 4, 0 \leq j \leq 1)$$

is a homomorphism. Since ϑ is invertible, it is an isomorphism. Thus, $H = \langle x, y \rangle \cong D_{10}$.

Cosets

Let G be a group and let H be a subgroup of G . For x in G , the subset

$$Hx = \{hx : h \in H\}$$

of G is called a *right coset* of H in G . The distinct right cosets of H in G form a partition of G (that is, every element of G is in precisely one of the cosets).

Suppose now that G is finite, and let Hx_1, \dots, Hx_r be all the distinct right cosets of H in G . For all i , the function

$$h \rightarrow hx_i \quad (h \in H)$$

is a bijection from H to Hx_i , and so $|Hx_i| = |H|$. Since

$$G = Hx_1 \cup \dots \cup Hx_r, \text{ and}$$

$$Hx_i \cap Hx_j \text{ is empty if } i \neq j,$$

we deduce that

$$|G| = r|H|.$$

In particular, we have

1.6 Lagrange's Theorem

If G is a finite group and H is a subgroup of G , then $|H|$ divides $|G|$.

The number r of distinct right cosets of H in G is called the *index* of H in G , and is written as $|G:H|$. Thus

$$|G:H| = |G|/|H|$$

when G is finite.

Normal subgroups

A subgroup N of a group G is said to be a *normal* subgroup of G if $g^{-1}Ng = N$ for all $g \in G$ (where $g^{-1}Ng = \{g^{-1}ng: n \in N\}$); we write $N \triangleleft G$ to indicate that N is a normal subgroup of G .

Suppose that $N \triangleleft G$ and let G/N be the set of right cosets of N in G . The importance of the condition $g^{-1}Ng = N$ (for all $g \in G$) is that it can be used to show that for all $g, h \in G$, we have

$$\{xy: x \in Ng \text{ and } y \in Nh\} = Ngh.$$

Hence we can define a product operation on G/N by

$$(Ng)(Nh) = Ngh \quad \text{for all } g, h \in G.$$

This makes G/N into a group, called the *factor group* of G by N .

1.7 Examples

(1) For every group G , the sub-groups $\{1\}$ and G are normal sub-groups of G .

(2) For $n \geq 1$, we have $A_n \triangleleft S_n$. If $n \geq 2$ then there are just two right cosets of A_n in S_n , namely

$$A_n = \{g \in S_n: g \text{ even}\}, \text{ and}$$

$$A_n(1\ 2) = \{g \in S_n: g \text{ odd}\}.$$

Thus $|S_n:A_n| = 2$, and so $S_n/A_n \cong C_2$.

(3) Let $G = D_8 = \langle a, b: a^4 = b^2 = 1, b^{-1}ab = a^{-1} \rangle$ and let $N = \langle a^2 \rangle = \{1, a^2\}$. Then $N \triangleleft G$ and

$$G/N = \{N, Na, Nb, Nab\}.$$

Since $(Na)^2 = (Nb)^2 = (Nab)^2 = N$, we see that $G/N \cong C_2 \times C_2$.

The subgroup $\langle a \rangle$ is also normal in G , but the subgroup $H = \langle b \rangle$ is not normal in G , since $b \in H$ while $a^{-1}ba = a^2b \notin H$.