

Graduate Texts in Mathematics

Peter Walters

An Introduction to Ergodic Theory

遍历性理论引论

Springer-Verlag

世界图书出版公司

Peter Walters

An Introduction to Ergodic Theory

With 8 Illustrations



Springer

书 名: An Introduction to Ergodic Theory
作 者: P. Walters
中 译 名: 遍历性理论引论
出 版 者: 世界图书出版公司北京公司
印 刷 者: 北京世图印刷厂
发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)
联系电话: 010-64015659, 64038347
电子信箱: kjsk@vip.sina.com
开 本: 24 印 张: 11
出版年代: 2003 年 6 月
书 号: 7-5062-6009-3/O · 398
版权登记: 图字: 01-2003-3600
定 价: 28.00 元

世界图书出版公司北京公司已获得 Springer-Verlag 授权在中国
大陆独家重印发行。

Peter Walters
Mathematics Institute
University of Warwick
Coventry CV4 7AL
England

Editorial Board

S. Axler
Mathematics Department
San Francisco State University
San Francisco, CA 94132
USA

F.W. Gehring
Mathematics Department
East Hall
University of Michigan
Ann Arbor, MI 48109-1109
USA

K.A. Ribet
Mathematics Department
University of California at Berkeley
Berkeley, CA 94720-3840
USA

AMS Subject Classification: 28-01, 28DXX, 47A35, 54H20

Library of Congress Cataloging-in-Publication Data
Walters, Peter, 1943–

An introduction to ergodic theory.

(Graduate texts in mathematics; 79)

Previously published as: Ergodic theory. 1975.

Bibliography: p.

Includes index.

I. Ergodic theory. I. Title. II. Series.

QA313.W34 1981 515.4'2 81-9319

ISBN 0-387-95152-0

AACR2

Printed on acid-free paper.

First softcover printing, 2000.

© 1982 Springer-Verlag New York, Inc.

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer-Verlag New York, Inc., 175 Fifth Avenue, New York, NY 10010, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the People's Republic of China only and not for export therefrom.

Reprinted in China by Beijing World Publishing Corporation, 2003

9 8 7 6 5 4 3 2 1

ISBN 0-387-95152-0

SPIN 10783040

Springer-Verlag New York Berlin Heidelberg
A member of BertelsmannSpringer Science+Business Media GmbH

Preface

In 1970 I gave a graduate course in ergodic theory at the University of Maryland in College Park, and these lectures were the basis of the Springer Lecture Notes in Mathematics Volume 458 called "Ergodic Theory—Introductory Lectures" which was published in 1975. This volume is now out of print, so I decided to revise and add to the contents of these notes. I have updated the earlier chapters and have added some new chapters on the ergodic theory of continuous transformations of compact metric spaces. In particular, I have included some material on topological pressure and equilibrium states. In recent years there have been some fascinating interactions of ergodic theory with differentiable dynamics, differential geometry, number theory, von Neumann algebras, probability theory, statistical mechanics, and other topics. In Chapter 10 I have briefly described some of these and given references to some of the others. I hope that this book will give the reader enough foundation to tackle the research papers on ergodic theory and its applications.

I would like to dedicate this volume to the memory of Rufus Bowen who died on July 30, 1978 at the age of 31. He made outstanding contributions to ergodic theory and his friendship enhanced the lives of all who knew him.

April, 1981

PETER WALTERS

Contents

Chapter 0	
Preliminaries	1
§0.1 Introduction	1
§0.2 Measure Spaces	3
§0.3 Integration	6
§0.4 Absolutely Continuous Measures and Conditional Expectations	8
§0.5 Function Spaces	9
§0.6 Haar Measure	11
§0.7 Character Theory	12
§0.8 Endomorphisms of Tori	14
§0.9 Perron–Frobenius Theory	16
§0.10 Topology	17
Chapter 1	
Measure-Preserving Transformations	19
§1.1 Definition and Examples	19
§1.2 Problems in Ergodic Theory	23
§1.3 Associated Isometries	24
§1.4 Recurrence	26
§1.5 Ergodicity	26
§1.6 The Ergodic Theorem	34
§1.7 Mixing	39
Chapter 2	
Isomorphism, Conjugacy, and Spectral Isomorphism	53
§2.1 Point Maps and Set Maps	53
§2.2 Isomorphism of Measure-Preserving Transformations	57
§2.3 Conjugacy of Measure-Preserving Transformations	59
§2.4 The Isomorphism Problem	62

§2.5 Spectral Isomorphism	63
§2.6 Spectral Invariants	66
Chapter 3	
Measure-Preserving Transformations with Discrete Spectrum	68
§3.1 Eigenvalues and Eigenfunctions	68
§3.2 Discrete Spectrum	69
§3.3 Group Rotations	72
Chapter 4	
Entropy	75
§4.1 Partitions and Subalgebras	75
§4.2 Entropy of a Partition	77
§4.3 Conditional Entropy	80
§4.4 Entropy of a Measure-Preserving Transformation	86
§4.5 Properties of $h(T, \mathcal{A})$ and $h(T)$	89
§4.6 Some Methods for Calculating $h(T)$	94
§4.7 Examples	100
§4.8 How Good an Invariant is Entropy?	103
§4.9 Bernoulli Automorphisms and Kolmogorov Automorphisms	105
§4.10 The Pinsker σ -Algebra of a Measure-Preserving Transformation	113
§4.11 Sequence Entropy	114
§4.12 Non-invertible Transformations	115
§4.13 Comments	116
Chapter 5	
Topological Dynamics	118
§5.1 Examples	118
§5.2 Minimality	120
§5.3 The Non-wandering Set	123
§5.4 Topological Transitivity	127
§5.5 Topological Conjugacy and Discrete Spectrum	133
§5.6 Expansive Homeomorphisms	137
Chapter 6	
Invariant Measures for Continuous Transformations	146
§6.1 Measures on Metric Spaces	146
§6.2 Invariant Measures for Continuous Transformations	150
§6.3 Interpretation of Ergodicity and Mixing	154
§6.4 Relation of Invariant Measures to Non-wandering Sets, Periodic Points and Topological Transitivity	156
§6.5 Unique Ergodicity	158
§6.6 Examples	162
Chapter 7	
Topological Entropy	164
§7.1 Definition Using Open Covers	164
§7.2 Bowen's Definition	168
§7.3 Calculation of Topological Entropy	176

Chapter 8	
Relationship Between Topological Entropy and Measure-Theoretic Entropy	182
§8.1 The Entropy Map	182
§8.2 The Variational Principle	187
§8.3 Measures with Maximal Entropy	191
§8.4 Entropy of Affine Transformations	196
§8.5 The Distribution of Periodic Points	203
§8.6 Definition of Measure-Theoretic Entropy Using the Metrics d_n	205
Chapter 9	
Topological Pressure and Its Relationship with Invariant Measures	207
§9.1 Topological Pressure	207
§9.2 Properties of Pressure	214
§9.3 The Variational Principle	217
§9.4 Pressure Determines $M(X, T)$	221
§9.5 Equilibrium States	223
Chapter 10	
Applications and Other Topics	229
§10.1 The Qualitative Behaviour of Diffeomorphisms	229
§10.2 The Subadditive Ergodic Theorem and the Multiplicative Ergodic Theorem	230
§10.3 Quasi-invariant Measures	236
§10.4 Other Types of Isomorphism	238
§10.5 Transformations of Intervals	238
§10.6 Further Reading	239
References	240
Index	247

MAG52/04

CHAPTER 0

Preliminaries

§0.1 Introduction

In its broadest interpretation ergodic theory is the study of the qualitative properties of actions of groups on spaces. The space has some structure (e.g. the space is a measure space, or a topological space, or a smooth manifold) and each element of the group acts as a transformation on the space and preserves the given structure (e.g. each element acts as a measure-preserving transformation, or a continuous transformation, or a smooth transformation).

To see how this type of study arises consider a system of k particles moving in R^3 under known forces. Suppose that the state of the system at a given time is determined by knowing the positions and the momenta of each of the k particles. Thus at a given time the system is determined by a point in R^{6k} . As time continues the system alters according to the differential equations governing the motion, e.g., Hamilton's equations

$$\frac{dq_i}{dt} = \frac{\partial H}{\partial p_i}, \quad \frac{dp_i}{dt} = -\frac{\partial H}{\partial q_i}.$$

If we are given an initial condition and if the equations can be uniquely solved then the corresponding solution gives us the entire history of the system, which is determined by a curve in R^{6k} .

If x is a point in the state space representing the system at a time t_0 , let $T_t(x)$ denote the point of the state space representing the system at time $t + t_0$. From this we see that T_t is a transformation of the state space and, moreover, $T_0 = id$ and $T_{t+s} = T_t \circ T_s$. Thus $t \rightarrow T_t$ is an action of the group R on the state space. Because the Hamiltonian H is constant along solution curves, each energy surface $H^{-1}(e)$ is invariant for the transformation T_t .

so that we get an action of R on each energy surface. One is interested in the asymptotic properties of the action i.e. in T_t for large t . The transformations $T_t|H^{-1}(e)$ are continuous and are smooth if $H^{-1}(e)$ is smooth. Measure theory enters this picture via a theorem of Liouville which tells us that if the forces are of a certain type one can choose coordinates in the state space so that the usual $6k$ -dimensional measure in these coordinates is preserved by each transformation T_t .

The word "ergodic" was introduced by Boltzmann to describe a hypothesis about the action of $\{T_t|t \in R\}$ on an energy surface $H^{-1}(e)$ when the Hamiltonian H is of the type that arises in statistical mechanics. Boltzmann had hoped that each orbit $\{T_t(x)|t \in R\}$ would equal the whole energy surface $H^{-1}(e)$ and he called this statement the ergodic hypothesis. The word "ergodic" is an amalgamation of the Greek words *ergon* (work) and *odos* (path). Boltzmann made the hypothesis in order to deduce the equality of time means and phase means which is a fundamental algorithm in statistical mechanics. The ergodic hypothesis, as stated above, is false. The property the flow needs to satisfy in order to equate time means and phase means of real-valued functions is what is now called ergodicity.

It is common to use the name ergodic theory to describe only the qualitative study of actions of groups on measure spaces. The actions on topological spaces and smooth manifolds are often called topological dynamics and differentiable dynamics. This measure theoretic study began in the early 1930's and the ergodic theorems of Birkhoff and von Neumann were proved then. The next major advance was the introduction of entropy by Kolmogorov in 1958. The proof, by Ornstein in 1969, that entropy was complete for Bernoulli shifts revitalised the work on the isomorphism problem. During recent years ergodic theory had been used to give important results in other branches of mathematics.

We shall study actions of the group Z of integers on a space X i.e. we study a transformation $T: X \rightarrow X$ and its iterates T^n , $n \in Z$. This is simpler than studying the actions of R . Of course, if $\{T_t|t \in R\}$ is an action of R on X , then by choosing $t_0 \neq 0$ and observing the system at the times $\dots, -t_0, 0, t_0, 2t_0, 3t_0, \dots$, we are considering $(T_{t_0})^n$, $n \in Z$.

In the following sections we summarise some of the background ideas and notation we shall be using.

We shall use Z to denote the set of integers, Z^+ to denote the non-negative integers, R to denote the real numbers, R^+ to denote the non-negative reals, and C to denote the complex numbers. The empty set will be denoted by \emptyset .

If A, B are subsets of a set X , then $B \setminus A$ denotes the difference set $\{x \in X | x \in B, x \notin A\}$, and $A \triangle B$ denotes the symmetric difference $(A \setminus B) \cup (B \setminus A)$. We use 2^X to denote the collection of all subsets of X .

We use "iff" to denote "if and only if." We number lemmas and theorems in a single sequence (Theorem 5.6 is the sixth theorem in Chapter 5) but give a corollary the same number as the corresponding theorem (Corollary 5.6.2 is the second corollary of Theorem 5.6).

§0.2 Measure Spaces

We shall generally refer to Kingman and Taylor [1] and Parthasarathy [2]. Let X be a set. A σ -algebra of subsets of X is a collection \mathcal{B} of subsets of X satisfying the following three conditions: (i) $X \in \mathcal{B}$; (ii) if $B \in \mathcal{B}$ then $X \setminus B \in \mathcal{B}$; (iii) if $B_n \in \mathcal{B}$ for $n \geq 1$ then $\bigcup_{n=1}^{\infty} B_n \in \mathcal{B}$.

We then call the pair (X, \mathcal{B}) a *measurable space*. A *finite measure* on (X, \mathcal{B}) is a function $m: \mathcal{B} \rightarrow \mathbb{R}^+$ satisfying $m(\emptyset) = 0$ and $m(\bigcup_{n=1}^{\infty} B_n) = \sum_{n=1}^{\infty} m(B_n)$ whenever $\{B_n\}_1^{\infty}$ is a sequence of members of \mathcal{B} which are pairwise disjoint subsets of X . (Actually the latter condition implies $m(\emptyset) = 0$ since m is finite-valued.) A finite measure space is a triple (X, \mathcal{B}, m) where (X, \mathcal{B}) is a measurable space and m is a finite measure on (X, \mathcal{B}) . We say (X, \mathcal{B}, m) is a *probability space*, or a *normalised measure space*, if $m(X) = 1$. We then say m is a *probability measure* on (X, \mathcal{B}) . We shall usually consider only probability spaces.

A *finite signed measure* on a measurable space (X, \mathcal{B}) is a function $m: \mathcal{B} \rightarrow \mathbb{R}$ satisfying $m(\emptyset) = 0$ and $m(\bigcup_{n=1}^{\infty} B_n) = \sum_{n=1}^{\infty} m(B_n)$ whenever $\{B_n\}_1^{\infty}$ is a sequence of members of \mathcal{B} which are pairwise disjoint subsets of X . The *Jordan decomposition* says that a finite signed measure m on (X, \mathcal{B}) can be written as the difference $m = m_1 - m_2$ of two finite measures on (X, \mathcal{B}) which are uniquely determined by m (see Kingman and Taylor [1], pp. 62 and 64).

Measurable spaces are usually constructed by having a collection \mathcal{S} of interesting subsets of a set X (such as the collection of all subintervals of $[0, 1]$) and then considering the smallest σ -algebra \mathcal{B} containing all these subsets. This makes sense because 2^X is a σ -algebra and any intersection of σ -algebras of subsets of X is also a σ -algebra of subsets of X . It is then usually difficult to decide which subsets of X are in \mathcal{B} . When constructing a measure on a measurable space (X, \mathcal{B}) obtained in this way, one usually knows what values the measure should take on members of \mathcal{S} and then one needs to extend it to be defined on \mathcal{B} . We now describe the basic extension theorem of this type. This involves discussion of the properties the collection \mathcal{S} should have.

A collection \mathcal{S} of subsets of X is called a *semi-algebra* if the following three conditions hold: (i) $\emptyset \in \mathcal{S}$; (ii) if $A, B \in \mathcal{S}$, then $A \cap B \in \mathcal{S}$; (iii) if $A \in \mathcal{S}$, then $X \setminus A = \bigcup_{i=1}^n E_i$ where each $E_i \in \mathcal{S}$ and E_1, \dots, E_n are pairwise disjoint subsets of X . For example, the collection of all subintervals of $[0, 1]$ is a semi-algebra. Also, the collection of all subintervals of $[0, 1]$ of the forms $[0, b]$ and $(a, b]$, with $0 \leq a < b \leq 1$, forms a semi-algebra.

A collection \mathcal{A} of subsets of X is called an *algebra* if the following three conditions hold: (i) $\emptyset \in \mathcal{A}$; (ii) if $A, B \in \mathcal{A}$, then $A \cap B \in \mathcal{A}$; (iii) if $A \in \mathcal{A}$, then $X \setminus A \in \mathcal{A}$. Clearly every algebra is a semi-algebra and every σ -algebra is an algebra. In the definition of an algebra we can replace (ii) by the condition that whenever $A_1, \dots, A_n \in \mathcal{A}$ then $\bigcup_{i=1}^n A_i \in \mathcal{A}$.

Since the intersection of any family of algebras of a set X is again an algebra of subsets of X it makes sense to speak of the algebra generated by

any given collection of subsets of X . There is the following simple theorem (Parthasarathy [2], p. 19)

Theorem 0.1. *Let \mathcal{S} be a semi-algebra of subsets of X . The algebra, $\mathcal{A}(\mathcal{S})$, generated by \mathcal{S} consists precisely of those subsets of X that can be written in the form $E = \bigcup_{i=1}^n A_i$ where each $A_i \in \mathcal{S}$ and A_1, \dots, A_n are disjoint subsets of X .*

Suppose \mathcal{S} is a semi-algebra of subsets of X . A function $\tau: \mathcal{S} \rightarrow R^+$ is called *finitely additive* if $\tau(\emptyset) = 0$ and $\tau(\bigcup_{i=1}^n E_i) = \sum_{i=1}^n \tau(E_i)$ whenever E_1, \dots, E_n are members of \mathcal{S} which are pairwise disjoint subsets of X and $\bigcup_{i=1}^n E_i \in \mathcal{S}$. Such a map τ is called *countably additive* if the second condition is replaced by the requirement that $\tau(\bigcup_{i=1}^\infty E_i) = \sum_{i=1}^\infty \tau(E_i)$ whenever $\{E_i\}_{i=1}^\infty$ are members of \mathcal{S} which are pairwise disjoint subsets of X and $\bigcup_{i=1}^\infty E_i \in \mathcal{S}$. If \mathcal{S} is the semi-algebra of all subintervals of $[0, 1]$ of the form $[0, b]$ and $(a, b]$ then the length function is countably additive. A simple application of Theorem 0.1 gives the following (Parthasarathy [2], pp. 20 and 59).

Theorem 0.2. *If \mathcal{S} is a semi-algebra of subsets of X and $\tau: \mathcal{S} \rightarrow R^+$ is finitely additive then there is a unique finitely additive function $\tau_1: \mathcal{A}(\mathcal{S}) \rightarrow R^+$ which is an extension of τ (i.e. $\tau_1 = \tau$ on \mathcal{S}). If τ is countably additive then so is τ_1 .*

It could be that $X \notin \mathcal{S}$ but we always have that X is a disjoint union of a finite number of members E_1, \dots, E_n of \mathcal{S} so $\tau_1(X) = 1$ if $\sum_{i=1}^n \tau(E_i) = 1$.

There is the following theorem on extension from an algebra \mathcal{A} to the σ -algebra $\mathcal{B}(\mathcal{A})$ generated by \mathcal{A} . ($\mathcal{B}(\mathcal{A})$ is the intersection of all σ -algebras that contain \mathcal{A} .) (See Parthasarathy [2], pp. 70 and 71).

Theorem 0.3. *Let \mathcal{A} be an algebra of subsets of X and let $\tau_1: \mathcal{A} \rightarrow R^+$ be countably additive and $\tau_1(X) = 1$. Then there is a unique probability measure τ_2 on $(X, \mathcal{B}(\mathcal{A}))$ which extends τ_1 .*

By combining Theorems 0.2 and 0.3 we see that a countably additive function τ on a semi-algebra \mathcal{S} can be uniquely extended to a probability measure on $(X, \mathcal{B}(\mathcal{S}))$ if $\sum_{i=1}^n \tau(E_i) = 1$ when $X = \bigcup_{i=1}^n E_i$ is a disjoint union of members of \mathcal{S} . As an example the length function defined on the semi-algebra of all subintervals of $[0, 1]$ of the form $[0, b]$ and $(a, b]$ can be uniquely extended to a probability measure, called the Lebesgue measure, defined on the Borel subsets of $[0, 1]$.

In checking that the extension works for particular examples the most difficult part is usually showing countable additivity. This can sometimes be done for τ on the semi-algebra \mathcal{S} (as for Lebesgue measure) but it is sometimes more convenient to prove that τ is finitely additive and that the finitely additive extension $\tau_1: \mathcal{A}(\mathcal{S}) \rightarrow R^+$ is countably additive. The main tool for this is the following theorem (Kingman and Taylor [1], p. 56).

Theorem 0.4. Let \mathcal{A} be an algebra of subsets of X and let $\tau_1: \mathcal{A} \rightarrow R^+$ be finitely additive and let $\tau_1(X) = 1$. Then τ_1 will be countably additive if for every decreasing sequence $E_1 \supset E_2 \supset E_3 \supset \cdots$ of members of \mathcal{A} with $\bigcap_{n=1}^{\infty} E_n = \emptyset$ we have $\tau_1(E_n) \rightarrow 0$.

One situation where this theorem is used is in defining product measures on a countable product of probability spaces. For $i \in Z$ let $(X_i, \mathcal{B}_i, m_i)$ be a probability space. Let $X = \prod_{i=-\infty}^{\infty} X_i$. So a point of X is a bisequence $\{x_i\}_{i=-\infty}^{\infty}$ with $x_i \in X_i$ for each i . We now define a σ -algebra \mathcal{B} of subsets of X called the product of the σ -algebras \mathcal{B}_i . Let $n \geq 0$, let $A_j \in \mathcal{B}_j$ for $|j| \leq n$, and consider the set

$$\prod_{i=-\infty}^{-(n+1)} X_i \times \prod_{j=-n}^n A_j \times \prod_{i=n+1}^{\infty} X_i = \{(x_i)_{i=-\infty}^{\infty} \in X \mid x_j \in A_j \text{ for } |j| \leq n\}.$$

Such a set is called a *measurable rectangle* and the collection of all such subsets of X forms a semi-algebra \mathcal{S} . The σ -algebra \mathcal{B} is the σ -algebra generated by \mathcal{S} . We write $(X, \mathcal{B}) = \prod_{i=-\infty}^{\infty} (X_i, \mathcal{B}_i)$. If we define $\tau: \mathcal{S} \rightarrow R^+$ by giving the above rectangle the value $\prod_{j=-n}^n m_j(A_j)$, then one can use Theorems 0.2 and 0.4 (see Kingman and Taylor [1], p. 140) to extend τ to a probability measure m on (X, \mathcal{B}) . The probability space (X, \mathcal{B}, m) is called the *direct product* of the spaces $(X_i, \mathcal{B}_i, m_i)$ and is sometimes denoted $\prod_{i=-\infty}^{\infty} (X_i, \mathcal{B}_i, m_i)$. The corresponding construction holds for a product $\prod_{i=0}^{\infty} (X_i, \mathcal{B}_i, m_i)$.

A special type of product space will be important for us. Here each space $(X_i, \mathcal{B}_i, m_i)$ is the same space (Y, \mathcal{C}, μ) and Y is the finite set $\{0, 1, \dots, k-1\}$, $\mathcal{C} = 2^Y$, and μ is given by a probability vector $(p_0, p_1, \dots, p_{k-1})$ where $p_i = \mu(\{i\})$. We can take elementary rectangles where each A_j (in the description above) is taken to be one point of Y . So if $n \geq 0$ and $a_j \in Y$, $|j| \leq n$, such an elementary rectangle has the form $\{(x_i)_{i=-\infty}^{\infty} \mid x_j = a_j \text{ for } |j| \leq n\}$. We shall denote this set by ${}_n[a_{-n}, a_{-(n+1)}, \dots, a_{n-1}, a_n]_n$ and call it a *block* with end points $-n$ and n . The collection of all these sets form a semi-algebra which generates the product σ -algebra \mathcal{B} . We have $m({}_n[a_{-n}, \dots, a_n]_n) = \prod_{i=-n}^n p_{a_i}$. The measure m is called the (p_0, \dots, p_{k-1}) -product measure. Sometimes we consider *blocks* with end points h and l where $h \leq l$. Such a set is one of the form ${}_h[a_h, \dots, a_l]_l = \{(x_i)_{i=-\infty}^{\infty} \mid x_i = a_i \text{ for } h \leq i \leq l\}$. It has measure $\prod_{i=h}^l p_{a_i}$.

Theorem 0.4 can also be used to obtain further measures on the space (X, \mathcal{B}) where $X = \prod_{i=-\infty}^{\infty} Y$, $Y = \{0, 1, \dots, k-1\}$, and \mathcal{B} is the product σ -algebra described above. The following is a special case of the Daniell-Kolmogorov consistency theorem (Parthasarathy [2], p. 119).

Theorem 0.5. Fix $k \geq 1$ and let $Y = \{0, 1, \dots, k-1\}$ and $(X, \mathcal{B}) = \prod_{i=-\infty}^{\infty} (Y, 2^Y)$. For each natural number n and $a_0, \dots, a_n \in Y$ suppose a non-negative real number $p_n(a_0, \dots, a_n)$ is given so that

$$(a) \quad \sum_{a_0 \in Y} p_0(a_0) = 1$$

and

$$(b) \quad p_n(a_0, \dots, a_n) = \sum_{a_{n+1} \in Y} p_{n+1}(a_0, \dots, a_n, a_{n+1}).$$

Then there is a unique probability measure m on (X, \mathcal{B}) with $m([a_h, \dots, a_l]_i) = p_{l-h}(a_h, \dots, a_l)$ for all $h \leq l$ and all $a_i \in Y$, $h \leq i \leq l$.

The proof boils down to showing that the function naturally defined on the algebra of all finite unions of elementary rectangles is countable additive by using Theorem 0.4.

There is another way, which is useful in some proofs, of describing the σ -algebra $\mathcal{B}(\mathcal{A})$ generated by an algebra \mathcal{A} . A collection M of subsets of X is called a *monotone class* if whenever $E_1 \subset E_2 \subset E_3 \subset \dots$ all belong to M then so does $\bigcup_{n=1}^{\infty} E_n$ and whenever $F_1 \supset F_2 \supset F_3 \supset \dots$ all belong to M then so does $\bigcap_{n=1}^{\infty} F_n$. Since the intersection of any family of monotone classes is a monotone class, we can speak of the monotone class generated by any given collection of subsets of X .

Theorem 0.6. *Let \mathcal{A} be an algebra of subsets of X . Then $\mathcal{B}(\mathcal{A})$ equals the monotone class generated by \mathcal{A} .*

As we have seen we usually know the elements of an algebra \mathcal{A} but we do not know which subsets of X belong to $\mathcal{B}(\mathcal{A})$. This problem can sometimes be overcome by using the following approximation theorem (Kingman and Taylor [1], p. 84).

Theorem 0.7. *Let (X, \mathcal{B}, m) be a probability space and let \mathcal{A} be an algebra of subsets of X with $\mathcal{B}(\mathcal{A}) = \mathcal{B}$. Then for each $\varepsilon > 0$ and each $B \in \mathcal{B}$ there is some $A \in \mathcal{A}$ with $m(A \triangle B) < \varepsilon$.*

Note that when $m(A \triangle B) < \varepsilon$ then $|m(A) - m(B)| < \varepsilon$ because $m(A) = m(A \setminus B) + m(A \cap B)$ and $m(B) = m(B \setminus A) + m(A \cap B)$, so that $|m(A) - m(B)| \leq m(A \triangle B)$.

§0.3 Integration

Let $\mathcal{B}(R)$ denote the σ -algebra of Borel subsets of R . This is the σ -algebra generated by all open subsets of R and is also generated by the collection of all intervals, or by the collection of all intervals of the form (c, ∞) .

Let (X, \mathcal{B}, m) be a measure space. A function $f: X \rightarrow R$ is *measurable* if $f^{-1}(D) \in \mathcal{B}$ whenever $D \in \mathcal{B}(R)$ or equivalently if $f^{-1}(c, \infty) \in \mathcal{B}$ for all $c \in R$. A function $f: X \rightarrow \mathbb{C}$ is measurable if both its real and imaginary parts are

measurable. If X is a topological space and \mathcal{B} the σ -algebra generated by the open subsets of X , then any continuous function $f: X \rightarrow \mathbb{C}$ is measurable. We say $f = g$ a.e. if $m(\{x: f(x) \neq g(x)\}) = 0$. Suppose X is a topological space, $\mathcal{B}(X)$ its σ -algebra of Borel sets and m a measure on $(X, \mathcal{B}(X))$ with the property that each non-empty open set has non-zero measure. Then for two continuous functions $f, g: X \rightarrow \mathbb{R}$, $f = g$ a.e. implies $f = g$ because $\{x: f(x) - g(x) \neq 0\}$ is an open set of zero measure.

Let (X, \mathcal{B}, m) be a probability space. A function $f: X \rightarrow \mathbb{R}$ is a *simple function* if it can be written in the form $\sum_{i=1}^n a_i \chi_{A_i}$, where $a_i \in \mathbb{R}$, $A_i \in \mathcal{B}$, the sets A_i are disjoint subsets of X , and χ_{A_i} denotes the characteristic function of A_i . Simple functions are measurable. We define the integral for simple functions by:

$$\int f \, dm = \sum_{i=1}^n a_i m(A_i).$$

This value is independent of the representation $\sum_i a_i \chi_{A_i}$.

Suppose $f: X \rightarrow \mathbb{R}$ is measurable and $f \geq 0$. Then there exists an increasing sequence of simple functions $f_n \nearrow f$. For example, we could take

$$f_n(x) = \begin{cases} \frac{i-1}{2^n}, & \text{if } \frac{i-1}{2^n} \leq f(x) < \frac{i}{2^n} \quad i = 1, \dots, n2^n \\ n, & \text{if } f(x) \geq n. \end{cases}$$

We define $\int f \, dm = \lim_{n \rightarrow \infty} \int f_n \, dm$ and note that this definition is independent of the chosen sequence $\{f_n\}$. We say f is *integrable* if $\int f \, dm < \infty$.

Suppose $f: X \rightarrow \mathbb{R}$ is measurable. Then $f = f^+ - f^-$ where $f^+(x) = \max\{f(x), 0\} \geq 0$ and $f^-(x) = \max\{-f(x), 0\} \geq 0$. We say that f is *integrable* if $\int f^+ \, dm, \int f^- \, dm < \infty$ and we then define

$$\int f \, dm = \int f^+ \, dm - \int f^- \, dm.$$

We say $f: X \rightarrow \mathbb{C}$ is *integrable* ($f = f_1 + i f_2$) if f_1 and f_2 are integrable and we define

$$\int f \, dm = \int f_1 \, dm + i \int f_2 \, dm.$$

Observe that f is integrable if and only if $|f|$ is integrable. If $f = g$ a.e. then one is integrable if the other is and $\int f \, dm = \int g \, dm$.

The two basic theorems on integrating sequences as functions are the following.

Theorem 0.8 (Monotone Convergence Theorem). Suppose $f_1 \leq f_2 \leq f_3 \leq \dots$ is an increasing sequence of integrable real-valued functions on (X, \mathcal{B}, m) . If $\{\int f_n \, dm\}$ is a bounded sequence of real numbers then $\lim_{n \rightarrow \infty} f_n$ exists a.e. and is integrable and $\int(\lim f_n) \, dm = \lim \int f_n \, dm$. If $\{\int f_n \, dm\}$ is an unbounded sequence then either $\lim_{n \rightarrow \infty} f_n$ is infinite on a set of positive measure or $\lim_{n \rightarrow \infty} f_n$ is not integrable.

Theorem 0.9 (Fatou's Lemma). *Let $\{f_n\}$ be a sequence of measurable real-valued functions on (X, \mathcal{B}, m) which is bounded below by an integrable function. If $\liminf_{n \rightarrow \infty} \int f_n dm < \infty$ then $\liminf_{n \rightarrow \infty} f_n$ is integrable and $\int \liminf_{n \rightarrow \infty} f_n dm \leq \liminf_{n \rightarrow \infty} \int f_n dm$.*

Corollary 0.9.1 (Dominated Convergence Theorem). *If $g: X \rightarrow \mathbb{R}$ is integrable and $\{f_n\}$ is a sequence of measurable real-valued functions with $|f_n| \leq g$ a.e. ($n \geq 1$) and $\lim_{n \rightarrow \infty} f_n = f$ a.e. then f is integrable and $\lim \int f_n dm = \int f dm$.*

We denote by $L^1(X, \mathcal{B}, m)$ (or $L^1(m)$) the space of all integrable functions $f: X \rightarrow \mathbb{C}$ where two such functions are identified if they are equal a.e. However we write $f \in L^1(X, \mathcal{B}, m)$ to denote that $f: X \rightarrow \mathbb{C}$ is integrable. The space $L^1(X, \mathcal{B}, m)$ is a Banach space with norm $\|f\|_1 = \int |f| dm$.

If $f \in L^1(X, \mathcal{B}, m)$, then $\int_A f dm$ denotes $\int f \cdot \chi_A dm$.

If m is a finite signed measure on (X, \mathcal{B}) and $m = m_1 - m_2$ is its unique Jordan decomposition into the difference of two finite measures, then we can define $\int f dm = \int f dm_1 - \int f dm_2$ for $f \in L^1(m_1) \cap L^1(m_2)$.

§0.4 Absolutely Continuous Measures and Conditional Expectations

Let (X, \mathcal{B}) be a measurable space and suppose μ, m are two probability measures on (X, \mathcal{B}) . We say μ is *absolutely continuous* with respect to m ($\mu \ll m$) if $\mu(B) = 0$ whenever $m(B) = 0$. The measures are *equivalent* if $\mu \ll m$ and $m \ll \mu$. The following theorem characterises absolute continuity.

Theorem 0.10 (Radon–Nikodym Theorem). *Let μ, m be two probability measures on the measurable space (X, \mathcal{B}) . Then $\mu \ll m$ iff there exists $f \in L^1(m)$, with $f \geq 0$ and $\int f dm = 1$, such that $\mu(B) = \int_B f dm \forall B \in \mathcal{B}$. The function f is unique a.e. (in the sense that any other function with these properties is equal to f a.e.).*

The function f is called the Radon–Nikodym derivative of μ with respect to m and denoted by $d\mu/dm$.

The “opposite” notion to absolute continuity is as follows. Two probability measures μ, m on (X, \mathcal{B}) are said to be *mutually singular* ($\mu \perp m$) if there is some $B \in \mathcal{B}$ with $\mu(B) = 0$ and $m(X \setminus B) = 0$. There is the following decomposition theorem.

Theorem 0.11 (Lebesgue Decomposition Theorem). *Let μ, m be two probability measures on (X, \mathcal{B}) . There exists $p \in [0, 1]$ and probability measures μ_1, μ_2 on (X, \mathcal{B}) such that $\mu = p\mu_1 + (1 - p)\mu_2$ and $\mu_1 \ll m, \mu_2 \perp m$. ($\mu = p\mu_1 + (1 - p)\mu_2$ means $\mu(B) = p\mu_1(B) + (1 - p)\mu_2(B) \forall B \in \mathcal{B}$). The number p and probabilities μ_1, μ_2 are uniquely determined.*

The Radon-Nikodym theorem allows us to define conditional expectations. Let (X, \mathcal{B}, m) be a measure space and let \mathcal{C} be a sub- σ -algebra of \mathcal{B} . We now define the conditional expectation operator $E(\cdot/\mathcal{C}): L^1(X, \mathcal{B}, m) \rightarrow L^1(X, \mathcal{C}, m)$. If $f \in L^1(X, \mathcal{B}, m)$ takes non-negative real values then $\mu_f(C) = a^{-1} \int_C f dm$ (where $a = \int_X f dm$) defines a probability measure, μ_f , on (X, \mathcal{C}, m) and $\mu_f \ll m$. By Theorem 0.10 there is a function $E(f/\mathcal{C}) \in L^1(X, \mathcal{C}, m)$ such that $E(f/\mathcal{C}) \geq 0$ and $\int_C E(f/\mathcal{C}) dm = \int_C f dm \forall C \in \mathcal{C}$. Moreover $E(f/\mathcal{C})$ is unique a.e. If f is real-valued we can consider the positive and negative parts of f and define $E(f/\mathcal{C})$ linearly. Similarly when f is complex-valued we can use the real and imaginary parts to define $E(f/\mathcal{C})$ linearly. Therefore if $f \in L^1(X, \mathcal{B}, m)$ then $E(f/\mathcal{C})$ is the only \mathcal{C} -measurable function with $\int_C E(f/\mathcal{C}) dm = \int_C f dm \forall C \in \mathcal{C}$. The following properties of the map $E(\cdot/\mathcal{C}): L^1(X, \mathcal{B}, m) \rightarrow L^1(X, \mathcal{C}, m)$ hold (Parthasarathy [2], p. 225):

- (i) $E(\cdot/\mathcal{C})$ is linear.
- (ii) If $f \geq 0$, then $E(f/\mathcal{C}) \geq 0$.
- (iii) If $f \in L^1(X, \mathcal{B}, m)$ and g is \mathcal{C} -measurable and bounded,

$$E(fg/\mathcal{C}) = gE(f/\mathcal{C}).$$

- (iv) $|E(f/\mathcal{C})| \leq E(|f|/\mathcal{C}), \quad f \in L^1(X, \mathcal{B}, m)$
- (v) If $\mathcal{C}_2 \subset \mathcal{C}_1$, then $E(E(f/\mathcal{C}_1)/\mathcal{C}_2) = E(f/\mathcal{C}_2), f \in L^1(X, \mathcal{B}, m)$.

§0.5 Function Spaces

One way to deal with some problems on a measure space is to use certain natural Banach spaces of functions associated with the measure space.

Let (X, \mathcal{B}, m) be a measure space and let $p \in \mathbb{R}$ with $p \geq 1$. Consider the set of all measurable functions $f: X \rightarrow \mathbb{C}$ with $|f|^p$ integrable. This space is a vector space under the usual addition and scalar multiplication of functions. If we define an equivalence relation on this set by $f \sim g$ iff $f = g$ a.e. then the space of equivalence classes is also a vector space. Let $L^p(X, \mathcal{B}, m)$ denote the space of equivalence classes, although we write $f \in L^p(X, \mathcal{B}, m)$ to denote that the function $f: X \rightarrow \mathbb{C}$ has $|f|^p$ integrable. The formula $\|f\|_p = [\int |f|^p dm]^{1/p}$ defines a norm on $L^p(X, \mathcal{B}, m)$ and this norm is complete. Therefore $L^p(X, \mathcal{B}, m)$ is a Banach space. If $L^p_R(X, \mathcal{B}, m)$ denotes those equivalence classes containing real-valued functions then $L^p_R(X, \mathcal{B}, m)$ is a real Banach space. The bounded measurable functions are dense in $L^p(X, \mathcal{B}, m)$. If $m(X) < \infty$ and $1 \leq p < q$ then $L^q(X, \mathcal{B}, m) \subset L^p(X, \mathcal{B}, m)$. We sometimes write $L^p(m)$ or $L^p(\mathcal{B})$ instead of $L^p(X, \mathcal{B}, m)$ when no confusion can arise.

A Hilbert space \mathcal{H} is a Banach space in which the norm is given by an inner product, i.e., \mathcal{H} is a Banach space and there is a map $(\cdot, \cdot): \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ such that (\cdot, \cdot) is bilinear, $(f, g) = (\bar{g}, f) \forall g, f \in \mathcal{H}$, $(f, f) \geq 0 \forall f \in \mathcal{H}$, and $\|f\| = (f, f)^{1/2}$ is the norm on \mathcal{H} .