

*Hollinshead*

Functional Anatomy  
of the  
Limbs and Back

# the Limbs and Back

*W. Henry Hollinshead, Ph.D.*

*Head of the Section of Anatomy, Mayo Clinic, Rochester;  
Professor of Anatomy, Mayo Foundation,  
University of Minnesota*

*Second Edition*

*Illustrated*

© 1960 by W. B. Saunders Company

*Copyright, 1951 by W. B. Saunders Company*

Copyright under the International Copyright Union

*All rights reserved.* This book is protected by copyright. No part of it may be duplicated or reproduced in any manner without written permission from the publisher. *Made in the United States of America at the press of W. B. Saunders Company.*

*Library of Congress Catalog Card Number: 60-5420*

# Preface to the Second Edition

In revising *Functional Anatomy of the Limbs and Back* I have kept in mind its original purpose, to provide a readable account of the anatomy of most concern to those particularly interested in the functions of muscles and the movements of the body, with special reference to the needs of the student of physical therapy. The division of the book into five sections, of which the first and the last present general information deemed useful to the student while the three intervening sections contain more detailed information on the limbs and back, has therefore been retained.

Within this general framework the revision of both text and figures has been thorough. The N.A. (Nomina Anatomica) of 1955 has been used throughout, thus eliminating the necessity for the beginning student to burden his memory with synonyms, and only when the student might obviously be handicapped by ignorance of an older but now obsolete term has that been given. (With few exceptions, however, N.A. terms are readily understandable to those who are already familiar with the B.N.A.) With the beginning student in mind, more pains have

been taken in the explanation of terms, cross references to both figures and text have been added where it seemed they might help in understanding the text, and parts of the text have been rearranged and when necessary rewritten to fit it better with the study of laboratory material. The more important descriptive chapters have also had added to them brief reviews through which the student can consolidate the knowledge gained by observation in the laboratory, and review the surface anatomy of the part studied. As a part of these, tables of the nerves and muscles are included.

New also are a number of figures, most of them showing vessels and nerves. Figure 93 is borrowed from *Clinical Examinations in Neurology*, by the Sections of Neurology and the Section of Physiology, Mayo Clinic and Mayo Foundation, W. B. Saunders Co., 1956. Figures 19, 25-27, 29, 30, 44, 49-53, 57, 60, 61, 67, 71, 74-76, 96-98, 101, 103, 120, 123, 128-130, 132, 137, 141, 147, 150, 154, and 156 have been taken from my larger three-volume work, *Anatomy for Surgeons* (New York, Paul B. Hoeber Inc., 1954, 1956, and 1958), and I am much indebted to its publishers for permission to use them here.

My thanks are due to a number of persons who have been kind enough to call my attention to errors in the first edition or to offer suggestions for this one. I believe I have made good use of their criticism, and hope that such helpful letters and comments continue to come in. I am particularly indebted to Mr. Darrel D. Hunt, Educational Administrator, and Mr. Dale V. Shaffer, Instructor, of the School of Physical Therapy, Mayo Clinic, who have offered many suggestions on the basis of their use of the book, and both of whom have read for me the entire text.

W. HENRY HOLLINSHEAD

*Rochester, Minnesota*

# Contents

## SECTION I. THE ORGANIZATION OF THE BODY

1. ANATOMICAL TERMINOLOGY . . . . .	3
2. THE TISSUES OF THE BODY . . . . .	8
3. THE ORGANS AND ORGAN SYSTEMS . . . . .	22

## SECTION II. THE UPPER LIMB

4. GENERAL SURVEY OF THE UPPER LIMB . . . . .	71
5. THE SHOULDER . . . . .	77
6. THE ARM . . . . .	117
7. FOREARM AND HAND: GENERAL CONSIDERATIONS . . . . .	132
8. FLEXOR FOREARM . . . . .	141
9. EXTENSOR FOREARM . . . . .	152
10. RADIO-ULNAR AND WRIST MOVEMENTS . . . . .	161
11. THE HAND . . . . .	169
12. MOVEMENTS OF THE DIGITS . . . . .	189

**SECTION III. THE BACK**

13. THE BACK . . . . .	199
------------------------	-----

**SECTION IV. THE LOWER LIMB**

14. GENERAL SURVEY OF THE LOWER LIMB . . . . .	229
15. THE BONY PELVIS, FEMUR, AND HIP JOINT . . . . .	235
16. THE THIGH AND KNEE . . . . .	242
17. BUTTOCK AND POSTERIOR THIGH . . . . .	263
18. MOVEMENTS OF THE THIGH AND LEG . . . . .	278
19. THE LEG . . . . .	294
20. THE FOOT . . . . .	319

**SECTION V. THE HEAD, NECK, AND TRUNK**

21. THE HEAD AND NECK . . . . .	341
22. THE THORAX . . . . .	365
23. THE ABDOMEN . . . . .	373

<b>SYNONYMS . . . . .</b>	<b>387</b>
---------------------------	------------

<b>INDEX . . . . .</b>	<b>391</b>
------------------------	------------

# The Organization of the Body





# Anatomical Terminology

The beginning student of anatomy is confronted with the necessity of mastering a largely new, cumbersome and complicated vocabulary. The difficulties are increased by the fact that, in common with most scientific terminology, anatomical names are given a Latin form. In this country we tend to depart from the strict Latin by anglicizing many expressions or using a direct English translation, but the Latin terminology is nevertheless the foundation upon which our scientific vocabulary is built. The problem is further complicated for most of us through the fact that, regardless of their endings, most anatomical terms have Greek or Latin roots. As most of them convey very definite meanings it is well worth the effort to consult a medical dictionary, when necessary, to discover the original meaning of the word and thus translate it in one's mind from a term that must be merely memorized to one that is understood. If the student will make a conscientious effort to understand the terminology of anatomy, he will find it much easier to learn the facts and concepts of anatomy.

A further difficulty in anatomical terminology is the abundance of synonyms that have accumulated for generations. The new internationally adopted anatomical terminology recognizes practically no synonyms, but even anatomists, because they are used to an older

terminology, have difficulty in eliminating synonyms from their talking and writing. Certainly, many of the clinicians with whom the therapist will come in contact will use the synonyms with which they are most familiar, and it is impossible to eliminate synonyms from older texts and figures that the student may consult. In this book the use of synonyms has been reduced to an approximate minimum. Only where it seemed obvious that the student might be handicapped through ignorance of another commonly employed although officially outdated term has that term been given as a synonym.

The major subdivisions of the body are the head, neck, trunk, and limbs. Although it is perfectly proper to use these English names, they also have Latin names that are used in many terms that the student will meet. Thus the head is "caput," and "capitis" therefore means "of the head." The neck is "collum"; cervix also means neck, especially its anterior part, and nucha means its posterior part. Thus "colli" means "of the neck," and we also find the terms "cervical" and "nuchal" or "nuchae" used in referring to structures in the neck.

Our word "trunk" is obviously the same as the Latin "truncus," but we have no particular reason to use the latter word. However, the subdivisions of the trunk need to be understood. The Latin word for chest is thorax, and this word will be met often both in this form, in its possessive "thoracis," which means "of the chest," and in its adjectival form "thoracic." The abdomen is the part of the trunk with muscular walls that lies below the thorax. It is most easily translated into our word "belly," but since this is considered inelegant we are left with no acceptable translation, and therefore use the words "abdomen" and "abdominal." The supposed synonym "stomach," as used popularly in "stomach ache" or "shot in the stomach," is not really a synonym; the stomach is one particular part of the digestive tract, almost all of which lies in the abdomen. The lowest part of the trunk is the pelvis (meaning "basin") and the Latin term is always used for this; "pelvic" is the adjective pertaining to the pelvis.

The Latin for the limbs is "membra" (member), but has no common usage. "Appendage" is a term long used by zoologists to describe limbs in general, and sometimes appears in human anatomy in the adjectival form "appendicular." Names of smaller subdivisions of the limbs are best reserved for the times when the limbs are studied.

Since anatomy is a descriptive science, many anatomical terms are in themselves descriptive, referring to shape, size, location or function of a part, or its fancied resemblance to some nonanatomical structure. Muscles, especially, are usually so named, but since many muscles may have a given shape or a given function it is usual to use more than one descriptive adjective in naming a muscle. Thus there are two muscles called biceps, or muscles with two heads of origin, so we distinguish one as the biceps brachii, or two-headed muscle of the arm, the other

as the biceps femoris, or two-headed muscle of the thigh. Similarly, the pronator quadratus is a quadrilateral muscle that pronates or turns the palm of the hand backward, the quadratus lumborum is a quadrilateral muscle located in the lumbar region; the rectus abdominis is a muscle running vertically (rectus means straight) in the abdominal wall, the rectus capitis anterior is an anteriorly situated muscle running vertically to attach to the caput, or head, and so forth.

In addition to the various technical names of structures in the body there are also certain general terms describing surfaces of the body, planes through the body, relative positions of one structure to another, and so forth, that must be understood from the very beginning. The surfaces of the trunk may be conveniently described as the dorsum, or back, the ventral or belly surface, and the two sides, or lateral surfaces. The cranium is the skull, and cephalon is the Greek word for head, so cranial and cephalic both mean toward the head; similarly, caudal means toward the tail, or in the human being toward where the tail would be had it persisted from embryonic life.

The above-defined terms are all understandable regardless of the position of the body, but there are others that require agreement as to what position of the body we are referring to before they can be understood. For instance, superior, meaning up or upward, implies a relation to gravity, and therefore might differ entirely in meaning according to whether one was erect, lying upon one's back, or standing upon one's head. For this reason, anatomists have agreed that such terms of relative position should always be used in relation to a fixed position of the body termed the anatomical position; the anatomical position is the erect one, with the heels together and the feet pointing somewhat outward, the arms by the sides and the palms facing forward. With reference, then, to the anatomical position, superior always means toward the head, and is therefore used interchangeably with cephalic or cranial; similarly, inferior means toward the feet, and is usually synonymous with caudal. Anterior, referring to the part of the body habitually carried forward in progression, is thus synonymous with ventral in the human being, and posterior and dorsal are also synonymous. Even in the new anatomical terminology there is some inconsistency in the use of the terms "anterior" and "ventral," and similarly in "posterior" and "dorsal"; in essence, however, dorsal and ventral are used in human anatomy only in referring to parts of spinal nerves and of the hand and foot (compare our English "back of the hand"). Elsewhere anterior and posterior are preferred, so what was once, for example, the dorsal interosseous artery is now the posterior interosseous artery. The term "ventral" was rarely used in regard to the limbs (except in developmental stages), the more common term here being "volar" (vola = the palm or sole); this has now been abandoned as the

antithesis of posterior, so what was once the volar interosseous artery is now called the anterior interosseous artery.

Other terms of relative position are medial and lateral, that is, toward the midline or toward a side. In the case of the limbs, however, some confusion might exist as to whether these terms refer to the body as a whole or to the limb itself; thus the little finger is medial to the other fingers in regard to the midline of the body, but is lateral, as is the thumb, in regard to the midline of the hand. It is best in this case, therefore, to use medial and lateral only when referring to the limb as a whole, and its relation to the body in the anatomical position; if relative mediolateral relationships of structures within the limb are to be described, radial and ulnar, and tibial and fibular are the terms best employed. These terms refer to the paired bones of the limbs. In the upper limb, the radius is on the thumb side, the ulna on the little finger side, and in the lower limb the tibia is on the side of the big toe while the fibula is on that of the little toe; the sides of the limbs are thus named from the positions of these bones. Additional terms especially useful in regard to the limbs are proximal, meaning toward the attachment of the limb to the trunk; distal, or away from the base of the limb; palmar, referring to the palm of the hand; and plantar, referring to the sole of the foot.

In regard to the planes of the body, the sagittal plane is one passing through the midline of the body so as to divide it into right and left halves; a parasagittal plane is one parallel to the sagittal, but to one side of it. A coronal or frontal plane is one dividing the body into an anterior and a posterior (ventral and dorsal) portion, thus running roughly parallel with the front of the body and with a suture of the skull called the coronal suture. A horizontal or transverse plane divides the body or limbs into upper and lower parts, in relation to gravity and the anatomical position.

In describing movements we may refer to them as being toward or away from a given plane, or we may find it convenient to speak of the axis of motion. For any given movement the axis of motion is an imaginary line about which a part describes a rotatory movement. These axes of motion typically make angles with one of the chief planes of the body. For instance, the axis for bringing the forearm up against the arm ("bending" the arm) is a line passing through the elbow region from lateral to medial sides, and is therefore at approximate right angles to the sagittal plane, and correspondingly approximately parallel to the frontal plane.

In addition to the terms of position, we must also agree in general upon the meaning of terms describing movement. In general, bending of the trunk or the limbs occurs most freely toward the original ventral surfaces of these members, therefore flexion, which means bending, is usually a bending in the ventral direction; if we want to designate a

bending as being in the dorsal direction, it is proper to say dorsiflexion. Extension, the straightening out of a bent part, is a movement opposed to that of flexion, and therefore occurs typically in the dorsal direction; if, at the joints where this is possible, we continue the movement of extension beyond that necessary to straighten the part, we can designate it as hyperextension. Dorsiflexion and hyperextension are plainly synonymous, as they both designate movement in the same direction. From these terms of movement we have acquired terms of position that are frequently used colloquially, although not a part of official terminology. Thus we may speak of the flexor or extensor surfaces of a limb, and these correspond to the original ventral and dorsal surfaces of the developing limb.

Abduction means moving apart, or away from the midline, and adduction, the reverse, is moving together, or toward the midline; both of these are particularly useful in describing movements of the limbs. Protraction is moving a part forward, retraction moving it back; elevation is lifting a part, that is, moving it superiorly, while depression is moving it inferiorly. Rotation is the twisting of a part around its longitudinal axis, and is described as lateral or external rotation if the anterior surface of the part is turned laterally, and medial or internal rotation if it is turned medially. Circumduction is a combination of successive movements of flexion, abduction, extension and adduction in such a fashion that the distal end of the part being moved describes a circle.

In relation to the limbs, especially, some of the terms defined above are apt to be confusing in their specific applications. For this reason an effort has been made in the following chapters of this book to so define terms of relation and movement, as they apply specifically to the part being considered, that there can be no ambiguity as to what is meant. With the proper use of the dictionary, and with due regard to the correct usage of anatomical terms as displayed by your instructor and in textbooks, the bugaboo of anatomical terminology can be largely overcome, and what may seem at first an almost incomprehensible jargon will become a useful part of your scientific language.

# The Tissues of the Body

The human body, like most of the better-organized forms of animal life, consists of various types of specialized cells and a varying amount of intercellular substance; much of the actual weight of the body is water, both within the cells and without. The cells represent the living portion of the organism, while the intercellular substance, regardless of its nature, represents nonliving material that owes its existence to the activities of the cells.

Most types of cells tend to occur in groups in which the constituents are somewhat similar both in appearance and in function; such organized groups of cells are known as tissues. The tissues of the body are in turn not independent of one another, but rather various types of tissues are interwoven to form more complex anatomical and functional units known as organs or organ systems.


According to their general appearance and functions, the various tissues of the body are usually classified into a few great groups: epithelial tissue, connective tissue, muscular tissue, nervous tissue, and blood.

## *Epithelia*

Epithelial tissue (fig. 1) occurs most commonly in sheets, and is adapted especially for covering other tissues. It serves the general func-

tions of protection, absorption and secretion. An epithelium is characterized by the fact that its cells are closely packed together with a minimum of inert intercellular cement substance between them. The cells may vary in shape from extremely flat ones resembling paving stones (called squamous cells; squama means a scale, such as that of a fish), to cuboidal ones shaped somewhat like children's blocks, to tall columnar ones. There are many subvarieties in shape, general appearance and function.

An epithelium may take the form of a single-layered sheet of cells, a multilayered sheet, or of essentially tubular outgrowths (glands) from such sheets. One type of epithelium covers the external surface of the body where, as the outer layer of the skin, a stratified or multilayered epithelium with dead outer cells protects the more delicate deeper-lying cells and serves as a membrane to seal off intercellular spaces from contact with the outside. Another type of epithelium, adapted for absorption and secretion, lines the digestive tract, and outgrowths from this epithelium form the digestive glands, including also the characteristic cells of such large organs as the liver and pancreas. Other types of epithelium line the tubules of the kidneys, the ureters, and the urinary bladder, and of course continue along the urethra (the tube leading from the bladder) to unite with the epithelium of the skin. Thus epithelium occurs primarily either on the outside of the body or as a lining of those cavities of the body that communicate with the exterior.



*Figure 1.* Several types of epithelium. *a.* Simple squamous, surface view. *b.* Simple cuboidal. *c.* Ciliated columnar. *d.* Stratified squamous. *b, c,* and *d* represent views of sections.



Two specialized types of epithelium are also found lining closed cavities within the body. Mesothelium, a single-layered very flattened epithelium of the pavement or squamous type, lines the four great cavities of the trunk, namely, the two pleural cavities surrounding the lungs, the pericardial cavity surrounding the heart, and the peritoneal cavity surrounding the abdominal viscera. Endothelium, essentially similar to mesothelium in appearance, forms the inner lining of the heart, of all blood vessels, and of the lymphatics.

### **Connective Tissue**

Connective tissue stands in the sharpest contrast to epithelial tissue in its composition, since in connective tissue the cells are more or less widely dispersed and separated from each other by nonliving intercellular material; it is the presence and character of this intercellular material that gives connective tissue its specific characteristics.

The most pervasive type of connective tissue in the body is **fibrous connective tissue** (fig. 2). In this type the spaces between the cells are occupied by numerous fibers that make the tissues tough and capable of withstanding distortions and strains. The fibers between the cells may be of several types, and may occur either in the form of a loosely woven net with large quantities of fluid in the interstices of the net, or they may be closely packed with little interfibrillar space. The most common type of fiber found in connective tissues is the collagenous fiber. These fibers are nonelastic; therefore, when they occur in places in which some deformation must be possible they are arranged in wavy bundles that allow movement until the slack of these bundles is taken up. Elastic fibers are the other important type of intercellular fibers; these fibers are actually elastic as their name implies: they may be stretched, and when the tension upon them is relaxed, they will shorten again. Their cut ends often curl. They occur frequently mixed in with more numerous collagenous fibers, but in certain locations great bundles of almost pure elastic tissue are found. At intervals, in the interstices between connective tissue fibers, connective tissue cells occur. The most important of these, known as fibroblasts, are responsible for the formation and repair of the connective tissue fibers. Others possess the property of ingesting formed material; in this duty they may be aided by cells from the blood, some of which pass freely into the fibrous connective tissues as a part of the reaction of inflammation.

Collagenous tissue, with or without the admixture of elastic fibers, is the most ubiquitous of all tissues. Taking various forms, it permeates and surrounds practically all the tissues of the body, serving as a binding agent for these tissues. Indeed, it has been aptly said that if it were possible to dissolve out all of the tissues of the body so as to leave only the fibrous connective tissues, the essential organization of the body