Monographs in Virology

Vol. 7

The Agent of Trachoma

Recent Studies on the Biology, Biochemistry and Immunology of a Prokaryotic Obligate Parasite of Eukaryocytes

YECHIEL BECKER

The Agent of Trachoma

Recent Studies on the Biology, Biochemistry and Immunology of a Prokaryotic Obligate Parasite of Eukaryocytes

YECHIEL BECKER

Laboratory for Molecular Virology, Institute of Microbiology, Hebrew University-Hadassah Medical School, Jerusalem

12 figures and 5 tables

session and straight of our to be are as to 177



Monographs in Virology

Vol. 3

and hamagology of a Probaryous Obligate Parasite

HOTCHIN, J. (Albany, N.Y.): Persistent and Slow Virus Infections X + 210 p., 26 fig., 11 tab., 1971

ISBN 3-8055-1176-0

Vol. 4

Neurath, A. Robert and Rubin, Benjamin A. (Philadelphia, Pa.): Viral Structural Components as Immunogens of Prophylactic Value VI + 87 p., 11 fig., 1 cpl., 1971 ISBN 3-8055-1203-1

Vol. 5

WILDY, P. (Birmingham): Classification and Nomenclature of Viruses VIII + 82 p., 1971 ISBN 3-8055-1196-5

Vol. 6

TINSLEY, FHOMAS W. and HARRAP, KEITH A. (Oxford):

Moving Frontiers in Invertebrate Virology

VIII + 66 p., 23 fig., 3 tab., 1972

ISBN 3805-5-1464-6

S. Karger · Basel · München · Paris · London · New York · Sydney Arnold-Böcklin-Strasse 25, CH-4011 Basel (Switzerland)

All rights, including that of translation into other languages, reserved.

Photomechanic reproduction (photocopy, microcopy) of this book or parts thereof without special permission of the publishers is prohibited.

Copyright 1974 by S. Karger AG, Verlag für Medizin und Naturwissenschaften, Basel Printed in Switzerland by Buchdruckerei Merkur AG, Langenthal ISBN 3-8055-1657-6

Editor's Preface Faitor's Profines

Although the agent of trachoma has now been generally accepted as nonviral, it was for so long considered a virus that it seems appropriate to include this review as a Monograph in Virology. Hence, this volume should be useful not only to those working directly with the trachoma agents, but also to virologists who need an overview of the developments during the last decade which have distinguished these chlamydial agents from the viruses. JOSEPH L. MELNICK

blobers, because the complete parties on Alleger within the contract of the assess and an oran manufacture had a their armanent. Or full entire

Editor's Preface

Although the agent of trachoma has now been generally accepted as non-viral, it was for so long considered a virus that it seems appropriate to include this review as a Monograph in Virology. Hence, this volume should be useful not only to those working directly with the trachoma agents, but also to virologists who need an overview of the developments during the last decade which have distinguished these chlamydial agents from the viruses.

Altenush the agest of true points has now been generally pointpled as non-

JOSEPH L. MELNICK

The review of the floorager was completed July 1972. However, come

The opportunity to write a review on the recent developments in the research on the agent of trachoma had arisen from the invitation of Professor J. L. Melnick to revise the review written in 1962, by the late Professor Hans Bernkopf. As a result of marked developments in research on the immunology and biochemistry of the agent, a new review which will compile the knowledge added in the last decade seemed necessary. I have, therefore, referred mainly to studies on *Chlamydia trachomatis* which were published during the last decade. The very important studies on *Chlamydia psittaci* are beyond the scope of this monograph.

I wish to thank my colleagues, Professor I. C. MICHAELSON and Dr. B. MAYTHAR from the Department of Ophthalmology, Hadassah University Hospital, Jerusalem; Drs. Z. ZAKAY-RONES, I. SAROV, from the Department of Virology, B. GUTTER, and Y. ASHER, Y. COHEN, and H. LOKER from the Laboratory for Molecular Virology, for their continuous and enthusiastic collaboration; Professor S. E. Luria, Department of Biology, Massachusetts Institute of Technology, Cambridge, Mass., for the critical discussions on the nature of obligate parasitism of viruses and prokaryotes in eukaryotic cells; Professor E. JAWETZ, Department of Microbiology, University of California, San Francisco, Calif., for his continuous interest and support; Dr. M. L. TARIZZO, World Health Organization, Geneva, for his continuous interest and help; Professor A. L. BARRON, on sabbatical leave from the Department of Microbiology, School of Medicine, State University of Buffalo, N.Y.; Professor H. M. JENKIN, Department of Microbiology, Hormel Institute, University of Minnesota, Austin, Minn., for the critical reading of the manuscript and for their comments; Dr. Julia Hadar for proofreading the manuscript; and my wife Miriam Becker for her continuous encouragement.

The preparation of the review was supported by a grant (No. 06-501-1.38) from the National Library of Medicine, USA, to the Israel Journal of Medical Sciences.

The research on trachoma in the author's laboratory was supported by grants from the National Institutes of Health, Bethesda, Md., USA, and the World Health Organization, Geneva, Switzerland.

The review of the literature was completed July 1972. However, some manuscripts published during 1973, while the Monograph was prepared, are also included.

the apportunity is write a leview on the count developments in the

- Lastinto that any collecture. Professor L.C. Market son and Dr. R.

treat of Virology, R. Gross, and Y. Asiro, Y. Court, and H. Lotter

Manush and institute of Lexandrey, Cambridge, Massel for the critical discussions on the nature of obligate parasition of curious and prokeryotes

loave from the Department of Memoricology, School of Medicine, State University of Bulleton, N. V., or George M. M., have to Department of Apertment of Apertment of Apertment of Apertment of the control of the comments; i.e. Turn the Location for produced as a few comments; i.e. Turn the Location for produced as the control of the con

Contents

the Adequation of Continuents of

4. Commission Presugential of Littlement N. Co.

Editor's Preface	. XII
Editor's Preface Preface	. XIII
III United being Pt III to make the world to be the members of the	17
I. Introduction	
B. Recent Developments	
II. Definitions and Classification of the Chlamydia	
A. Definitions of the Chlamydia	4
The Agents of Psittacosis, Lymphogranuloma Venereum and Trachom	19
(PLT) – Members of the Order Chlamydiales	
2. The Classification of the Chlamydia (PLT) Agents	
3. The Non-Viral Nature of the Chlamydia	The second second
4. The Bacterial Aspects of the Chlamydia	
B. Relationship between C. trachomatis and C. psittaci	
Division of Chlamydia into Two Distinct Subgroups	
a) Based on Biological Properties	
b) Based on Susceptibility to Sulfadiazine and p-Cycloserine	
c) Based on Antigenicity	
d) Based on DNA Homology	
2. Further Subdivision of the C. trachomatis Agents	
C. Evolution of the Chlamydial Agents	
1. Evolutionary Relationship between C. trachomatis and C. psittaci	
2. Evolution of C. trachomatis Strains	
III. Isolation and Propagation of TRIC and Related Agents	
A. Isolation of TRIC Agents from the Conjunctivae of Trachoma Patients .	
1. Difficulties in the Diagnosis of Trachoma and Inclusion Conjunctivitis	
2. Isolation of TRIC Agents in the Yolk Sac of Embryonated Eggs	13
a) Studies by T'ANG and Collaborators	13
b) Further Isolations of Agents from the Conjunctivae of Trachon	
Patients	
c) Factors Affecting the Isolation of Trachoma Strains	
B. Direct Isolation of TRIC Agents in Cultured Cells	14
1. The Use of McCoy Cells for the Isolation of the Trachoma Agents	
2. Isolation of TRIC Agents from Human Disorders	15
a) Isolation of Trachoma Strains from the Urogenital Tract and the	he
Rectum	15
b) Isolation of Chlamydia from LGV and Other Human Disorders .	16

VI Contents

C. Growth of C. trachomatis in Cultured Tissues	16
1. Explants of Chick Embryos	16
2. Adaptation of C. trachomatis Agents to Growth in Conjunctival Cells in	
vitro	16
3. Adaptation of Trachoma to Mouse Macrophages	17
4. Continuous Propagation of Chlamydia in Cell Lines	17
5. Enhancement of Trachoma Agent Infectivity by DEAE-Dextran	17
D. Properties of the T'ang (TE55) Strain	18
1. Early Passages of the TE55 Strain in Chick Embryos	18
2. Adaptation of TE55 to Growth in Cell Cultures	18
3. Properties of the TE55 Strain Propagated in Other Cell Lines	19
4. Propagation of TE55 in Mice	19
5. Development of Fast Killing (f) Strains of TE55 and Related Agents	20
6. Antigenic Differences of TE 55 Strains	21
7. Antigenic Properties of the T'ang TE55 Strain	21
IV. The Biology of the Trachoma Agent	23
A. The Developmental Cycle	23
1. The Interaction of e.b. with Host Cells	23
2. Site of Development in the Infected Cell	24
3. The Biological Activity of Trachoma Elementary Bodies - Determination	-
of Infectivity	24
a) Titration in Embryonated Eggs	24
b) Titration in Cultured Cells	
4. Stages of the Developmental Cycle	25
B. Factors Affecting C. trachomatis Development	26
Nutritional Factors 1. Nutritional Factors	26
	26
a) 1 mmo 1 m	27
b) Serum 2. Temperature of Incubation	27
3. The Host Cell	
C. Effects of C. trachomatis infection on the flost Cen	28
1. Effect off Flost Cell Metabolishi	28
2. Induction of Interferon Synthesis	
D. Morphology of the Developing C. trachomatis Inclusion Bodies	
1. Correlation octween wiorphology and infectivity	29
2. Morphology of C. trachomatis Inclusions during the Noninfective Phase	
3. The Morphology of C. trachomatis Inclusion Bodies during the Infectivity-	
Increase Phase	30
4. Comparison between C. trachomatis Inclusions in Cultured Cells and	
Conjunctival Cells from Trachoma Patients	30
5. Isolation and Characterization of C. trachomatis Initial Bodies	
a) Sucrose Gradient Analysis	30
b) Characterization of the Initial Bodies	32
6. Initial Bodies Are Precursors to Elementary Bodies. The Transition of the	
Initial Bodies to Elementary Bodies	
V. The Composition of Trachoma Flementary Bodies (e.b.)	34

		Contents	VI
		Tooletion and Duriffersion of a b	2
	A.	Isolation and Purification of e.b. 1. Purification of e.b. from Infected Yolk Sacs	34
			34
		a) Sucrose Gradient Centrifugation	34
		b) DEAE-Sephadex Column Chromatography	34
		c) Genetron Treatment	
	_		35
	B.	The Chemical Composition of Trachoma e.b.	35
Ē.		1. The DNA Genome	35
		a) The Molecular Weight of the DNA Genome	35
		b) Properties of C. trachomatis DNA	36
		2. RNA in the e.b	37
ľ		a) Ribosomal Subunits	37
ì		b) Ribosomal RNA	37
		c) Transfer RNA	
		d) Messenger RNA	31
Š		3. Proteins in e.b	38
		4. Lipids in the e.b.	38
		5. Cell Membranes and Cell Walls in Trachoma e.b.	39
	C.	Antigenic Composition of Trachoma e.b.	39
		1. Antigens	
Č		2. Chemical Composition of the Group-Specific Antigen	39
à		3. Hemagglutinin	40
		4. Toxin	40
į	D.	Enzymatic Activities in Trachoma e.b.	40
		1. DNA-Dependent RNA Polymerase	40
		2. Polynucleotide Phosphorylase	42
		3. Enzymes for Glucose Metabolism	42
		4. Transaminase Activity	42
I.	Mo	olecular Aspects of the Developmental Cycle of C. trachomatis	43
k	A.	Initiation of e.b. Development after Infection of Eukaryotic Cells	43
		1. Initiation of RNA Synthesis in the e.b.	43
		2. Activation of the e.bAssociated DNA-Dependent RNA Polymerase	43
		3. The Role of Host Cell Mitochondria in e.b. Development	43
		4. Initiation of Protein Synthesis in the e.b.	44
Š	B.	RNA Synthesis during the Trachoma Developmental Eycle	44
		1. Cytoplasmic Site of RNA Synthesis in the C. trachomatis-Infected Cell .	44
100		2. Rate of C. trachomatis RNA Synthesis	44
		3. Synthesis of Precursors to Ribosomal RNA	45
3		4. Maturation of Trachoma Ribosomal RNA	40
		75. Synthesis of Trachoma 5S and 4S RNA Species	4
į		6. RNA Synthesis in the Initial (Secondary) Bodies	4
		7. DNA-Dependent RNA Polymerase in the Purified Initial Bodies	49
	C.	DNA Synthesis	49
þ		Time Course of DNA Synthesis	49
		2 Site of DNA Synthesis	49
		3. DNA Synthesis and Morphogenesis	- 50
į		Sulforesteld: Traction in first and Area and Are	
		그림을 하는 것이 보고 있다. 이번 사람들이 되었다면 가장 하는 것이 없었다면 하는데 없지 않는데 없다면 하는데 없다면 하는데 없다면	

·VIII Contents

	D.	Protein Synthesis	50
		1. Time Course of C. trachomatis Protein Synthesis	50
ME.		2. Characterization of Trachoma Proteins Synthesized in Infected Cells	51
NE.	E.	Glycogen Synthesis	51
		1. Demonstration of Glycogen in the Inclusions of C. trachomatis	51
		2. Relationship between Glycogen Synthesis and the C. trachomatis Growth	
355		Cycle	52
35		3. Effect of Inhibitors on Glycogen Biosynthesis	52
at		a) Effect of Penicillin	52
		b) Effect of Fluorouracil on Glycogen Synthesis and Degradation	53
		c) Effect of Phlorizin	53
12		4. Glycogen Synthetase	53
	F.	Lipid Metabolism and Membrane Biosynthesis	54
	•	1. ³ H-Choline Incorporation into Developing Inclusion Bodies	54
SE		2. The Inclusion-Body Limiting Membrane	54
w		3. The Membranes of the Initial and Elementary Bodies	54
05	G	Cell Wall Synthesis	55
VI		ntibiotics against C. trachomatis and Trachoma	56
RE		Mode of Action of Antitrachoma Drugs	56
92	A.		56
		Sulfonamides	56
			56
		b) Mode of Action of Sulfa Drugs c) Mutations to Sulfonamide-Resistant Trachoma Strains	57
4			-
		2. Inhibitors of Folic Acid Metabolism	58
	73	a) Aminopterin	58
		b) Trimethoprim	58
	В.	Mode of Antitrachoma Action of Antibiotics	58
23		1. Penicillin	58
-EA		a) Inhibition of Trachoma Development	58
E.P.		b) Mode of Action	59
-53-		2. Tetracyclines	59
		3. Chloramphenicol	60
75		4. Rifampicin	61
- 64		a) Effect on Trachoma Development	
, M		b) Irreversibility of Rifampicin Inhibition	61
- 80	1	c) Mode of Action of Rifampicin	62
45		d) Inhibitory Effects by Rifampicin Derivatives	62
		5. Streptovaricin	63
173		6. Macrolides	63
		7. Hydroxyurea	63
		8. Metabolic Analogs	64
-85		9. Antibiotics Inhibitory at High Concentrations	64
-69	C	Drugs and Antibiotics in the Treatment of Trachoma Patients	64
		1. Treatment with Sulfonamides	65
		a) Long-Acting Sulfonamides	65
		b) Sulfonamide Treatment in Endemic Areas	65

此为试读,需要完整PDF请访问:,www.ertongbook.co

Contents	IX

			of Actification Veryletter Total Chiesian Control	
		2.	Treatment with Antibiotics	65
		3.	Doxycycline Treatment of Patients with Chronic Trachoma	66
	D.	Eff	ect of Interferon on Experimental Trachoma Infections	66
		1.	Effect of Interferon on Trachoma Development	66
		2.	Effect of Interferon Inducers on Infection	66
VII	II. I	mm	nunobiology	67
			munological Responses to the Trachoma Agent	67
			Local Antibodies in Humans	67
			a) Detection of Antibodies	67
			b) Characterization of the Antitrachoma Antibodies	67
			c) The Role of Antitrachoma Antibodies in the Eyes	68
		2.	Local Antibodies in Experimental Infections of Animals	68
			a) Antibody Response to Trachoma Antigens in Guinea Pig Eyes	68
			b) Local Antibodies in Monkeys' Eyes	68
		3.	Antitrachoma Antibodies in Human Sera	69
			a) Serum Antibodies in Trachoma Patients	69
			b) Serum Antibodies in Experimental Animals	69
			c) Effect of Route of Immunization on the Antibodies in Guinea Pigs'	
			Eyes	70
		4.	Delayed Hypersensitivity	70
			a) Role in the Pathogenesis of the Disease	70
			b) In Humans	70
			c) Experimental Delayed Hypersensitivity	70
			d) Induction of Delayed Hypersensitivity in Monkeys	71
			e) Differentiation of Chlamydial Agents by a Delayed-Type Hyper-	
			sensitivity Test	71
	B.	Ch	aracterization of Antitrachoma Antibodies	71
h		1.	Neutralizing Antibodies	71
			a) Neutralization of Homologous Strains	71
			b) Neutralization of Heterologous Strains	72
		2.	Antibodies to Specific Antigens	72
			a) Antibodies to Group-Specific Antigen	72
			b) Antibodies to Cell Wall Antigens	72
	C.		perimental Immunization against Trachoma	73
		1.	Immunological Responses to Trachoma Vaccines in Animals	73
			a) Guinea Pigs	73
			b) Owl Monkeys	73
		2.	Other Species of Monkeys	73
			a) Dissemination of Infective Trachoma Elementary Bodies in Baboons	73
			b) Importance of Content of Elementary Bodies in the Vaccine	74
			c) Immunization of Monkeys with a Trivalent Vaccine	74
		2	d) Conclusions	74
		3.	a) Antitrachoma Vaccination Trials in Taiwan	75
			b) Vaccination of Children in Punjab Indian Villages	75
			c) A Field Trial in Saudi Arabia	75 76
			C) A FICH IIIdi III DAUGI AIADIA	10

X	Contents
	d) Antitrachoma Vaccination Trials in Ethiopia
	e) Vaccinotherapy Studies in USSR
	f) Vaccination Trials in Gambia
IX.	Summary and Conclusions
X.	References
	Photosia in the control of the contr
10	The second secon
10.	the state of the party of the party of the state of the s
	Local Andbaco in Burdan
1.00	estindant to emestic, (g
	 2) Distriction of Application 3) Character flow of the anti-radional Application
	c) The Role of Antigrehoons antibooks in the Pierc.
83	stamma, to ending the transmission of the tage in
23	of Animals Removally Traditions Authority in the least the Elest.
	The second of th
John State	Andrew one activities in Touring 27
40	a) Serum Auri, idies in , rachoma for esta, A
to the	Martin A company of the Company of the Martin of the Marti
	from change of Rouge of Boundaries of the Authorities in Council to
	The court of Arterior markets of the court of
	Delived Hygoman analog
	to Role to the Publishers of the Contact of the Property of th
	the second of th
100	the formation of the second of the second of the
	of Industries of Delay of Name and Architecture of Architecture of Delay of Name and Architecture of the A
	as differentiation of Coloregial Aniche et a Cristiania in the control of the coloregian
	full this sections
	Bursclements of A - traceons with the For a street with
	pollodDnA grinteracki
	Spring analogue to management to
	Amibo as so Special Amigent
	at Ambodise to Great Secular Addition
	ASSISTA BUY II O TO COROCITIS, 18
TES	organization and the recommon along the forms
	between logical responses Treatment Vaccional Amortis.
	benieveloged reported Technica Vaccionia Amerika ki Gulini Pigi.
8.1	to the land of the second land to
	Citing Species of Montells
11	at Place of the Place of the Stone Place of the Interest.
	in many X-sa of Contain of Literatury further in the Youth Land
	in 2, 2 to graff a day term to be constituted to
	A Aportonia trializa e al meta Chicanorle da accordante do
	A PART THE STATE OF THE PART O

1. 150

I. Introduction

A. Historical Notes

Étiological studies on the trachoma agent date back to the observation by HALBERSTAEDTER and VON PROWAZEK [1907] who reported on the presence, in conjunctival cells, of inclusion bodies which contain elementary particles (designated 'Chlamydozoa' or mantle bodies). They believed, on the basis of resemblance in the morphology of the developmental cycle, that the particles in the inclusion bodies were the infectious agent of trachoma. The trachoma agent was grouped together with psittacosis and lymphogranuloma venereum into the psittacosis-lymphogranuloma-trachoma (PLT) group of organisms [Thygeson, 1962] now defined as members of the order Chlamydiales which includes one family Chlamydiaceae and two species Chlamydia trachomatis and Chlamydia psittaci (fig. 1) [PAGE, 1966, 1968, 1973]. However, 50 years elapsed from the discovery of the trachoma agent until the etiological agent of trachoma was isolated by T'ANG and his collaborators [T'ANG et al., 1957, 1958], propagated in embryonated eggs, and studied with respect to its properties. Although it has been known since 1938 that the agent of trachoma and similar agents of the PLT group resembled bacteria in their sensitivity to sulfa drugs and antibacterial antibiotics, these agents were, for many years, considered as viruses. The consideration of the trachoma agent as a virus resulted in a remarkable delay in the development of trachoma research. Only after the agents of trachoma were cultivated in vitro was the prokaryotic nature of the agent elucidated. In 1962, Bernkopf indicated in his review on trachoma that 'the more we learn of the nature of trachoma and related agents the less fitting the term "virus" becomes'. The indecisiveness in defining the PLT group is indicated in the constant changes in the name of the group of agents and its systematic position in the microbial world. In the 1962 symposium on the 'Biology of the Trachoma Agent', the PLT agents were regarded as viruses. In the 1966 conference on 'Trachoma and Allied Diseases', the agents were named Bedsonia. The name Chlamydia was used in the 1970 symposium on 'Tra-






Fig. 1. The chlamydia (PLT group of agents).

choma and Related Disorders Caused by Chlamydial Agents'. However, MOULDER [1966] clearly defined the psittacosis group of agents as microorganisms which are obligate intracellular parasites of eukaryotic cells. Recent developments in the research on the molecular biology of the trachoma agent have confirmed and extended MOULDER's ideas and furthered the understanding of the chlamydial agents as prokaryotic obligate parasites. In the present review, mainly studies on *C. trachomatis* will be reviewed.

B. Recent Developments

was his probuty offe nature of the agent clo-

Research on the trachoma agent has gone through several phases since the discovery by T'ANG et al. [1957, 1958]: the initial attempts were to isolate numerous strains of trachoma from patients in different parts of the world. This was followed by studies on the morphology and life cycle of the agent and on the effects of antibiotics on the trachoma agent. In the second phase, studies were concerned with the immunological properties of the trachoma

strains and attempts were made to develop potent antitrachoma vaccines. The developments in the research on the molecular biology of microorganisms made possible studies on the molecular processes which occur during the developmental cycle of the trachoma agent. These research activities are well reflected in the scientific reports presented in three international conferences (1962, 1966 and 1970)1 which were devoted to studies on the trachoma agent. The developments in the understanding of the chlamydia (trachoma and psittacosis agents) were reviewed by BERNKOPF [1962a, b], BIETTI and VOZZA [1963], JAWETZ [1964] and MOULDER [1966, 1971]. These reviews contain the basic information on the medical aspects of trachoma as well as the basic concepts of the chlamydial agents. The present review will deal with some of the developments in trachoma research which took place during the last decade (1962-1972)2. The aim of the present review is to present the current knowledge on the classification, molecular biology and immunological properties of C. trachomatis, the agents which cause trachoma, a disease of the human eye. These agents are members of a unique group of obligate microbial parasites of eukaryotic cells.

the index of the succession of the countries and pribates associated and the countries and the countries as the countries and the countries are considered as the countries are considered as

trees a reason temporar and Resona (\$10), continues (Rossean, 1940));

¹ a. The biology of the trachoma agent. Ann. N.Y. Acad. Sci 98 (1962). b. Conference on Trachoma and Allied Diseases. Amer. J. Ophthal. 63 (1967). c. NICHOLS R. L. (ed.): Trachoma and related disorders caused by chlamydial agents. Excerpta med. Int. Congr. Series, No. 223 (1971).

² Since the number of publications on trachoma is exceedingly high, it has become virtually impossible to quote all of them. Hence, only a relatively small number of the pertinent studies were included in the present review. A complete list of publications on trachoma is published by the Centre International de Documentation sur le Trachome, in the Revue Internationale du Trachome.

II. Definitions and Classification of the Chlamydia

developmental cycle of the traditions around These contents activities are well

(1962, 1966 and 1870) which core described to studies on the trudents a sont

(1963) Jan et a [1964] and Modittee [1996, 1971], those reviews control the

A. Definitions of the Chlamydia

The Agents of Psittacosis,

Lymphogranuloma Venereum and Trachoma (PLT) –

Members of the Order Chlamydiales

In contrast to most bacterial and viral agents, whose distinct nature was easily recognized, the members of the PLT group of agents were constantly shifted from one class to another until they resumed a position on the border between viruses and bacteria [RAKE et al., 1942; Weiss, 1955; MOULDER, 1966]. The indecisiveness regarding the nature of the agents and their relationship to other classes of biological entities was due to the absence of a clear definition of the group. The chlamydial agents were isolated in many laboratories during a period of 30 years from excretions of respiratory tracts (e.g. psittacosis [BEDSON and BLAND, 1932], ornithosis [Pollard, 1947]); from genital tracts (e.g. lymphogranuloma venereum [FINDLAY et al., 1938]); and from the eyes (e.g. trachoma [RAKE et al., 1942; T'ANG et al., 1957], inclusion conjunctivitis [FURNESS et al., 1960b]). The agents isolated from humans and other vertebrates were maintained by serial transfers in the yolk sacs of developing chick embryos. Some of the agents were later adapted to growth in cultured cells in vitro. Numerous studies on the isolated chlamydial agents [summarized by MEYER, 1952; Cox, 1953; THYGESON, 1953; WENNER, 1958; COLLIER, 1959a, b; LEPINE, 1960; BERNKOPF, 1962a, b; JAWETZ, 1964; MOULDER, 1966], as well as advancement in defining the class of viruses [LURIA, 1953; LWOFF, 1957; LWOFF and TOURNIER, 1966] and the recognition of the unique features of bacteria as a group [STANIER and VAN NIEL, 1962], led to the suggestion [PAGE, 1966, 1968] that the chlamydiae differ from Rickettsiae and constitute a specific order, Chlamydiales.