

国外数学名著系列(续一)

(影印版) 56

E. B. Vinberg (Ed.)

Geometry II

Spaces of Constant Curvature

几何 II

常曲率空间



科学出版社
www.sciencep.com

国外数学名著系列(影印版) 56

Geometry II

Spaces of Constant Curvature

几 何 II

常曲率空间

E. B. Vinberg (Ed.)

科 学 出 版 社

北 京

图字: 01-2008-5376

E. B. Vinberg(Ed.): Geometry II: Spaces of Constant Curvature

© Springer-Verlag Berlin Heidelberg 1993

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the People's Republic of China only and not for export therefrom

本书英文影印版由德国施普林格出版公司授权出版。未经出版者书面许可, 不得以任何方式复制或抄袭本书的任何部分。本书仅限在中华人民共和国销售, 不得出口。版权所有, 翻印必究。

图书在版编目(CIP)数据

几何 II: 常曲率空间 = Geometry II: Spaces of Constant Curvature/ (俄罗斯)温贝格(Vinberg, E. B.) 编著. —影印版. —北京: 科学出版社, 2009
(国外数学名著系列; 56)

ISBN 978-7-03-023499-5

I. 几… II. 温… III. ①几何-英文 ② 常曲率空间-英文 IV. O18

中国版本图书馆 CIP 数据核字(2008) 第 186179 号

责任编辑:范庆奎/责任印刷:钱玉芬/封面设计:黄华斌

科学出版社出版

北京东黄城根北街 16 号

邮政编码: 100717

<http://www.sciencep.com>

双青印刷厂印刷

科学出版社发行 各地新华书店经销

*

2009 年 1 月第 一 版 开本: B5(720×1000)

2009 年 1 月第一次印刷 印张: 17

印数: 1—2 500 字数: 320 000

定价: 60.00 元

(如有印装质量问题, 我社负责调换〈双青〉)

《国外数学名著系列》(影印版)专家委员会

(按姓氏笔画排序)

丁伟岳 王元 文兰 石钟慈 冯克勤 严加安
李邦河 李大潜 张伟平 张继平 杨乐 姜伯驹
郭雷

项目策划

向安全 林鹏 王春香 吕虹 范庆奎 王璐

执行编辑

范庆奎

《国外数学名著系列》(影印版)序

要使我国的数学事业更好地发展起来,需要数学家淡泊名利并付出更艰苦地努力。另一方面,我们也要从客观上为数学家创造更有利的发展数学事业的外部环境,这主要是加强对数学事业的支持与投资力度,使数学家有较好的工作与生活条件,其中也包括改善与加强数学的出版工作。

从出版方面来讲,除了较好较快地出版我们自己的成果外,引进国外的先进出版物无疑也是十分重要与必不可少的。从数学来说,施普林格(Springer)出版社至今仍然是世界上最具权威的出版社。科学出版社影印一批他们出版的好的新书,使我国广大数学家能以较低的价格购买,特别是在边远地区工作的数学家能普遍见到这些书,无疑是对推动我国数学的科研与教学十分有益的事。

这次科学出版社购买了版权,一次影印了 23 本施普林格出版社出版的数学书,就是一件好事,也是值得继续做下去的事情。大体上分一下,这 23 本书中,包括基础数学书 5 本,应用数学书 6 本与计算数学书 12 本,其中有些书也具有交叉性质。这些书都是很新的,2000 年以后出版的占绝大部分,共计 16 本,其余的也是 1990 年以后出版的。这些书可以使读者较快地了解数学某方面的前沿,例如基础数学中的数论、代数与拓扑三本,都是由该领域大数学家编著的“数学百科全书”的分册。对从事这方面研究的数学家了解该领域的前沿与全貌很有帮助。按照学科的特点,基础数学类的书以“经典”为主,应用和计算数学类的书以“前沿”为主。这些书的作者多数是国际知名的大数学家,例如《拓扑学》一书的作者诺维科夫是俄罗斯科学院的院士,曾获“菲尔兹奖”和“沃尔夫数学奖”。这些大数学家的著作无疑将会对我国的科研人员起到非常好的指导作用。

当然,23 本书只能涵盖数学的一部分,所以,这项工作还应该继续做下去。更进一步,有些读者面较广的好书还应该翻译成中文出版,使之有更大的读者群。

总之,我对科学出版社影印施普林格出版社的部分数学著作这一举措表示热烈的支持,并盼望这一工作取得更大的成绩。

王 元

2005 年 12 月 3 日

List of Editors, Authors and Translators

Editor-in-Chief

R. V. Gamkrelidze, Russian Academy of Sciences, Steklov Mathematical Institute, ul. Vavilova 42, 117966 Moscow, Institute for Scientific Information (VINITI), ul. Usievicha 20 a, 125219 Moscow, Russia

Consulting Editor

E. B. Vinberg, Department of Mathematics, Moscow University, 119899 Moscow, Russia

Authors

D. V. Alekseevskij, Sophus Lie Centre, Krasnokazarmennaya 6, 111250 Moscow, Russia

O. V. Shvartsman, Department of Applied Mathematics, Moscow Institute of Communications, Aviamotornaya ul. 8a, 105855 Moscow, Russia

A. S. Solodovnikov, State Financial Academy, 2nd Volkonskij per. 1, kv. 9, 103473 Moscow, Russia

E. B. Vinberg, Department of Mathematics, Moscow University, 119899 Moscow, Russia

Translator

V. Minachin, Cybernetics Council of the Russian Academy of Sciences, ul. Vavilova 40, 117966 Moscow, Russia

Foreword to the English Edition

The authors have introduced some improvements into the English edition. The literature list has been enlarged (mainly that to Part 2) and a number of short additions have been included in the text referring, in particular, to recent works which appeared after the publication of the Russian edition. We have also made every effort to correct misprints and occasional inaccuracies in the Russian version.

The process of translating the book into English has revealed that the use of geometric terminology in the English language literature on the subject is rather haphazard (which, incidentally, is also the case in Russian). In each case, after consulting many relevant English language publications, we and the translator have made our choice. However, in the footnotes we provide alternative terms also used in the literature.

The authors are extremely grateful to Dr. Minachin who performed the laborious work of translating the book with a sense of responsibility and interest.

E. B. Vinberg

Contents

I. Geometry of Spaces of Constant Curvature
D. V. Alekseevskij, E. B. Vinberg, A. S. Solodovnikov
1

**II. Discrete Groups of Motions of Spaces
of Constant Curvature**
E. B. Vinberg, O. V. Shvartsman
139

Author Index
249

Subject Index
251

I. Geometry of Spaces of Constant Curvature

D.V. Alekseevskij, E.B. Vinberg,
A.S. Solodovnikov

Translated from the Russian
by V. Minachin

Contents

Preface	6
Chapter 1. Basic Structures	8
§ 1. Definition of Spaces of Constant Curvature	8
1.1. Lie Groups of Transformations	8
1.2. Groups of Motions of a Riemannian Manifold	9
1.3. Invariant Riemannian Metrics on Homogeneous Spaces	10
1.4. Spaces of Constant Curvature	11
1.5. Three Spaces	11
1.6. Subspaces of the Space $\mathbb{R}^{n,1}$	14
§ 2. The Classification Theorem	15
2.1. Statement of the Theorem	15
2.2. Reduction to Lie Algebras	16
2.3. The Symmetry	16
2.4. Structure of the Tangent Algebra of the Group of Motions ..	17
2.5. Riemann Space	19
§ 3. Subspaces and Convexity	20
3.1. Involutions	20
3.2. Planes	21
3.3. Half-Spaces and Convex Sets	23
3.4. Orthogonal Planes	24

§ 4. Metric	26
4.1. General Properties	26
4.2. Formulae for Distance in the Vector Model	26
4.3. Convexity of Distance	27
Chapter 2. Models of Lobachevskij Space	29
§ 1. Projective Models	29
1.1. Homogeneous Domains	29
1.2. Projective Model of Lobachevskij Space	29
1.3. Projective Euclidean Models. The Klein Model	31
1.4. "Affine" Subgroup of the Group of Automorphisms of a Quadric	32
1.5. Riemannian Metric and Distance Between Points in the Projective Model	33
§ 2. Conformal Models	36
2.1. Conformal Space	36
2.2. Conformal Model of the Lobachevskij Space	39
2.3. Conformal Euclidean Models	42
2.4. Complex Structure of the Lobachevskij Plane	46
§ 3. Matrix Models of the Spaces \mathbb{H}^2 and \mathbb{H}^3	47
3.1. Matrix Model of the Space \mathbb{H}^2	47
3.2. Matrix Model of the Space \mathbb{H}^3	48
Chapter 3. Plane Geometry	50
§ 1. Lines	51
1.1. Divergent and Parallel Lines on the Lobachevskij Plane	51
1.2. Distance Between Points of Different Lines	54
§ 2. Polygons	56
2.1. Definitions. Conditions of Convexity	56
2.2. Elementary Properties of Triangles	57
2.3. Polar Triangles on the Sphere	59
2.4. The Sum of the Angles in a Triangle	59
2.5. Existence of a Convex Polygon with Given Angles	60
2.6. The Angular Excess and the Area of a Polygon	62
§ 3. Metric Relations	63
3.1. The Cosine Law for a Triangle	63
3.2. Other Relations in a Triangle	65
3.3. The Angle of Parallelism	67
3.4. Relations in Quadrilaterals, Pentagons, and Hexagons	68
3.5. The Circumference and the Area of a Disk	69
§ 4. Motions and Homogeneous Lines	70
4.1. Classification of Motions of Two-Dimensional Spaces of Constant Curvature	70

4.2. Characterization of Motions of the Lobachevskij Plane in the Poincaré Model in Terms of Traces of Matrices	71
4.3. One-Parameter Groups of Motions of the Lobachevskij Plane and Their Orbits	72
4.4. The Form of the Metric in Coordinates Related to a Pencil ..	75
Chapter 4. Planes, Spheres, Horospheres and Equidistant Surfaces ...	77
§ 1. Relative Position of Planes	77
1.1. Pairs of Subspaces of a Euclidean Vector Space	77
1.2. Some General Notions	79
1.3. Pairs of Planes on the Sphere	79
1.4. Pairs of Planes in the Euclidean Space	80
1.5. Pseudo-orthogonal Transformations	80
1.6. Pairs of Hyperbolic Subspaces of the Space $\mathbb{R}^{n,1}$	81
1.7. Pairs of Planes in the Lobachevskij Space	82
1.8. Pairs of Lines in the Lobachevskij Space	82
1.9. Pairs of Hyperplanes	83
§ 2. Standard Surfaces	84
2.1. Definitions and Basic Facts	84
2.2. Standard Hypersurfaces	86
2.3. Similarity of Standard Hypersurfaces	87
2.4. Intersection of Standard Hypersurfaces	89
§ 3. Decompositions into Semi-direct Products	90
3.1. Spherical, Horospherical, and Equidistant Decompositions ..	90
3.2. Spherical-equidistant Decomposition	90
Chapter 5. Motions	92
§ 1. General Properties of Motions	92
1.1. Description of Motions	92
1.2. Continuation of a Plane Motion	92
1.3. Displacement Function	93
§ 2. Classification of Motions	94
2.1. Motions of the Sphere	94
2.2. Motions of the Euclidean Space	95
2.3. Motions of the Lobachevskij Space	95
2.4. One-parameter Groups of Motions	97
§ 3. Groups of Motions and Similarities	98
3.1. Some Basic Notions	98
3.2. Criterion for the Existence of a Fixed Point	99
3.3. Groups of Motions of the Sphere	99
3.4. Groups of Motions of the Euclidean Space	100
3.5. Groups of Similarities	101
3.6. Groups of Motions of the Lobachevskij Space	101

Chapter 6. Acute-angled Polyhedra	104
§ 1. Basic Properties of Acute-angled Polyhedra	104
1.1. General Information on Convex Polyhedra	104
1.2. The Gram Matrix of a Convex Polyhedron	104
1.3. Acute-angled Families of Half-spaces and Acute-angled Polyhedra	106
1.4. Acute-angled Polyhedra on the Sphere and in Euclidean Space	107
1.5. Simplicity of Acute-angled Polyhedra	108
§ 2. Acute-angled Polyhedra in Lobachevskij Space	108
2.1. Description in Terms of Gram Matrices	108
2.2. Combinatorial Structure	109
2.3. Description in Terms of Dihedral Angles	110
Chapter 7. Volumes	112
§ 1. Volumes of Sectors and Wedges	113
1.1. Volumes of Sectors	113
1.2. Volume of a Hyperbolic Wedge	114
1.3. Volume of a Parabolic Wedge	116
1.4. Volume of an Elliptic Wedge	116
§ 2. Volumes of Polyhedra	117
2.1. Volume of a Simplex as an Analytic Function of the Dihedral Angles	117
2.2. Volume Differential	118
2.3. Volume of an Even-dimensional Polyhedron. The Poincaré and Schläfli Formulae	120
2.4. Volume of an Even-dimensional Polyhedron. The Gauss-Bonnet Formula	122
§ 3. Volumes of 3-dimensional Polyhedra	123
3.1. The Lobachevskij Function	123
3.2. Double-rectangular Tetrahedra	124
3.3. Volume of a Double-rectangular Tetrahedron. The Lobachevskij Formula	125
3.4. Volumes of Tetrahedra with Vertices at Infinity	126
3.5. Volume of a Pyramid with the Apex at Infinity	128
Chapter 8. Spaces of Constant Curvature as Riemannian Manifolds ..	130
1.1. Exponential Mapping	130
1.2. Parallel Translation	131
1.3. Curvature	131
1.4. Totally Geodesic Submanifolds	132

I. Geometry of Spaces of Constant Curvature	5
1.5. Hypersurfaces	133
1.6. Projective Properties	134
1.7. Conformal Properties	134
1.8. Pseudo-Riemannian Spaces of Constant Curvature	135
References	135

Preface

Spaces of constant curvature, i.e. Euclidean space, the sphere, and Lobachevskij space, occupy a special place in geometry. They are most accessible to our geometric intuition, making it possible to develop elementary geometry in a way very similar to that used to create the geometry we learned at school. However, since its basic notions can be interpreted in different ways, this geometry can be applied to objects other than the conventional physical space, the original source of our geometric intuition.

Euclidean geometry has for a long time been deeply rooted in the human mind. The same is true of spherical geometry, since a sphere can naturally be embedded into a Euclidean space. Lobachevskij geometry, which in the first fifty years after its discovery had been regarded only as a logically feasible by-product appearing in the investigation of the foundations of geometry, has even now, despite the fact that it has found its use in numerous applications, preserved a kind of exotic and even romantic element. This may probably be explained by the permanent cultural and historical impact which the proof of the independence of the Fifth Postulate had on human thought.

Nowadays modern research trends call for much more businesslike use of Lobachevskij geometry. The traditional way of introducing Lobachevskij geometry, based on a kind of Euclid-Hilbert axiomatics, is ill suited for this purpose because it does not enable one to introduce the necessary analytical tools from the very beginning. On the other hand, introducing Lobachevskij geometry starting with some specific model also leads to inconveniences since different problems require different models. The most reasonable approach should, in our view, start with an axiomatic definition, but it should be based on a well-advanced system of notions and make it possible either to refer to any model or do without any model at all.

Their name itself provides the description of the property by which spaces of constant curvature are singled out among Riemannian manifolds. However, another characteristic property is more important and natural for them — the property of maximum mobility. This is the property on which our exposition is based.

The reader should realize that our use of the term “space of constant curvature” does not quite coincide with the conventional one. Usually one understands it as describing any Riemannian manifold of constant curvature. Under our definition (see Chap. 1, Sect. 1) any space of constant curvature turns out to be one of the three spaces listed at the beginning of the Preface.

Although Euclidean space is, of course, included in our exposition as a special case, we have no intention of introducing the reader to Euclidean geometry. On the contrary, we make free use of its basic facts and theorems. We also assume that the reader is familiar with the basics of linear algebra and affine geometry, the notion of a smooth manifold and Lie group, and the elements of Riemannian geometry.

For the history of non-Euclidean geometry and the development of its ideas the reader is referred to relevant chapters in the books of Klein [1928], Kagan [1949, 1956], Coxeter [1957], and Efimov [1978].

Chapter 1

Basic Structures

§ 1. Definition of Spaces of Constant Curvature

This chapter provides the definition of spaces of constant curvature and of their basic structures, and describes their place among homogeneous spaces on the one hand and Riemannian manifolds on the other. If the reader's main aim is just to study Lobachevskij geometry, no great damage will be done if he skips Theorems 1.2 and 1.3 and the proof of Theorem 2.1.

1.1. Lie Groups of Transformations. We assume that the reader is familiar with the notions of a (real) smooth manifold and of a (real) Lie group. The word "smooth" (manifold, function, map etc.) always means that the corresponding structure is C^∞ . All smooth manifolds are assumed to have a countable base of open subsets. By $T_x(X)$ we denote the tangent space to a manifold X at a point x , and by $d_x g$ the differential of the map g at a point x . If no indication of the point is necessary the subscript is omitted.

We now recall some basic definitions of Lie group theory. (For more details see, e.g. Vinberg and Onishchik [1988].)

A group G of transformations¹ of a smooth manifold X endowed with a Lie group structure is said to be a *Lie group of transformations* of the manifold X if the map

$$G \times X \rightarrow X, \quad (g, x) \mapsto gx,$$

is smooth, which means that the (local) coordinates of the point gx are smooth functions of the coordinates of the element g and the point x . Then the stabilizer

$$G_x = \{g \in G : gx = x\}$$

of any point $x \in X$ is a (closed) Lie subgroup of the group G . Its linear representation $g \mapsto d_x g$ in the space $T_x(X)$ is called the *isotropy representation* and the linear group $d_x G_x$ is called the *isotropy group* at the point x .

The stabilizers of equivalent points x and $y = gx$ ($g \in G$) are conjugate in G , i.e.

$$G_y = gG_x g^{-1}.$$

The corresponding isotropy groups are related in the following way:

$$d_x G_y = (d_x g)(d_x G_x)(d_x g)^{-1}.$$

In other words, if tangent spaces $T_x(X)$ and $T_y(Y)$ are identified by the isomorphism $d_x g$, then the group $d_x G_x$ coincides with the group $d_y G_y$.

¹ By a group of transformations we understand an effective group of transformations, i.e. we assume that different transformations correspond to different elements of the group.

If G is a transitive Lie group of transformations of a manifold X , then for each point $x \in X$ the map

$$G/G_x \rightarrow X, \quad gG_x \mapsto gx$$

is a diffeomorphism commuting with the action of the group G . (The group G acts on the manifold G/G_x of left cosets by left shifts.) In this case the manifold X together with the action of G on it can be reconstructed from the pair (G, G_x) .

Definition 1.1. A smooth manifold X together with a given transitive Lie group G of its transformations is said to be a *homogeneous space*.

We denote a homogeneous space by (X, G) , or simply X .

A homogeneous space (X, G) is said to be connected or simply-connected² if the manifold X has this property.

1.2. Group of Motions of a Riemannian Manifold. A Riemannian metric is said to be defined on a smooth manifold X if a Euclidean metric is defined in each tangent space $T_x(X)$, and if the coefficients of this metric are smooth functions in the coordinates of x . A diffeomorphism g of a Riemannian manifold X is called a *motion* (or an *isometry*) if for each point $x \in X$ the linear map

$$d_x g : T_x(X) \rightarrow T_{gx}(X)$$

is an isometry. The set of all motions is evidently a group.

Each motion g takes a geodesic into a geodesic, and therefore commutes with the exponential map, i.e.

$$g(\exp \xi) = \exp dg(\xi)$$

for all $\xi \in T(X)$. Hence each motion g of a connected manifold X is uniquely defined by the image gx of some point $x \in X$ and the differential $d_x g$ at that point. This enables us to introduce coordinates into the group of motions, turning it into a Lie group. To be more precise, the following theorem holds.

Theorem 1.2 (Kobayashi and Nomizu [1981]). *The group of motions of a Riemannian manifold X is uniquely endowed with a differentiable structure, which turns it into a Lie group of transformations of the manifold X .*

If the group of motions of a Riemannian manifold X is transitive, then X is complete. Indeed, in this case there exists $\varepsilon > 0$, which does not depend on x , such that for any point $x \in X$ and for any direction at that point there exists a geodesic segment of length ε issuing from x in that direction. This implies that each geodesic can be continued indefinitely in any direction.

A Riemannian manifold X is said to have *constant curvature* c if at each point its sectional curvature along any plane section equals c .

² We assume that any simply-connected space is, by definition, connected.