


Series in Contemporary Applied Mathematics  
CAM 10

# Industrial and Applied Mathematics in China

中国的工业与应用数学

Ta-Tsien Li  
Pingwen Zhang  
*editors*



高等教育出版社  
Higher Education Press

Series in Contemporary Applied Mathematics CAM 10

# Industrial and Applied Mathematics in China

中国的工业与应用数学



Higher Education Press



World Scientific

NEW JERSEY • LONDON • SINGAPORE • BEIJING • SHANGHAI • HONG KONG • TAIPEI • CHENNAI

**Ta-Tsien Li**

School of Mathematical Sciences  
Fudan University  
220, Handan Road  
Shanghai, 200433  
China

**Pingwen Zhang**

School of Mathematical Sciences  
Peiking University  
5, Yiheyuan Road  
Beijing, 100871  
China

**Editorial Assistant:** Zhou Chun-Lian

## 图书在版编目 (CIP) 数据

中国的工业与应用数学 = Industrial and Applied  
Mathematics in China: 英文 / 李大潜, 张平文主编.  
北京: 高等教育出版社, 2008.12  
(现代应用数学丛书)  
ISBN 978-7-04-023722-1

I. 中... II. ①李...②张... III. ①工程数学—英文  
②应用数学—英文 IV. TB11

中国版本图书馆 CIP 数据核字 (2008) 第 193153 号

Copyright © 2009 by

**Higher Education Press**

4 Dewai Dajie, Beijing 100011, P. R. China, and

**World Scientific Publishing Co Pte Ltd**

5 Toh Tuch Link, Singapore 596224

*All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without permission in writing from the Publisher.*

**ISBN 978-7-04-023722-1**

Printed in P. R. China

## Preface

The China Society for Industrial and Applied Mathematics (CSIAM) had its 9th Annual Conference entitled “Industrial and Applied Mathematics in China” with 12 plenary talks from August 14 to 18, 2006 in Nanjing, China. Later on, in the 6th International Congress on Industrial and Applied Mathematics (ICIAM 2007) held from July 16 to 20, 2007 in Zurich, Switzerland, CSIAM organized an embedded meeting with the same title on July 18, 2007, which consists of two two-hour sessions with six lectures. Since all these talks concern the topic “Industrial and Applied Mathematics in China”, we gather a large part of them in this volume for publication. We hope that the readers can get an impression on the present situation and trends of the industrial and applied mathematics in China from this volume and the researchers and graduate students in applied mathematics and in applied sciences can benefit from the mathematical models and methods with applications presented in this book.

We would like to take this opportunity to give our sincere thanks to all the speakers and, in particular, to those who gave their contribution to this volume for their kind help and support.

Ta-Tsien Li  
July 2008

# Contents

## Preface

<i>Xiaoshan Gao, Ziming Li</i> : Mechanized Methods for Differential and Difference Equations .....	1
<i>Song Jiang, Feng Xie, Jianwen Zhang</i> : A Global Existence Result in Radiation Hydrodynamics .....	25
<i>Shi Jin</i> : Recent Computational Methods for High Frequency Waves in Heterogeneous Media .....	49
<i>Ying Bao, Zhiming Ma, Yanhong Shang</i> : Some Recent Results on Ranking Webpages and Websites .....	65
<i>Lifeng Chen, Shige Peng</i> : Report on Testing and Finding the Generating Functions $g$ of an Option Pricing Mechanism through Market Data .....	79
<i>Jun Hu, Zhongci Shi</i> : Analysis of Nonconforming Rotated $Q_1$ Element for the Reissner-Mindlin Plate Problem .....	101
<i>Yongji Tan</i> : Monitoring the Corrosion of the Blast Furnace by Perturbation Method .....	112
<i>Yong Xiao, Sufen Zhao, Xiaoping Wang</i> : Numerical Study of Magnetic Properties of Nanowire Arrays .....	129
<i>Zongmin Wu</i> : Generalized B-spline .....	141
<i>Xuan Zeng, Hengliang Zhu, Fan Yang, Jun Tao, Yi Wang, Jintao Xue</i> : Mathematical Problems in System-on-Chip Design and Manufacture .....	166

*Weiwei Qi, Ming Chen, Huitao Zhang, Peng Zhang:*

A New Reconstruction Algorithm for Cone-beam CT with Unilateral  
Off-centered RT Multi-scan ..... 215

*Tie Zhou, Jiantao Cheng, Ming Jiang:*

Bioluminescence Tomography Reconstruction by Radial Basis  
Function Collocation Method ..... 229

# Mechanized Methods for Differential and Difference Equations\*

Xiaoshan Gao, Ziming Li

*Key Laboratory of Mathematics Mechanization*

*Institute of Systems Science, AMSS*

*Academia Sinica, Beijing 100080, China*

*Email: {xgao, zml}@mmrc.iss.ac.cn*

## Abstract

Some recent results on the mechanized methods for differential and difference equations are surveyed. The results include: the characteristic set method for differential and difference equation systems, algorithms for computing closed-form solutions of differential and difference equations, and algorithms for solving and factoring finite-dimensional linear functional systems.

## 1 Introduction

This paper provides a survey of some recent work on differential and difference equations by researchers at the Key Laboratory of Mathematics Mechanization and their collaborators. The work under review is greatly stimulated by Wu's method for mechanical theorem-proving in differential geometries, finding closed-form solutions of differential (difference) equations, and handling analytic and discrete mathematical objects by computers.

Differential equations describe physical laws in mechanics and geometric properties of manifolds. The characteristic set method for differential equations enables us to search for physical laws and geometric properties by computers [52]. For example, Newton's gravitational law is automatically derived from Kepler's laws [51], and "Theorema Egregium" is rediscovered by computing a characteristic set of the fundamental equations of surface theory [30].

The notion of characteristic sets for differential ideals was introduced by Ritt [42]. It plays a fundamental role in differential algebra, because

---

\*Partially supported by a National Key Basic Research Project of China (2004CB318000).

the Hilbert Basissatz does not hold for differential ideals. The notion and algorithm of characteristic sets for polynomial and differential polynomial sets were introduced by Wu [48,50] to prove theorems in geometries and to manipulate systems of differential and algebraic equations [48,49]. Wu's work inspired a great deal of research in the communities of symbolic computation and automated reasoning. Later on, the success of Gröbner bases for polynomial ideals led to methods to characterizing radical differential ideals [4]. The reader may consult [44] for more details on the recent developments of the differential characteristic set method. In this paper we briefly review Wu's scheme for differential characteristic sets and point out its recent extension to difference polynomial systems.

Integrals, special functions and combinatorial sequences are often considered as "infinite" objects. To specify them in terms of a finite amount of information on computers, one uses the differential (difference) equations annihilating these objects. For instances, automatic proofs of combinatorial identities need to find hypergeometric solutions of difference equations [37], while algorithms for symbolic integration need to compute elementary functions satisfying Risch's equation [7]. Great efforts have been made to compute closed-form solutions of linear ordinary differential (difference) equations (see, [26,39] and the references therein). There are two ways to go further: one is to look for closed-form solutions of nonlinear ordinary differential (difference) equations of some kind; the other is to develop symbolic algorithms for linear partial differential (difference) equations. We will summarize recent theoretical and algorithmic results concerning this subject.

Nonlinear differential equations arise from physics. Their analytic solutions are important for the understanding of the physical phenomena. Interesting methods to search for analytic solutions of nonlinear PDEs are given in [16,53].

Factoring polynomials helps us to solve algebraic equations. Likewise, we want to decompose differential and difference equations into those of lower orders. There have been efficient algorithms for decomposing linear ordinary differential operators [6,24,25,43]. Recent work on extending these methods to linear partial differential and difference equations [33] will be surveyed. We also mention that a decomposition algorithm for nonlinear ordinary differential equations is presented in [23].

The rest of this paper is organized as follows. In Section 2, we outline the differential characteristic method. Methods for computing rational and algebraic solutions of first-order ordinary differential and difference equations are presented in Section 3. An algebraic setting and a factorization algorithm for finite-dimensional linear functional systems are described in Sections 4 and 5, respectively.



## 2 The characteristic set method

The characteristic set method plays a central role in the theory and applications of mathematics mechanization. In this section, we will introduce its main features and applications in automated reasoning.

### 2.1 Properties of ascending chains

Let  $\mathbb{K}$  be an ordinary differential field,  $\mathbb{X} = \{x_1, \dots, x_n\}$  a set of differential indeterminates, and  $\mathbb{K}\{\mathbb{X}\}$  the set of *differential polynomials* in  $\mathbb{X}$  with coefficients in  $\mathbb{K}$ . We denote  $x_{i,j}$  to be the  $j$ -derivative of  $x_i$ . The *universal field*  $\mathbb{E}$  over  $\mathbb{K}$  is a differentially closed field containing  $\mathbb{K}$  and infinitely many indeterminates. For a polynomial  $D$  and a polynomial set  $\mathbb{P} \subset \mathbb{K}[\mathbb{X}]$ ,

$$\text{Zero}(\mathbb{P}) = \{\eta \in \mathbb{E}^n \mid P(\eta) = 0, \forall P \in \mathbb{P}\}$$

is called a *variety*, and  $\text{Zero}(\mathbb{P}/D) = \text{Zero}(\mathbb{P}) \setminus \text{Zero}(D)$  is called a *quasi variety*.

A set  $\mathcal{A}$  of differential polynomials is called an *ascending chain* (triangular set), or simply a chain, if after renaming the indeterminates in  $\mathbb{X}$  as  $\mathbb{U} = \{u_1, \dots, u_q\}$  and  $\mathbb{Y} = \{y_1, \dots, y_p\}$ , we can write  $\mathcal{A}$  in the following form:

$$\begin{aligned} A_1(\mathbb{U}, y_1) &= I_1 y_{1,o_1}^{d_1} + \text{terms of lower orders and degrees in } y_1, \\ &\dots \\ A_p(\mathbb{U}, y_1, \dots, y_p) &= I_p y_{p,o_p}^{d_p} + \text{terms of lower orders and degrees in } y_p. \end{aligned} \tag{1}$$

As a matter of terminologies,  $o_i$  is called the order of  $A_i$ ;  $I_i$  is called the *initial* of  $A_i$ ,  $S_i = \frac{\partial A_i}{\partial y_{i,o_i}}$  is called the *separant* of  $A_i$ . Write  $\mathbf{I}_{\mathcal{A}} = \prod_i I_i S_i$ . The *dimension* of  $\mathcal{A}$  is defined to be  $|\mathbb{U}| = q$ , which is denoted  $\dim(\mathcal{A})$ . The *order* of  $\mathcal{A}$  is defined to be  $\text{ord}(\mathcal{A}) = \sum_{i=1}^p o_i$ . The *degree* of  $\mathcal{A}$  is defined to be  $\deg(\mathcal{A}) = \prod_{i=1}^p d_i$ .

We could say that the formal solutions for a chain is basically determined. Intuitively, for a set of given values of the parameters  $\mathbb{U}$ , the  $y_i$  can be determined iteratively by solving univariate equations  $A_i = 0$ . In order to show the properties of chains, we first introduce several concepts. The *saturation ideal* of  $\mathcal{A}$  is defined to be

$$\text{sat}(\mathcal{A}) = \{P \in \mathbb{K}\{\mathbb{X}\} \mid \exists k \in \mathbb{N}, \mathbf{I}_{\mathcal{A}}^k P \in (\mathcal{A})\}.$$

We may define a partial ordering among the chains in a nature way [42, 52]. It is known that any set of chains contains one with lowest order. A *characteristic set* of a differential polynomial set  $\mathbb{P}$  is any chain of lowest ordering contained in  $\mathbb{P}$ .

A chain  $\mathcal{A}$  is called *irreducible* if  $A_1$  is an irreducible polynomial in  $y_{1,o_1}$  and  $A_k$  is an irreducible polynomial modulo  $A_1, \dots, A_{k-1}$ .

**Theorem 2.1.** [42, 52] *Let  $\mathcal{A}$  be an irreducible chain. Then  $\text{sat}(\mathcal{A})$  is a prime ideal of dimension  $\dim(\mathcal{A})$ , order  $\text{ord}(\mathcal{A})$  wrt  $\mathbb{U}$ , and degree  $\deg(\mathcal{A})$  wrt  $\mathbb{U}$ . Conversely, a characteristic set of a prime ideal is irreducible.*

The following result shows that the dimension, order and degree of a chain are intrinsic properties.

**Theorem 2.2.** [19, 22] *Let  $\mathcal{A}$  be a chain of form (1). If  $\text{Zero}(\text{sat}(\mathcal{A})) \neq \emptyset$ ,  $\text{Zero}(\text{sat}(\mathcal{A}))$  and  $\text{Zero}(\mathcal{A}/\mathbb{I}_{\mathcal{A}})$  are unmixed. More precisely, write  $\text{Zero}(\text{sat}(\mathcal{A}))$  as an irredundant decomposition:  $\text{Zero}(\text{sat}(\mathcal{A})) = \cup_{i=1}^r \text{Zero}(\text{sat}(\mathcal{C}_i))$ . Then*

- (1)  $\mathcal{C}_i$  is also of form (1). As a consequence,  $\dim(\text{sat}(\mathcal{C}_i)) = \dim(\mathcal{A})$  and  $\text{ord}(\mathcal{C}_i) = \text{ord}(\mathcal{A})$ .
- (2)  $\deg(\mathcal{A}) \geq \sum_{i=1}^r \deg(\mathcal{C}_i)$ . Furthermore,  $\deg(\mathcal{A}) = \sum_{i=1}^r \deg(\mathcal{C}_i)$  iff  $\mathcal{A}$  is saturated, that is, the initials and seprants of  $\mathcal{A}$  are invertible wrt  $\mathcal{A}$ .

Another important property for chains is

**Theorem 2.3.** [52] *An irreducible chain admits a formal power series solution which can be computed algorithmically.*

In order to make the paper shorter, we limit to the ordinary differential case. Similar results for the partial differential case were also established, where we need to assume that the chains are either passive [49, 52] or coherent [4, 5, 27].

Similar results are also proved in the case of algebraic difference polynomials [21, 22]. However, in the difference case, we do not have algorithms to decide whether a chain is irreducible. In order to have a constructive theory, proper irreducible chains are introduced [21]. Also, Theorem 2.2 is proved only for proper irreducible chains.

## 2.2 Characteristic set method

The characteristic set method decomposes the zero set for a differential polynomial system in general form into the union of zero sets for chains. Since the zero set of a chain is considered to be known, this method gives a general tool to deal with differential equation systems.

Let  $\mathbb{P}$  be a finite set of differential polynomials. Then we can perform the following operations:

$$\begin{aligned} \mathbb{P} &= \mathbb{P}_0 \ \mathbb{P}_1 \ \cdots \ \mathbb{P}_i \ \cdots \ \mathbb{P}_m, \\ \mathcal{B}_0 \ \mathcal{B}_1 \ \cdots \ \mathcal{B}_i \ \cdots \ \mathcal{B}_m &= \mathcal{C}, \\ \mathbb{R}_0 \ \mathbb{R}_1 \ \cdots \ \mathbb{R}_i \ \cdots \ \mathbb{R}_m &= \emptyset, \end{aligned} \quad (2)$$

where  $\mathcal{B}_i$  is a lowest chain in  $\mathbb{P}_i$  with respect to a pre-selected partial ordering;  $\mathbb{R}_i$  is the set of nonzero remainders of the polynomials in  $\mathbb{P}_i$  wrt  $\mathcal{B}_i$ ; and  $\mathbb{P}_{i+1} = \mathbb{P}_0 \cup \mathcal{B}_i \cup \mathbb{R}_i$ . In scheme (2),  $\mathcal{B}_m = \mathcal{C}$  verifies

$$\text{prem}(\mathbb{P}, \mathcal{C}) = \{0\} \text{ and } \text{Zero}(\mathbb{P}) \subset \text{Zero}(\mathcal{C}), \quad (3)$$

where  $\text{prem}$  denotes the differential pseudo-remainder. Any chain  $\mathcal{C}$  verifying the property (3) is called a *Wu characteristic set* of  $\mathbb{P}$ .

**Theorem 2.4** (Wu's Well-ordering Principle). [49, 52] *Let  $\mathcal{C}$  be a Wu characteristic set of a finite set  $\mathbb{P}$  of differential polynomials. Then:*

$$\begin{aligned} \text{Zero}(\mathbb{P}) &= \text{Zero}(\mathcal{C}/\mathbb{I}_{\mathcal{C}}) \bigcup \bigcup_i \text{Zero}(\mathbb{P} \cup \mathcal{C} \cup \{I_i\}), \\ \text{Zero}(\mathbb{P}) &= \text{Zero}(\text{sat}(\mathcal{C})) \bigcup \bigcup_i \text{Zero}(\mathbb{P} \cup \mathcal{C} \cup \{I_i\}), \end{aligned}$$

where  $I_i$  are the initials and separants of the polynomials in  $\mathcal{C}$ .

Using the well-ordering principle recursively, we obtain the following key result.

**Theorem 2.5** (Ritt-Wu's Zero Decomposition Theorem). [42, 52] *There is an algorithm which permits to determine, for a given finite set  $\mathbb{P}$  of differential polynomials, a finite set of (irreducible) chains  $\mathcal{A}_j$  such that*

$$\text{Zero}(\mathbb{P}) = \bigcup_j \text{Zero}(\mathcal{A}_j/\mathbb{I}_{\mathcal{A}_j}) = \bigcup_j \text{Zero}(\text{sat}(\mathcal{A}_j)).$$

Let  $\mathbb{P}$  be a finite subset of  $\mathbb{K}\{\mathbb{U}, \mathbb{X}\}$ , and  $D \in \mathbb{K}\{\mathbb{U}, \mathbb{X}\}$ , where  $\mathbb{U} = \{u_1, \dots, u_m\}$  and  $\mathbb{X} = \{x_1, \dots, x_n\}$ . The projection of  $\text{Zero}(\mathbb{P}/D)$  to  $\mathbb{U}$  is defined as follows:

$$\text{Proj}_{\mathbb{X}} \text{Zero}(\mathbb{P}/D) = \{e \in \mathbb{E}^m \mid \exists a \in \mathbb{E}^n \text{ s.t. } (e, a) \in \text{Zero}(\mathbb{P}/D)\}.$$

Projection for quasi-varieties can be computed with the characteristic set method.

**Theorem 2.6** (Projection Theorem). [19] *For a finite subset set  $\mathbb{P} \subset K\{\mathbb{U}, \mathbb{X}\}$  and  $D \in K\{\mathbb{U}, \mathbb{X}\}$ , we can compute chains  $\mathcal{A}_i$  and polynomials  $D_i$  in  $\mathbb{K}[\mathbb{U}]$  such that*

$$\text{Proj}_{\mathbb{X}} \text{Zero}(\mathbb{P}/D) = \bigcup_{i=1}^l \text{Zero}(\mathcal{A}_i/D_i \mathbb{I}_{\mathcal{A}_i}).$$

The concept of characteristic sets for prime ideals was introduced by Ritt [42]. The notion of characteristic sets given above, the well-ordering principle, and the current form of zero decomposition theorems were introduced by Wu [48,49,52]. An implementation of the method can be found in [46]. In order to improve the efficiency, new characteristic set methods were proposed [4,5,9,10,18,27,40,45]. The characteristic set method was used to solve certain problems for analytical functions [41].

A characteristic set method for algebraic difference equation systems was proposed in [21,22]. It is quit surprising that there are no essential progresses for the theory and algorithms of difference characteristic set methods since the early work of Ritt and his colleagues in the 1930s. In [21], an algorithm was proposed to decompose the zero set a difference polynomial system into the union of unmixed zero sets of difference polynomial systems represented by proper irreducible chains. In [22], a new resolvent theory for difference polynomial systems was proposed.

To solve a set of equations in triangular form, we need to solve univariate equations in a cascade form. The resolvent methods were introduced to reduce the solving of equation systems into the solving of one univariate equation plus a set of linear equations [13,22].

## 2.3 Wu's method of automated geometry theorem proving and discovering

A geometry theorem is called a *theorem of equality type*, if after introducing coordinates, the theorem can be expressed in the following form

$$\forall x_i[(H_1 = 0 \wedge \cdots \wedge H_s = 0 \wedge D_1 \neq 0 \wedge \cdots \wedge D_t \neq 0) \implies (C = 0)], \quad (4)$$

where  $H_i, D_i, C$  are in  $\mathbb{K}\{\mathbb{X}\}$ .

For theorems of equality type, we have the following principles of mechanical theorem proving, which are consequences of Theorems 2.1 and 2.4.

**Theorem 2.7.** [49] *For a geometry statement of form (4), let  $\mathcal{A}$  be a Wu-characteristic set of  $\{H_1, \dots, H_s\}$ . If  $\text{prem}(C, \mathcal{A}) = 0$ , then the statement is valid under the non-degenerate condition  $\mathbf{I}_{\mathcal{A}} \neq 0$ .*

Note that the non-degenerate condition  $\mathbf{I}_{\mathcal{A}} \neq 0$  is generated automatically by the algorithm.

**Theorem 2.8.** [52] *Let  $D = \prod_i D_i$ . By Theorem 2.5, we have*

$$\text{Zero}(\{H_1, \dots, H_s\}/D) = \cup_{i=1}^l \text{Zero}(\text{sat}(\mathcal{A}_i)/D).$$

*If  $\text{prem}(C, \mathcal{A}_i) = 0, i = 1, \dots, l$ , then the statement is true. If  $\mathcal{A}_i$  is irreducible and  $\text{prem}(C, \mathcal{A}_i) \neq 0$ , then the statement is not valid on  $\text{Zero}(\text{sat}(\mathcal{A}_i)/D)$ .*

As an example, let us show how to prove Newton's gravitational law with Kepler's laws. The first and second Kepler laws state that each planet describes an ellipse with the sun in one focus and the radius vector drawn from the sun to a planet sweeps out equal areas in equal times. The Newton's law states that the acceleration is reversely proportional to the distance from the planet to the sun. We may use differential equations  $K_1 = 0$ ,  $K_2 = 0$ , and  $N_1 = (ar^2)' = 0$  to represent these laws:

$$\begin{aligned} h_1 &= r^2 - x^2 - y^2 = 0, \\ h_2 &= a^2 - x''^2 - y''^2 = 0, \\ K_1 &= r - p - ex = 0 \wedge p' = 0 \wedge e' = 0, \\ K_2 &= y'x - yx' - h = 0 \wedge h' = 0, \\ d_1 &= p \neq 0 \quad (\text{The ellipse is not a line.}). \end{aligned}$$

Then, we need to show

$$\forall x, y, p, e, a, r[(K_1 = 0 \wedge K_2 = 0 \wedge h_1 = 0 \wedge h_2 = 0 \wedge d_1 \neq 0) \Rightarrow N_1 = 0].$$

By Theorem 2.5 ( $p < e < x < y < r < a$ ),

$$\text{Zero}(\{K_1, p', e', h_1, h_2, n_2\}/p) = \text{Zero}(\text{sat}(\mathcal{A}_1)p),$$

where  $\mathcal{A}_1$  is a chain. By computation, we have  $\text{prem}(n_1, ASC_1) = 0$ , which proves Newton's law.

There are two kinds of problems in differential geometry other than theorem proving. One is finding locus equations, the other is deriving geometry formulas. For a geometric configuration given by a set of polynomial equations  $h_1(\mathbb{U}, x_1, \dots, x_p) = 0, \dots, h_r(\mathbb{U}, x_1, \dots, x_p) = 0$ , we want to find a relation between arbitrarily chosen variables  $\mathbb{U}$  (parameters) and a dependent variable, say,  $x_1$ . Wu pointed out that the characteristic set method can be used to discover such unknown geometric formulas [51]. Actually, Newton's law can be deduced from Kepler's laws automatically in this way. More detailed accounts can be found in [10, 11, 30, 45].

The characteristic set method can be used to prove a much wider class of geometry theorems. Let  $\mathbb{E}$  be a differentially closed extension of  $\mathbb{K}$ , say, the field of meromorphic functions [42]. A *first order formula* over  $\mathbb{E}$  can be defined as follows.

1. If  $P \in \mathbb{K}[\mathbb{X}]$ , then  $P(\mathbb{X}) = 0$  is a formula.
2. If  $f, g$  are formulas, then  $\neg f$ ,  $f \wedge g$ , and  $f \vee g$  are formulas.
3. If  $f$  is a formula, then  $\exists x_i \in \mathbb{E}(f)$  and  $\forall x_i \in \mathbb{E}(f)$  are formulas.

A formula can always be written as a prefix canonical form

$$\phi = Q_1 y_1 \dots Q_m y_m \psi(u_1, \dots, u_d, y_1, \dots, y_m), \quad (5)$$

where  $Q_k$  is a quantifier  $\exists$  or  $\forall$  and  $\psi$  a formula free of quantifiers. For a first order formula  $\phi$  of form (5), there exists a fundamental problem:

**Quantifier Elimination:** Find a formula  $\theta(u_1, \dots, u_d)$  such that  $\theta$  is equivalent to  $\phi$ . If  $d = 0$ , we need to decide whether  $\phi$  is valid or not.

As a consequence of Theorem 2.6, we have

**Theorem 2.9.** *There exists a decision procedure for the first order theory over a differentially closed field.*

### 3 Rational and algebraic solutions of ODEs and OΔEs

For brevity we abbreviate ordinary difference equations as OΔE.

By decomposing the zero set of a differential polynomial system into the zero sets of chains, the characteristic set method gives a complete way to describe the structure for the zero sets of equation systems. In particular, finding the solutions of differential polynomial systems can be reduced to finding those of a single differential equation or a system of equations in a single variable.

Closed-form solutions of linear ODEs and OΔEs were widely studied. On the other hand, similar results to nonlinear ODEs are very limited. In this section, we summarize some recent results on finding rational and algebraic solutions to nonlinear ODEs and OΔEs. It is interesting to see whether these results can be treated uniformly with the differential Galois theory [35].

#### 3.1 Rational and algebraic solutions of algebraic ODEs

Let  $P \in \mathbb{K}\{y\} \setminus \mathbb{K}$  be an irreducible differential polynomial in an indeterminate  $y$  and

$$\Sigma_P = \{A \in \mathbb{K}\{y\} \mid SA \equiv 0 \pmod{\{P\}}\},$$

where  $S$  is the separant of  $P$  and  $\{P\}$  is the radical differential ideal generated by  $P$ . Then  $\Sigma_P$  is a prime ideal [42]. A generic zero of  $\Sigma_P$  is defined to be a *general solution* of  $P = 0$ . In particular, an *algebraic general solution* of  $P = 0$  is a general solution  $\hat{y}$  which satisfies the following equation

$$G(x, y) = \sum_{i=0}^n a_i(x)y^i = 0, \tag{6}$$

where  $a_i$  is a polynomial in  $x$  with degree  $\alpha_i$  and with constant coefficients, and  $G(x, y)$  is an irreducible polynomial in  $x, y$ . When  $n = 1$ ,  $\hat{y}$  is called a *rational general solution* of  $P = 0$ .

For  $\alpha_0, \alpha_1, \dots, \alpha_n \in \mathbb{Z}_{\geq 0}$ , we define the differential polynomial

$$\mathbb{D}_{(\alpha_0; \alpha_1, \dots, \alpha_n)} := \det(\mathcal{A}_{(h, \alpha_1; \alpha_0)}(y) | \mathcal{A}_{(h, \alpha_2; \alpha_0)}(y^2) | \dots | \mathcal{A}_{(h, \alpha_n; \alpha_0)}(y^n)),$$

where

$$\mathcal{A}_{(h, \alpha; k)}(y) := \begin{pmatrix} \binom{k+1}{0} y_{k+1} & \binom{k+1}{1} y_k & \cdots & \binom{k+1}{\alpha} y_{k+1-\alpha} \\ \binom{k+2}{0} y_{k+2} & \binom{k+2}{1} y_{k+1} & \cdots & \binom{k+2}{\alpha} y_{k+2-\alpha} \\ \vdots & \vdots & \cdots & \vdots \\ \binom{k+h+1}{0} y_{k+h+1} & \binom{k+h+1}{1} y_{k+h} & \cdots & \binom{k+h+1}{\alpha} y_{k+h+1-\alpha} \end{pmatrix}.$$

We have

**Lemma 3.1.** [1]  $y(x)$  satisfies an equation of the type (6) if and only if  $\mathbb{D}_{(\alpha_0; \alpha_1, \dots, \alpha_n)}(y(x)) = 0$ . As a consequence, we give a defining differential equation for algebraic functions.

When  $n = 2$ ,  $\alpha_1 = \alpha_2 = 1$  and  $\alpha_0 = 2$ ,

$$\mathbb{D}_{(2; 1, 1)} = \begin{vmatrix} y_3 & 3y_2 & (y^2)''' & 3(y^2)'' \\ y_4 & 4y_3 & (y^2)^{(4)} & 4(y^2)''' \\ y_5 & 5y_4 & (y^2)^{(5)} & 5(y^2)^{(4)} \\ y_6 & 6y_5 & (y^2)^{(6)} & 6(y^2)^{(5)} \end{vmatrix}.$$

We have  $\mathbb{D}_{(2; 1, 1)}(y(x)) = 0$  if and only if

$$(a_{2,1}x + a_{2,0})y^2(x) + (a_{1,1}x + a_{1,0})y(x) + a_{0,2}x^2 + a_{0,1}x + a_{0,0} = 0$$

for constants  $a_{i,j}$ .

The key to find a rational and algebraic function solutions is to give a degree bound for the solution. We can give these degree bounds for first order autonomous ODEs. In what follows, let  $F(y, y_1) = 0$  be a first order autonomous ODE. Then we have

**Theorem 3.2.** [1] If  $G(x, y) = 0$  defines a nontrivial algebraic solution of  $F = 0$ , then

- (1)  $\deg(G(x, y), x) = \deg(F, y_1)$ ,
- (2)  $\deg(G(x, y), y) \leq \deg(F, y) + \deg(F, y_1)$ .

The following example shows that the bound in (2) is optimal. Let  $n > m > 0$  and  $(n, m) = 1$ . Then  $G = y^n - x^m$  is irreducible.  $y^n - x^m = 0$  is an algebraic solution of  $F = y^{n-m}y_1^m - (m/n)^m = 0$ . Here,  $\deg(G(x, y), y) = \deg(F, y) + \deg(F, y_1)$ .

For rational solutions, we could give the exact degree bound [17].

**Theorem 3.3.** *If  $y = P(x)/Q(x)$  is a rational solution of  $F(y, y_1) = 0$ , then  $\deg(y(x)) = \deg(F, y_1)$ .*

These degree bounds are obtained by treating  $F(y, y_1) = 0$  as an algebraic curve and the solution as a parametrization of the curve. This idea also leads to the following algorithm to find a rational solution to a first order autonomous ODE [17].

**Theorem 3.4.** *Let  $y = r(x)$ ,  $y_1 = s(x)$  be a proper rational parametrization of  $F(y, y_1) = 0$ , where  $r(x), s(x)$  are rational functions in  $x$  with constant coefficients. Then  $F = 0$  has a rational general solution iff we have the following relations*

$$ar(x)' = s(x) \quad \text{or} \quad a(x - b)^2 r(x)' = s(x),$$

where  $a, b$  are constants and  $a \neq 0$ . If one of the above relations is true, then replacing  $x$  by  $a(x + c)$  (or  $b - \frac{1}{a(x+c)}$ ) in  $y = r(x)$ , we obtain a rational general solution of  $F = 0$ , where  $c$  is an arbitrary constant.

The above algorithm depends on the rational parametrization of algebraic curves. A more efficient algorithm is based on Hermite-Padé approximation.

Let  $A(x)$  be a formal power series. If a polynomial  $G(x, y)$  satisfies

$$G(x, A(x)) = O\left(x^{(n+1)(m+1)+1}\right),$$

where  $m = \deg(G, x)$ ,  $n = \deg(G, y)$ , then we call  $G(x, y) = 0$  Hermite-Padé approximant to  $A(x)$ . We find the algebraic solution for an first order autonomous ODEs as follows [1].

- (1) Find the first  $N$  terms  $f(x)$  of formal power series solution of  $F(y, y_1) = 0$ , where

$$N = 2(\deg(F, y) + \deg(F, y_1)).$$

- (2) Let  $d = \deg(F, y_1)$ . Construct the  $(d, d, \dots, d)$  Hermite-Padé approximant  $G(x, y) = 0$  to  $f(x)$ .
- (3) We need only to check whether  $G = 0$  is a nontrivial algebraic solution of  $F = 0$ .

The complexity of this algorithm is polynomial in terms of the number of the multiplications in the number field.

### 3.2 Rational solutions of algebraic OΔEs

The result about rational solutions of ODEs can be extended to OΔEs. Let  $\mathbb{K} = \mathbb{Q}(x)$  be the difference field with the difference operator  $\mathbf{E}(x) = x + 1$ ,  $y$  an indeterminate, and  $y_n = \mathbf{E}^n y$ .



Let  $P \in \mathbb{K}\{y\} \setminus \mathbb{K}$  be an irreducible difference polynomial in  $y$ , and

$$\Sigma_P = \{A \in \mathbb{Q}(x)\{y\} \mid SA \equiv 0 \pmod{\{P\}}\},$$

where  $S$  is the separant of  $P$ . Cohn proved that  $\Sigma_P$  is a perfect difference ideal and it could be decomposed into the intersection of the principle components of  $P$  [14]. Let  $\Lambda$  be one of the principle components of  $P$ . A general solution of  $P = 0$  is defined as a generic zero of one of the principle components of  $\Sigma_P$ . A *rational general solution* of  $P(y) = 0$  is defined as a general solution of  $P = 0$  with the following form:

$$\hat{y}(x) = \frac{a_n x^n + a_{n-1} x^{n-1} + \dots + a_0}{x^m + b_{m-1} x^{m-1} + \dots + b_0}, \quad (7)$$

where  $a_i, b_j$  are constants. In particular, if  $m = 0$ , we call  $\hat{y}(x)$  polynomial general solution. For instance, the difference equation  $(y - y_1)^2 - 2(y + y_1) + 1 = 0$  has two general solutions:  $y(x) = (x + c)^2$  and  $y(x) = (ce^{i\pi x} + \frac{1}{2})^2$  where  $c$  is an arbitrary constant.

The defining difference equations for polynomial and rational functions are given by the following lemmas [20].

**Lemma 3.5.** Let  $\mathcal{P}_n = \sum_{i=0}^{n+1} (-1)^i \binom{n+1}{i} y_i$ . Then  $y(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$  ( $\mathbf{E}(a_i) = a_i$ ) if and only if  $\mathcal{P}_n(y(x)) = 0$ .

Let

$$\mathcal{R}_{n,m} = \det \left( \sum_{i=0}^{n+1} (-1)^i \binom{n+1}{i} Y_i * M_i \right),$$

where  $Y_i = \text{diag}(y_i, y_{i+1}, \dots, y_{m+i})$ ,  $M_i = (H_{k,l}(i))_{(m+1) \times (m+1)}$ ,

$$\begin{aligned} & H_{k,l}(i) \\ &= \frac{(i+k-n)(i+k-n-1) \cdots (i+k-n-l)(i+k-n-l-2) \cdots (i+k-n-m)}{(-1)^{m-l}(m+1-l)!(l-1)!}. \end{aligned}$$

**Lemma 3.6.**  $y(x) = \frac{a_n x^n + a_{n-1} x^{n-1} + \dots + a_0}{b_m x^m + b_{m-1} x^{m-1} + \dots + b_0} \Leftrightarrow \mathcal{R}_{n,m}(y(x)) = 0$  where  $\mathbf{E}(a_i) = a_i$ ,  $\mathbf{E}(b_j) = b_j$ .

Using properties of the proper parametrization of algebraic curves, the degree bound for the rational solution can be given [20].

**Theorem 3.7.** Let  $F(y, y_1) = 0$  be a first order autonomous  $O\Delta E$ . If  $y(x) \in \bar{\mathbb{Q}}(x) \setminus \bar{\mathbb{Q}}$  is a rational solution of  $F = 0$ , then  $\deg(y(x)) = \deg(F, y_1) = \deg(F, y)$ .

Similar to the differential case, the rational solutions can be found with the help of rational parametrization of algebraic curves [20].