

THE INNER EAR

INCLUDING

OTONEUROLOGY, OTOSURGERY, AND
PROBLEMS IN MODERN WARFARE

By

JOSEPH FISCHER, M.D.

*Staff Member, Beth Israel Hospital, Boston;
formerly Associate of Adam Politzer and Senior
Otolaryngologist with Gustav Alexander,
Policlinic of Vienna*

AND

LOUIS E. WOLFSON, M.D.

*Instructor in Ear, Nose, and Throat, Tufts
Medical School; Senior Surgeon in Ear, Nose,
and Throat, Boston Dispensary*



1943

GRUNE & STRATTON

NEW YORK

COPYRIGHT 1943
GRUNE & STRATTON, INC.
443 Fourth Avenue
New York, N. Y.

PRINTED BY
WAVERLY PRESS, INC.
U. S. A.

Authors' Prefaces

Through twenty years of postgraduate lecturing, I have constantly been asked the same question: "What book would you recommend for studies on the labyrinth and on otoneurologic problems?" I could never answer satisfactorily, because all that is available are either the ordinary textbooks that deal with the subject in a rather brief and elementary way, or the encyclopedias and handbooks of anatomy, physiology, neurology, etc., in which the respective chapters comprise large-scale dissertations delving into trifling details, so that the reader soon finds himself lost in the depths of science. One could also use the most important original articles on the labyrinth by authors of the various countries. However, this requires a certain knowledge of the subject, and would not give a satisfactory survey of the knowledge of the labyrinth.

Repeatedly I was asked by my students to write down my lectures on the labyrinth, which included instruction on anatomy, physiology, clinical practice, and operative technic on cadavers. I am fully aware of the great difference between the cold printed words of a book and the living lecture using anatomic specimens or patients, with demonstration of surgical procedures, and followed by a general discussion. Despite this, I have finally decided to bridge the gap between the usual textbooks and the voluminous encyclopedias and to write a treatise on the subject. Since the diseases of the middle ear are exhaustively discussed in the various textbooks, and since the function of the cochlear apparatus is generally known, the main attention is given in this book to the labyrinth and its central pathways.

Before I began on this difficult task, I scrutinized my registration book, which contains the signatures of all the students who have taken the various courses with me. I found two main groups of students. One group consisted of rather older otolaryngologists with great clinical experience and skill, who could not, however, keep pace with current progress in theory and research. The other group was represented by well educated young physicians with great theoretic knowledge (chiefly interns and residents) but without clinical experience.

This differentiation gave me the basis for the structure of my book. Almost every topic occurs twice—once in the chapters on physiology, where the various theories and hypotheses are discussed and references to the bibliographic material are brought out, and again in the chapter on functional tests, where indications, technic, clinical significance, and evaluation of the various tests are critically described. In this manner each reader can find just what he is looking for without having to plow through pages of theory and literature.

In the discussion of the inflammatory diseases of the labyrinth, a new

principle of classification is used, as outlined by my former chief Alexander and myself. Particular stress has been placed on the pathologic anatomy of the various forms of labyrinthine disease; each is illustrated by photomicrographs. This emphasis stems from my conviction that correct diagnosis and proper treatment depend on an exact knowledge of the pathologic basis. The indications for surgical intervention, as used in our clinic, and the technics of the various operations have been outlined. In the complex matter of otoneurology, the important role played by the otologist in rendering diagnostic aid to the neurologist is exemplified in the differentiation between peripheral and central lesions.

The illustrations in the chapters on anatomy and physiology are, for didactic reasons, chiefly schematic and diagrammatic, while in the clinical sections photomicrographs are shown. The diagrams were made by Dr. Louis Bergmann, to whom I wish to express my appreciation for his artistic drawings. All macro and microphotographs used in this book are from specimens in my own collection.

JOSEPH FISCHER

As an otologist in active practice for more than twenty years, the various special problems of this field of medicine have been of particular interest to me. Every year or two I have managed to visit several of the special clinics either in this country or abroad. It has been especially interesting and instructive to see the various ways in which our present otologic problems are treated.

In this book, an effort has been made to review the important contributions in otology and also their relation to otoneurology. The subjects of Ménière's disease and otosclerosis are brought thoroughly up to date. In describing the intracranial complications, particular attention is given to the processes in the posterior cranial fossa, with special reference to their connection with the labyrinth. In the section on cerebellar abscess, the diversity of opinion between the otologist and neurosurgeon is critically discussed, and the interrelationship of their domains is shown in the discussion of the various diseases of the nervous system.

I wish to give grateful thanks to my many teachers and friends, many of whom are no longer with us: Neumann, Alexander, and Ruttin of Vienna; Jansen, Bruehl, and Passow of Berlin; Holmgren of Stockholm; Bárány of Upsala; Le Maître of Paris. And I wish to express my appreciation to Lempert in New York and Hughson in Abington, Pennsylvania, for their many kindnesses to me.

LOUIS E. WOLFSON

Boston, February, 1943

Contents

AUTHORS' PREFACES.....	xi
CHAPTER	
I. CLINICAL ANATOMY	
1. Nomenclature.....	1
2. Topographic relations within the petrous bone.....	1
3. The bony inner ear.....	4
4. The membranous inner ear.....	10
5. The nervous apparatus of the inner ear.....	11
The facial nerve. The cochlear nerve. The labyrinth nerve and its peripheral termination. The central labyrinthine ap- paratus	
6. Blood supply of the inner ear.....	17
7. The dura mater and the venous sinuses.....	18
8. Communications of the inner ear with the cranial fossa.....	21
9. Topographic relation of the inner ear to the brain.....	22
Relation to the cerebellum. Relation to the cerebrum	
References.....	25
II. GENERAL PHYSIOLOGY.....	28
1. Comparative physiology.....	28
2. Physiology of the human labyrinth.....	29
3. Histophysiologic factors.....	30
4. Function of the cristae.....	31
5. Function of the maculae (otolithic organ).....	34
6. Linear movements.....	36
7. Function of the saccule.....	37
8. Separation of macula and crista functions.....	38
9. Influence of the labyrinth on the striated musculature (tonus) ..	39
10. Influence of the labyrinth on the autonomous system.....	39
11. Kinetovisual function.....	40
12. Summary.....	41
References.....	41
III. APPLIED PHYSIOLOGY.....	45
SPONTANEOUS MANIFESTATIONS	
1. Vertigo.....	45
Definitions. Anatomic pathways. Differentiation of periph- eral and central vertigo	
2. Spontaneous disturbances of equilibrium and coordination.....	48
Comparative physiologic considerations. Sense of statics. In- terpretation of the various mechanisms	
3. Spontaneous nystagmus.....	50
Principles and definitions. Nystagmography and cinematogra- phy. Genesis of the slow phase. Genesis of the fast phase. Central localization. Influence of the cerebrum. Influence of the cerebellum. Labyrinthine reflex arc. Principles of clinical classification. Mechanism of end position nystagmus. Char- acter and significance of positional nystagmus	
4. Conjugate paresis and conjugate deviation.....	62
Optic centers. Central pathways. Vertical conjugate paresis	

5. Convergence spasm.....	64
6. Labyrinthine strabismus.....	65
References.....	66
INDUCED REACTIONS	
1. Optokinetic reaction.....	69
Mechanism of the nystagmus. Clinical significance	
2. Caloric reaction.....	71
Historical résumé. Theories of caloric reaction. Quantitative determinations. Peripheral and central excitability. Determination of labyrinthine threshold. Warm and cold stimulations	
3. Bilateral calorization.....	78
Clinical importance. Theories of the method	
4. Turning reaction.....	80
Etiology. Anatomic-physiologic factors. Rhythmic phases of postrotatory nystagmus. Phenomenon of interference. Theories of the short-turning methods. Quantitative determination	
5. Mechanical reaction (fistula symptom).....	86
Definition and pathology. Pseudo fistula symptom	
6. Galvanic reaction.....	87
The theories of galvanic reaction. Chronaxia	
7. Tonic reflexes and reactions.....	90
A. Postural reflexes (otolith reflexes). B. Movement reflexes (semicircular reflexes). C. Reflexes upon inadequate stimuli	
References.....	96
IV. FUNCTIONAL TESTS.....	100
SPONTANEOUS MANIFESTATIONS	
1. Vertigo.....	100
Types of labyrinthine vertigo. Clinical history. Relation of spontaneous to induced vertigo. Diseases accompanied by vertigo	
2. Spontaneous disturbances of equilibrium and coordination.....	104
Examination of the static labyrinth. Examination of the kinetic equilibrium. Examination for coordination disturbances. Evaluation of results of examination. Differential diagnosis of disturbances of equilibrium	
3. Spontaneous nystagmus.....	109
Clinical importance of nystagmus. Aids in local diagnosis. Diseases of the peripheral labyrinth associated with spontaneous nystagmus. Diseases of the central vestibular organ associated with spontaneous nystagmus. Diseases of the ocular organ associated with spontaneous nystagmus. Rare forms of nystagmus. Tests for positional nystagmus	
4. Conjugate paresis and conjugate deviation of the eyeballs.....	118
Clinical importance. The various types of conjugate paresis and deviation. Diseases associated with disturbances of the conjugate eye movements. Differential diagnosis of cortical and pontine lesions	
References.....	120

INDUCED REACTIONS

1. Optokinetic tests..... 122
Technic. Differentiation of labyrinthine and nonlabyrinthine nystagmus (inversion)
 2. Caloric tests..... 124
Mass or volume method. Minimal methods. Significance of the various tests. Evaluation of caloric excitability
 3. Bilateral calorization..... 131
Technic. Evaluation of results
 4. Turning tests..... 133
Forced-turning methods. Short-turning methods. Long-turning methods. Comment on the various methods. Evaluation of labyrinthine excitability
 5. Mechanical tests (fistula symptom)..... 139
Technic. Evaluation of results
 6. Galvanic test..... 141
Technic. Comment on the various methods
 7. Tonic reflexes and reactions..... 144
Postural reflexes and reactions (otolithic reflexes). Movement reactions and reflexes. Reactions and reflexes upon inadequate stimuli
- References..... 156

EVALUATION OF ABNORMAL REACTIONS

1. Changes in relation to certain stimuli..... 159
Loss of turning reaction, normal caloric reaction. Loss of caloric reaction, normal turning reaction. Loss of caloric and turning reaction, normal galvanic reaction. Loss of galvanic reaction, normal caloric reaction. Reactive dissociation
2. Changes in certain semicircular canals..... 160
3. Vestibular disharmony..... 161
Absence of primary reaction (nystagmus). Absent or atypical falling reaction. Absent or atypical past-pointing reaction. Absence of girdle reflex
4. Hyperexcitability..... 162
5. Hypo-excitability..... 163
6. Loss of excitability..... 164
7. Qualitative changes..... 164
Disturbance of rhythm. Decrease or loss of the quick component. Conjugate deviation plus nystagmus. Nystagmus plus conjugate deviation. Inverted nystagmus. Perverted nystagmus

DIFFERENTIATION OF PERIPHERAL AND CENTRAL LESIONS

1. Significance of spontaneous manifestations..... 166
Vertigo. Spontaneous disturbances of equilibrium and coordination. Spontaneous nystagmus. Positional nystagmus. Associated eye movements
2. Significance of induced reactions..... 169
Optokinetic nystagmus. Caloric nystagmus. Bilateral calorization. Turning nystagmus. Fistula symptoms. Galvanic

	reaction. Postural reflexes and reactions. Head-moving test. Falling reaction. Past-pointing reaction. Clinical significance of comparison of tests	
V. PRIMARY DISEASES OF THE LABYRINTHINE CAPSULE: OTOSCLEROSIS		176
1. Historical viewpoints		176
2. Pathologic anatomy		177
3. Etiology	Theories of inflammation. Theories of disturbances of circulation. Theory of tumor. Theory of congenital origin. Theory of constitution. Theory of trauma. Theory of systemic diseases of the skeleton. Theory of disturbed endocrine glands. Theory of vitamin deficiency	181
4. Clinical diagnosis		186
5. Treatment	Conservative methods. Surgical methods	187
References		193
VI. INFLAMMATORY DISEASES OF THE INNER EAR		195
LESIONS OF THE LABYRINTH		
1. Otitis interna serosa	Etiology. Pathologic anatomy. Clinical course. Differential diagnosis. Treatment	196
2. Para-otitis interna purulenta	Etiology. Pathologic anatomy. Clinical course. Treatment	199
3. Peri-otitis interna purulenta	Etiology. Pathologic anatomy. Clinical course. Treatment	201
4. Otitis interna purulenta	Etiology. Pathologic anatomy. Clinical course. Treatment	206
5. Otitis interna purulenta complicata	Etiology. Pathologic anatomy. Clinical course	216
6. Otitis interna traumatica	Skull fractures. Gunshot wounds. Foreign-body injuries. Operative damages	217
7. Indications for surgery		221
8. Operative technic	Hinsberg-Bourguet method. Jansen-Neumann method. Operative injuries, technical precautions	223
INFLAMMATION OF THE PETROUS PYRAMID		
1. Pathways of infection: pathology		231
2. Clinical diagnosis		232
3. Treatment		233
References		234
VII. INTRACRANIAL LABYRINTHOGENIC COMPLICATIONS		238
1. Viewpoints and classification		238
2. Extradural lesions (pachymeningitis externa)	Etiology and pathologic anatomy. Symptoms and diagnosis. Treatment	240
3. Intradural lesions	Empyema of the endolymphatic sac. Sinus thrombosis	245
4. Subdural lesions	Pachymeningitis interna (subdural abscess). Leptomeningitis. Serous meningitis. Purulent meningitis	252

5. Cerebellar abscess.....	262
General considerations. Etiology. Paths of infection. Pathologic anatomy. Bacteriology. Symptomatology. Diagnosis. Differential diagnosis. Operative treatment	
6. Temporoparietal abscess.....	283
Symptomatology. Treatment	
References.....	287
VIII. CHEMOTHERAPY.....	292
1. Classification and general use of the sulfa drugs.....	292
Sulfanilamide. Sulfapyridine. Sulfathiazole. Sulfadiazine	
2. Use of sulfa drugs in otology.....	296
References.....	300
IX. FACIAL PALSY.....	301
1. Anatomic and physiologic basis.....	301
2. Classification and etiology.....	303
Otogenic (inflammatory) facial palsy. Facial palsy in connection with skull trauma. Facial palsy due to operative injuries. Facial palsy in connection with tumors. Facial palsy of unknown origin	
3. Clinical signs and symptoms.....	306
4. Treatment.....	308
Causal treatment. Conservative treatment. Surgical repair	
References.....	312
X. CONGENITAL DISEASES.....	314
1. Changes of the bony inner ear.....	314
Systemic diseases of the bones. Otosclerosis. Arrested malformations	
2. Changes of the nerve and sensory apparatus.....	316
Deaf-mutism. Chronic progressive inner-ear deafness. Anomalies of the labyrinth	
References.....	319
XI. NEOPLASMS.....	320
1. Neoplasms of the inner ear.....	320
Endothelioma (meningeoma). Sarcoma. Carcinoma	
2. Neoplasms of the cerebellopontile angle.....	327
Acoustic tumors (neurinoma). Other angle tumors	
3. Tumors of the anterior and middle cranial fossae: congestive inner ear.....	331
Conclusions	
References.....	333
XII. VASCULAR LESIONS: MÉNIÈRE'S SYNDROME.....	334
1. Classification.....	334
2. Pathologic anatomy.....	334
3. Etiology.....	335
Theories of causation	
4. Symptomatology.....	338
5. Treatment.....	340
Medical therapy. Surgical therapy	
References.....	343

XIII. WAR TRAUMA.....	345
1. Skull fractures.....	345
Symptomatology. Treatment	
2. Concussion of the inner ear.....	355
Symptomatology	
3. Trauma due to explosions.....	360
Ear symptoms due to a sudden blast (acute trauma). Ear symptoms due to repeated short explosions (chronic trauma). Ear symptoms due to concussion of the brain. Ear symptoms due to concussion of the brain and of the inner ear. Ear symptoms due to brain concussion and fracture of the skull. Ear symptoms of psychogenic nature	
4. Gunshot injuries.....	366
Treatment	
References.....	368
XIV. THE ROLE OF THE INNER EAR IN AERONAUTICS.....	371
1. Physiology of the cochlea and labyrinth in the usual environment.....	371
Cochlea. Labyrinth	
2. Physiology and pathology of the cochlea and labyrinth in the unusual environment (air).....	376
Cochlea. Labyrinth. (Blind flight; Combat flight; Airsickness)	
3. Functional tests for aviation fitness.....	386
Cochlea. Labyrinth	
References.....	390
XV. EFFECTS OF ATMOSPHERIC PRESSURE CHANGES ON THE EAR.....	392
1. Effects of increased atmospheric pressure.....	393
2. Effects of decreased atmospheric pressure.....	394
3. Effects of sudden changes in atmospheric pressure.....	397
4. Regulations for examination of flyers.....	402
Eustachian tube patency. Nose and paranasal sinuses. Throat	
References.....	403
INDEX.....	405

I

Clinical Anatomy

By *Joseph Fischer*

1. NOMENCLATURE

THE DESIGNATION "clinical anatomy" is given to these studies to differentiate them from the descriptive treatment of anatomy found in most textbooks. The main attention is given to topographic relationships with respect to surgical procedures and to those anatomic details that have a clinical significance. Since it is often difficult to visualize anatomic relationships, illustrations are used chiefly in a schematic or half-schematic way to simplify matters, while for demonstration of pathology only photomicrographs are used.

The best way to get a proper conception of topographic relationships is to dissect petrous bones in various planes. Those who are interested in this subject are referred to the book about macroscopic and microscopic dissection of the ear by G. Alexander and J. Fischer.

In discussing the anatomy of the inner ear, I follow the nomenclature proposed by my former chief, Alexander. He divides the inner ear into cochlea and labyrinth. The latter consists of the vestibule and the semicircular canals. Hence the term "labyrinth" is applied to the non-cochlear part of the inner ear. The *nervus vestibularis* is spoken of as *nervus labyrinthicus*. The same holds true for the *ganglion vestibulare*, which he calls *ganglion labyrinthicum*.

2. TOPOGRAPHIC RELATIONS WITHIN THE PETROUS BONE

If it were possible to visualize a transparent petrous bone, with the inner-ear spaces filled with stained fluids, three spaces would be outlined: (1) a *perilabyrinthine* space, between the surface of the petrous bone and the inner ear; (2) a *perilymphatic* space, between the bony and the membranous inner ear; (3) an *endolymphatic* space within the membranous inner ear (FIG. 1).

The *perilabyrinthine space* is not very definite and shows great anatomic variation. There may be a whole system of pneumatic cells extending from the mastoid, antrum, epitympanum, or eustachian tube down to the tip of the pyramid, or there may be very few cells or none at all—so that it cannot properly be called a space in the same sense as the *perilymphatic* and *endolymphatic* spaces, which are real, definitely outlined spaces filled

with fluid. The clinical importance of this region can best be seen in the fact that most of the anatomic studies in the various countries are made not by anatomists but by otologists.

Mouret and Portmann (France) often found on the tip of the pyramid large cells communicating with the perilyabyrinthine space. They described two groups of cells in the apex: one, the external inferior group, originating from the eustachian tube; the other, the internal superior group, extending from the labyrinthine cells. A suppuration of the pyramidal tip can therefore come either directly from the eustachian




FIG. 1. SCHEMATIC ILLUSTRATION OF INNER EAR

black = endolymphatic space; white = perilymphatic space; *pl* = perilyabyrinthine cells; *im* = internal auditory meatus; *es* = endolymphatic sac; *ca* = cochlear aqueduct; *va* = vestibular aqueduct; *s* = saccule; *u* = utricle; *c* = cochlea; *sv* = superior vertical canal.

tube or indirectly (over the perilyabyrinthine cells) from the tympano-antral cells. Belinoff and Balan (Bulgaria) examined 40 petrous bones and found three different types of pyramidal structure, namely, the pneumatic, the diploetic, and the mixed type. In 62 per cent of these bones, the structure of the pyramidal tip did not correspond with the structure of the mastoid.

Myerson, Rubin, and Gilbert studied the anatomy of 200 petrous bones and found pneumatization of the apex in 11 per cent, while Hagens observed it in 34 per cent. Frenckner (Sweden) found in more than one-

fourth of his cases a pneumatized apex. He described the various cell passages, such as the posterosuperior, the inferior, and the tractus sub-arcuatus. The various cell systems joined and formed a common cell system within the pyramid.

J. R. Lindsay (Chicago) examined 100 petrous bones with the following results: pneumatization in the posterosuperior area (with cells from the epitympanum) in 36 per cent, pneumatization in the posteromedial area (with cells from the mastoid) in 25 per cent, pneumatization in the infralabyrinthine area (with cells from the mastoid to the jugular bulb or round window niche) in 25 per cent, pneumatization in the tip of the pyramid in 21 per cent.

The clinical importance of this region has been stressed by Eagleton, Friesner, Ramadier, Kopetzky, Brunner, etc. (p. 231).

The *perilymphatic space* lies between the bony and the membranous inner ear, and is filled with a fluid called the perilymph. Its communication outside the petrous pyramid is called the aquaeductus cochleae. It has two openings, one within the inner ear—*apertura interna*—in the *scala tympani*, or first turn of the cochlea, and the other—*apertura externa*—outside the pyramid on its inferior surface, as shown in FIGURE 1. There the duct debouches subdurally into the cerebrospinal spaces containing cerebrospinal fluid, and therefore communicates with the brain spaces. This anatomic fact is of great practical importance, since this is a pathway of infection from the inner ear to the cranial fossa in cases of meningitis and brain abscess. Conversely, pathologic conditions such as brain tumor, hydrocephalus, etc., which are accompanied by increased intracranial pressure, may likewise sooner or later produce changes in the inner ear (congestive inner ear or choked labyrinth), as I have shown in microscopic studies.

The *endolymphatic space* lies within the membranous inner ear and contains a fluid called the endolymph. Its communication is called the aquaeductus vestibuli. It has an internal aperture in the vestibule, where the canal begins from the ductus utriculosaccularis, and an external opening, located on the posterior surface of the petrous bone, halfway between the internal auditory meatus and the lateral sinus (FIG. 1). Studies on the anatomy of the endolymphatic duct and sac have been carried out recently by Guild, Anson, Bast, etc.

Bast described a valve that projects into the utricle and guards the utricular opening into the utriculo-endolymphatic duct. He believes that the valve prevents a sudden outflow of endolymph from the utricle. Perlman and Lindsay corroborate the presence of a utriculo-endolymphatic valve in man and in certain animals, which functions in maintaining the content and volume of the utricle and the semicircular canals.

The chemical composition of the peri- and endolymph fluids is the same, but anatomically they are separated, because the endolymphatic system is a closed system blindly intradural.

The dura in this region is very adherent and can easily be torn in the course of a labyrinth operation. It is therefore advisable to sever the




FIG. 2. DIAGRAM SHOWING TOPOGRAPHY OF INNER EAR

blue = sigmoid sinus; *ca* = carotid artery; *et* = eustachian tube; *VII* = facial nerve; *co* = cochlea; *l* = labyrinth.

attachment with a sharp instrument before chiseling on the posterior surface of the pyramid.

FIGURE 2 shows schematically the topographic relations of the various parts within an assumed transparent petrous bone.

3. THE BONY INNER EAR

The bony inner-ear structure consists of three parts, a central, an anterior, and a posterior part; these are, in order, the vestibule, the cochlea, and the semicircular canals.

The *vestibule* is oval-shaped. For purposes of demonstration, we shall assume a cube with six walls. The lateral wall constitutes also the medial wall of the middle ear, as seen in FIGURE 3, which shows the two windows—the fenestra ovalis, closed by the footplate of the stapes, and the fenestra rotunda, closed by the secondary membrane. Between these two windows




FIG. 3. VERTICAL SECTION THROUGH MIDDLE AND INNER EAR

pr = promontory; *rw* = round window; *ow* = oval window; *stf* = stapes footplate; *fa* = facial nerve; *cah* = crista ampullae horizontalis; *mu* = macula utriculi; *u* = utricle; *cp* = perilymphatic cistern; *ic* = inferior vertical canal.

is the promontory, which corresponds to the first turn of the cochlea. Above the oval window lies the fallopian canal, and above that, almost in the superior wall of the vestibule, the prominence of the horizontal semi-circular canal is visible. The medial wall of the vestibule (FIG. 4) shows two depressions in the bone, one lying superoposteriorly, called the re-

cessus ellipticus, and the other in front and below, called the recessus sphericus. In the medial wall the mouth of the vestibular aqueduct and the openings for the nerves coming from the internal auditory meatus are seen. The latter are called the maculae cribrosae. On the anterior wall of the vestibule is the large opening for the cochlea, and on the posterior wall the opening for the semicircular canals can be seen.

The *cochlea*, which represents the anterior part of the middle ear, consists of two parts: one is called the vestibular portion, the other the cochlear




FIG. 4. DIAGRAMMATIC SECTION SHOWING MEDIAL WALL OF BONY LABYRINTH WITH TWO DEPRESSIONS

re = recessus ellipticus; *rs* = recessus sphericus; *sct* = scala tympani; *scv* = scala vestibuli; *ap* = ampullar ends of semicircular canals; *1* = cochlear aqueduct; *2* = vestibular aqueduct; *3* = horizontal canal; *4* = crus commune; *5* = macula cribrosa superior; *6* = macula cribrosa media; *7* = macula cribrosa inferior.

body. The latter shows two and a half turns, i.e., one basal whorl, one middle whorl, and one half whorl at the top. The blind end on top is called the cupula. The cochlea contains, on the inner bony wall, a bony septum following the same contour as the cochlea itself, and called the *membrana spiralis ossea*. This bony septum divides the space in the cochlea into two compartments. The lower compartment is called the *scala tympani*, and the upper the *scala vestibuli*. There is a large communication between the two *scalae* at the top of the cochlea, because of the absence of the bony membrane at this point; this passage is called the