

TUMORS of the PARATHYROID GLANDS

by

BENJAMIN CASTLEMAN, M. D.


and

SANFORD I. ROTH, M. D.



AFIP

TUMORS OF THE PARATHYROID GLANDS



ATLAS OF TUMOR PATHOLOGY

Second Series

Fascicle 14

TUMORS OF THE PARATHYROID GLANDS

by

BENJAMIN CASTLEMAN, M. D.

Shattuck Professor of Pathological Anatomy, Emeritus
Harvard Medical School

and

Honorary Pathologist, Massachusetts General Hospital
Boston, Massachusetts 02114

and

SANFORD I. ROTH, M. D.

Professor and Chairman, Department of Pathology
University of Arkansas College of Medicine
Little Rock, Arkansas 72201

Published by the
ARMED FORCES INSTITUTE OF PATHOLOGY
Washington, D. C.

Under the Auspices of
UNIVERSITIES ASSOCIATED FOR RESEARCH AND EDUCATION IN PATHOLOGY, INC.
Bethesda, Maryland
1978

Accepted for Publication
1977

.....
For sale by the
Armed Forces Institute of Pathology
Washington, D. C. 20306

ATLAS OF TUMOR PATHOLOGY

Sponsored and Supported by

AMERICAN CANCER SOCIETY*
NATIONAL CANCER INSTITUTE, NATIONAL INSTITUTES OF HEALTH*
and
ARMED FORCES INSTITUTE OF PATHOLOGY

EDITOR

WILLIAM H. HARTMANN, M. D.

Professor and Chairman, Department of Pathology
Vanderbilt University School of Medicine
Nashville, Tennessee 37232
Area Code 615 - 322-6609

COEDITOR

WILLIAM R. COWAN, COLONEL, USAF, MC

Armed Forces Institute of Pathology
Washington, D. C. 20306

EDITORIAL ADVISORY COMMITTEE

Pelayo Correa, M.D.	Louisiana State University School of Medicine New Orleans, Louisiana 70112
Harlan I. Firminger, M.D.	University of Colorado Medical Center Denver, Colorado 80220
David G. Freiman, M.D.	Harvard Medical School, Beth Israel Hospital Boston, Massachusetts 02215
Myron R. Melamed, M.D.	Memorial Sloan-Kettering Cancer Center New York, New York 10021
Henry C. Pitot, M.D.	University of Wisconsin Medical Center Madison, Wisconsin 53706
Richard J. Reed, M.D.	Tulane University School of Medicine New Orleans, Louisiana 70112
Juan Rosai, M.D.	University of Minnesota Medical School Minneapolis, Minnesota 55455
John H. Yardley, M.D.	The Johns Hopkins School of Medicine Baltimore, Maryland 21205

*(Supported by Public Health Service grant no. CA08575 from the
National Cancer Institute and American Cancer Society grant no. MG-122)

EDITORS' NOTE

The Atlas of Tumor Pathology was originated by the Committee on Pathology of the National Academy of Sciences—National Research Council in 1947. The form of the Atlas became the brainchild of the Subcommittee on Oncology and was shepherded by a succession of editors. It was supported by a long list of agencies; many of the illustrations were made by the Medical Illustration Service of the Armed Forces Institute of Pathology; the type was set by the Government Printing Office; and the final printing was made by the press at the Armed Forces Institute of Pathology. The American Registry of Pathology purchased the fascicles from the Government Printing Office and sold them at cost, plus a small handling and shipping charge. Over a period of 20 years, 15,000 copies each of 40 fascicles were produced. They provided a system of nomenclature and set standards for histologic diagnosis which received worldwide acclaim. Private contributions by almost 600 pathologists helped to finance the compilation of an index by The Williams & Wilkins Company to complete the original Atlas.

Following the preparation of the final fascicle of the first Atlas, the National Academy of Sciences—National Research Council handed over the task of further pursuit of the project to Universities Associated for Research and Education in Pathology, Inc. Grant support for a second series was generously made available by both the National Cancer Institute and the American Cancer Society. The Armed Forces Institute of Pathology has expanded and improved its press facilities to provide for a more rapid and efficient production of the new series. A new Editor and Editorial Advisory Committee were appointed, and the solicitation and preparation of manuscripts continues.

This second series of the Atlas of Tumor Pathology is not intended as a second edition of the first Atlas and, in general, there will be variation in authorship. The basic purpose remains unchanged in providing an Atlas setting standards of diagnosis and terminology. Throughout the rest of this new series, the term chosen for the World Health Organization's series "International Histological Classification of Tumours" (when available) is shown by an asterisk if it corresponds to the authors' choice, or as the first synonym in bold print if it differs from the authors' heading. Hematoxylin and eosin stained sections still represent the keystone of histologic diagnosis; therefore, most of the photomicrographs will be of sections stained by this technic, and only sections prepared by other technics will be specifically designated in the legends. It is hoped that in many of the new series a broader perspective of tumors may be offered by the inclusion of special stains, histochemical illustrations, electron micrographs, data on biologic behavior, and other pertinent information when indicated for a better understanding of the disease.

The format of the new series is changed in order to allow better correlation of the illustrations with the text, and a more substantial cover is provided. An index will be included in each fascicle.

It is the hope of the Editors, past and present, the Editorial Advisory Committees, past and present, and the Sponsors that these changes will be welcomed by the readers. Constructive criticisms and suggestions will always be appreciated.

William H. Hartmann, M. D.
William R. Cowan, M. D.

ACKNOWLEDGMENTS

The authors are indebted to many individuals and to many of the staff of the Massachusetts General Hospital in the Departments of Medicine, Surgery, and Pathology who have helped and encouraged us over the years in our studies of the parathyroid glands. Dr. Chiu-An Wang, Associate Clinical Professor of Surgery at the Massachusetts General Hospital, has been especially generous in obtaining follow-up information on the patients and in allowing us to use his clinical data on his patients and those from the Massachusetts General Hospital. Dr. Oliver Cope, Professor of Surgery Emeritus, has always been a great help and inspiration in our study of these patients. Our co-workers and collaborators are too numerous to mention by name, but to each we express our gratitude. We are grateful to Mrs. Betty U. McNeely who edited the manuscript with her usual care and precision, and Mrs. Pauline Park and Mrs. Carol Kuykendall who typed the manuscript. The preparation of the electron micrographs was the result of the outstanding effort of Mrs. Sendy P. C. Su.

Many of the experimental studies described here have been the result of research support by the National Cancer Institute of the National Institutes of Health.

Finally, we wish to thank the following for having generously contributed microscopic slides, photographs, photomicrographs, or charts for use in this fascicle: Dr. A. von Albertini and Dr. F. Koller, Zurich, Switzerland; Dr. F. M. Vellios, Dallas, Texas; Dr. C-A. Wang, Boston, Massachusetts; and Dr. M. Wolman, Tel Aviv, Israel.

Benjamin Castleman, M. D.
Sanford I. Roth, M. D.

Permission to use copyrighted illustrations has been granted by:

American Cancer Society, Inc.:

Cancer 38:1668-1675, 1976. For figures 63, 65, and 66.

Dun-Donnelley Publishing Corporation:

Am. J. Med. 50:612-622, 1971. For figure 4.

Harper & Row Publishers, Inc.—Hoeber Medical Division:

Am. J. Pathol. 13:553-574, 1937. For figures 85 and 89.

Am. J. Pathol. 77:493-506, 1974. For figure 11.

Am. J. Pathol. 84:521-527, 1976. For plate III.

Hospital for Joint Diseases and Medical Center:

Bull. Hosp. Joint Dis. 12:368-378, 1951. For figure 70; plate IV.

J. B. Lippincott Company:

Ann. Surg. 104:9-35, 1936. For figure 70.

Ann. Surg. 175:448-453, 1972. For figure 96.

Ann. Surg. 183:271-275, 1976. For figures 1 and 2.

Prof. W. G. Spector, Editor, J. Pathol.

J. Pathol. & Bact. 44:431-462, 1937. For data in figure 4.

The Massachusetts Medical Society:

N. Engl. J. Med. 266:121-123, 1962. For figures 50 and 51.

N. Engl. J. Med. 285:422-429, 1971. For figure 82.

The Williams & Wilkins Company:

Lab. Invest. 10:354-360, 1961. For figure 10.

TUMORS OF THE PARATHYROID GLANDS

Contents

	Page No.
Introduction	1
History	1
Normal Parathyroid Gland	3
Embryology	3
Gross Anatomy	6
Microscopic Anatomy	6
Ultrastructure	11
Blood Supply	13
Primary Hyperparathyroidism	21
Adenoma	29
Lipoadenoma	52
Chief-cell Hyperplasia	54
Clear-cell Hyperplasia	67
Carcinoma	74
Therapy	79
Malignant Nonparathyroid Tumors Associated with Hypercalcemia and Hypophosphatemia	82
Secondary Hyperparathyroidism	83
Tertiary Hyperparathyroidism	89
Nonfunctional Enlargement	90
Oxyphil Adenoma	90
Primary Carcinoma	90
Metastatic Carcinoma	90
Parathyroid Cysts	91
Index	93

TUMORS OF THE PARATHYROID GLANDS

INTRODUCTION

The tumors referred to in the title of this Fascicle are generally those that cause enlargement of the parathyroid glands. The discussion, therefore, concerns lesions that produce primary hyperparathyroidism—adenomas, hyperplasias, and carcinomas. In addition, other functioning and nonfunctioning lesions of the parathyroid glands will be discussed briefly.

Since publication of the first Fascicle on this subject in 1952, there have been numerous advances in the understanding of the ultrastructure, biochemistry, and physiology of normal and abnormal parathyroid glands. The increased clinical recognition of hyperparathyroidism and the routine determination of the serum calcium level in the multiphasic screening of apparently healthy

persons have led to an earlier diagnosis of this disease, thus providing the pathologist with a large number of surgical specimens to broaden his experience. A new entity, primary chief-cell hyperplasia, involving all four parathyroid glands has been recognized (Cope et al.). The differentiation between adenoma and the rare carcinoma of the parathyroid gland has been refined.

References

- Castleman, B. Tumors of the Parathyroid Glands. Fascicle 15, Atlas of Tumor Pathology. Washington: Armed Forces Institute of Pathology, 1952.
- Cope, O., Keynes, W. M., Roth, S. I., and Castleman, B. Primary chief-cell hyperplasia of the parathyroid glands: A new entity in the surgery of hyperparathyroidism. *Ann. Surg.* 148:375-388, 1958.

HISTORY

The parathyroid glands were first described in the rhinoceros in 1862 by Owen. Virchow, in 1863, was the first to describe this gland in man, but it was Sandström, in 1880, who presented the first relatively complete comparative description of the gland and gave it the present name. Subsequently, there have been numerous comparative studies of the parathyroid glands in various species. In 1903, Askanazy re-

ported a parathyroid tumor (an adenoma) found at autopsy in a patient with von Recklinghausen's disease (osteitis fibrosa cystica generalisata). Since that time, other authors have made the connection between the parathyroid tumor and the bone disease, but it remained for Mandl, a Viennese surgeon, in 1925, to remove a parathyroid tumor from a living patient with von Recklinghausen's disease.

In the United States, the first patient operated on for primary hyperparathyroidism, Captain Charles Martell, required six neck explorations, the first in 1926 at the Massachusetts General Hospital, before the tumor was finally discovered in the mediastinum in 1932. The first successful parathyroid exploration in the United States was at the Barnes Hospital, St. Louis, in 1929 (Barr and Bulger). Albright and his co-workers, in 1934, recognized the entity of clear-cell hyperplasia of the parathyroid glands. Hall and Chaffin, in 1935, reported the first carcinoma of a parathyroid gland. Castleman and Wilens described the histologic changes in the secondary parathyroid hyperplasia that occurs as a result of chronic renal disease.

References

- Albright, F. A page out of the history of hyperparathyroidism. *J. Clin. Endocrinol. Metab.* 8:637-657, 1948.
-, Bloomberg, E., Castleman, B., and Churchill, E. D. Hyperparathyroidism due to diffuse hyperplasia of all parathyroid glands rather than adenoma of one. *Arch. Intern. Med.* 54:315-329, 1934.
- Askanazy, M. Ueber ostitis deformans ohne osteoides Gewebe. *Arb. Geb. Pathol. Anat. Inst. Tubingen Leipz.* 4:398-422, 1904.
- Axelrod, L. Bones, stones, and hormones: The contributions of Fuller Albright. *N. Engl. J. Med.* 283:964-970, 1970.
- Barr, D. P. and Bulger, H. A. The clinical syndrome of hyperparathyroidism. *Am. J. Med. Sci.* 179:449-476, 1930.
- Bauer, W. and Federman, D. D. Hyperparathyroidism epitomized: The case of Captain Charles E. Martell. *Metabolism* 11:21-29, 1962.
- Castleman, B. and Mallory, T. B. The pathology of the parathyroid gland in hyperparathyroidism. A study of 25 cases. *Am. J. Pathol.* 11:1-72, 1935.
- Cope, O. The story of hyperparathyroidism of the Massachusetts General Hospital. *N. Engl. J. Med.* 274:1174-1182, 1966.
- Hall, E. M. and Chaffin, L. Malignant tumors of the parathyroid glands. *West. J. Surg. Obstet. Gynecol.* 42:578-586, 1934.
- Mandl, F. Therapeutischer Versuch bei einem Falle von Ostitis fibrosa generalisata mittels Exstirpation eines Epithelkörperchentumors. *Centralb. f. Chir.* 53:260-264, 1926.
- Zur Frage der Exstirpation eines Epithelkörperchentumors bei der allgemeinen Osteitis fibrosa. *Z. Klin.* 56:1739-1745, 1929.
- Owen, Richard. On the anatomy of the Indian rhinoceros (*Rh. unicornis*, L.). *Trans. Zool. Soc. Lond.* 4:31-58, 1862.
- Pappenheimer, A. M. and Wilens, S. L. Enlargement of the parathyroid glands in renal disease. *Am. J. Pathol.* 11:73-91, 1935.
- Sandström, I. Om en ny körtel hos menniskan och atskilliga däggdjur. *Upsala Läkareförenings Förh.* 15:441-554, 1879-1880. (Translated by Seipel, C. M. with biographical notes by Hammar, J. H. *Bull. Hist. Med.* 6:179-222, 1938.)
- Virchow, R. *Die krankhaften Geschwulste.* Berlin: A. Hirschwald, 1863.
- Von Recklinghausen, F. Die fibröse oder deformierende Ostitis, die Osteomalacie und die osteoplastische Carcinose in ihren gegenseitigen Beziehungen. In: *Festschrift Rudolf Virchow zu seinem 71. Geburtstage.* Berlin: G. Reimer, 1891.



NORMAL PARATHYROID GLAND

Knowledge of the embryologic development and the normal anatomic variations of the parathyroid gland is necessary to recognize and understand pathologic alterations.

EMBRYOLOGY

The human parathyroid glands are first recognizable in the 8 to 9 mm. embryo as bilateral localized proliferations arising from the third and fourth branchial pouches. When the third pouch separates from the pharynx, parathyroid III* forms part of a bilobed body with the thymus III, and, as the heart descends into the thorax, it pulls thymus III and parathyroid III caudally with it[†]. In the 18 mm. embryo, when parathyroid III is approximately at the level of the lower pole of the thyroid gland, it usually separates from the thymus gland and remains near the lower pole of the thyroid gland as the lower parathyroid gland. However, one or both parathyroid IIIs may remain attached to the thymus gland and come to lie in the lower neck or the anterior mediastinum as far caudad as the pericardium, or even in the posterior

.....
*Parathyroid III refers to the parathyroid glands derived from the third branchial pouch, and parathyroid IV to the glands derived from the fourth pouch. Unless otherwise noted, the glands from each pouch develop approximately symmetrically, and, therefore, the singular is used.

†Although embryologic usage refers to structures rising and descending, the relative alterations in location of structures are due in large part to differential growth rates of adjacent organs.

mediastinum. If parathyroid III separates prematurely from the thymus gland, its final position may even be cephalad to the thyroid gland and parathyroid IV. In recent studies of the locations of normal parathyroid glands, Wang found that the lower gland was on the anterior or posterolateral surface of the thyroid gland in 42 percent of the cases (fig. 1); 39 percent were in the lower neck underneath the capsule of the thymic tongue, 15 percent were in the neck lateral to the lower pole of the thyroid gland, 2 percent were in the mediastinal thymic tissue, and the remaining 2 percent were at the carotid bifurcation above the upper pole of the thyroid gland or lateral to the midportion of the gland.

Parathyroid IV forms a bilobed complex with the ultimobranchial or postbranchial body (lateral thyroid), the gland associated with the calcitonin-secreting C cells. As the ultimobranchial body joins the lateral lobes of the thyroid gland, parathyroid IV separates from it and finds its adult position as the upper gland, near the point where the medial thyroid artery crosses the recurrent laryngeal nerve at the cricothyroid junction posteriorly. Since parathyroid IV (upper gland) is more medial and closely associated with the midline structures than parathyroid III (lower gland), the former has more movement in a cephalad direction and is more constant in position, although its position may vary in the dorsoventral plane with respect to the nerve and artery. In Wang's series, 77 percent of the upper

glands were found at the cricothyroid junction (fig. 2), 22 percent were behind the upper pole of the thyroid gland, invariably underneath the surgical capsule of the thyroid (also known as the pretracheal

lamina of the cervical fascial sheath), and 1 percent were in the midline behind the pharyngoesophageal junction.

Rarely, either parathyroid gland may remain in the pharyngeal wall or move more

ANATOMIC DISTRIBUTION OF 312 LOWER PARATHYROID GLANDS PARATHYROID III




Figure 1

NORMAL PARATHYROID GLAND

Anatomic distribution of 312 normal lower parathyroid glands. (Fig. 2 from Wang, C-A. The anatomic basis of parathyroid surgery. *Ann. Surg.* 183:271-275, 1976.)

ventrally to lie in the esophagotracheal groove or behind the esophagus. The glands tend to be bilaterally symmetrical in location. Although the number of parathyroid glands has been reported to vary between

two and eight, in the most careful studies the finding of less than four glands in either the embryo or the adult is rare. Supernumerary glands, which have been reported in 2 to 6.5 percent of adults, probably




Figure 2
NORMAL PARATHYROID GLAND

Anatomic distribution of 312 normal upper parathyroid glands. (Fig. 1 from Wang, C-A. The anatomic basis of parathyroid surgery. *Ann. Surg.* 183:271-275, 1976.)




Figure 3

EMBRYOLOGY OF PARATHYROID GLAND

Cervical region of a 12 cm. embryo: a small fragment of the lower gland has been separated by a small vessel to form a supernumerary gland. Hematoxylin and eosin stain.* X10.

result from embryonic division of one or more glands (fig. 3). Parathyroid rests (parathyromatosis) resulting from deposits of parathyroid cells during embryonic migration could also account for some of the smaller supernumerary glands (Reddick et al.).

GROSS ANATOMY

From infancy and childhood, the growth of the parathyroid gland continues until the late third or early fourth decade, when the weight of the total parathyroid gland levels off at 120 ± 3.5 mg. in men and 142 ± 5.2 mg. in women (fig. 4). There

may be a somewhat uneven distribution of weight in the individual glands, and the lower gland is generally heavier than the upper. In adults, each gland is from 3 to 6 mm. in length, 2 to 4 mm. in width, and 0.5 to 2 mm. in thickness; it varies in shape as it is molded by adjacent tissues. The gland is soft and malleable and is covered by a thin, colorless capsule. It is yellow-tan to orange-tan, depending upon the amount of fat, the degree of vascular congestion, and the number of oxyphil cells.

MICROSCOPIC ANATOMY

The adult gland is composed of chief and oxyphil cells and a stroma largely consisting of fat cells (fig. 5). The chief cell is polyhedral and poorly outlined and 6 to 8

.....
*Throughout the fascicle where the stain is not so designated, hematoxylin and eosin stain has been used.




Figure 4

CHART DEMONSTRATING GROWTH OF PARATHYROID GLANDS

The difference in the total and parenchymal weight in older persons is due to increased stromal fat. (From data of Gilmour, J. R. and Martin, W. J. The weight of parathyroid glands. *J. Pathol. & Bact.* 44:431-462, 1937; also fig. 3 from Roth, S. I. Recent advances in parathyroid gland pathology. *Am. J. Med.* 50:612-622, 1971.)




Figure 5

NORMAL PARATHYROID GLAND

Cross section of an entire normal parathyroid gland, demonstrating the usual amount and distribution of stromal fat in a middle-aged person. X15.




Figure 6
NORMAL PARATHYROID GLAND

The sheets of chief cells in this normal parathyroid gland of a middle-aged person are separated by mature fat cells. The chief cells have distinctive cell membranes and small, centrally located nuclei. X400. (Fig. 2 from Fascicle 15, First Series.)

μm . in diameter (fig. 6). The cytoplasm is amphophilic to slightly eosinophilic. Artifactual vacuolization due to fixation is frequently seen. Early workers referred to the vacuolated, often glycogen-rich, chief cells in normal glands and adenomas as "clear cells" (see page 36). We suggest, to avoid confusion, that the terms "clear cell" or "water-clear cell" be restricted to those cells with the membrane limited vacuoles seen in clear-cell hyperplasia (see page 70). The nuclear chromatin is well defined and abundant, and the nuclear membrane is sharply outlined, giving the nucleus a pyknotic appearance. Intracellular fat is

present in 70 to 80 percent of the normal chief cells of a euparathyroid person (fig. 7).

Near the time of puberty, the oxyphil cells appear, at first singly and then in groups, presumably representing clones. An occasional oxyphil cell has been seen in childhood. They increase in number with advancing age, often forming large islands and nodules. After 60 years of age, they may occupy a large proportion of the gland, forming large single nodules (fig. 8) that may be incorrectly interpreted as neoplastic. The oxyphil cells are 8 to 12 μm . in diameter, with well demarcated cell mem-




Figure 7

NORMAL PARATHYROID GLAND

Normal parathyroid gland, with intracellular fat droplets in 70 to 80 percent of the chief cells. Sudan IV and toluidine blue stain. X300.



Figure 8

PARATHYROID GLAND

(Figures 8 and 9 from same case)

Cross section of an entire parathyroid gland from an elderly person, demonstrating a large single island of oxyphil cells and several smaller oxyphil cell nodules. X25. (Fig. 3 from Fascicle 15, First Series.)