



FAMOUS PROBLEMS  
OF  
ELEMENTARY GEOMETRY

THE DUPLICATION OF THE CUBE  
THE TRISECTION OF AN ANGLE  
THE QUADRATURE OF THE CIRCLE

AN AUTHORIZED TRANSLATION OF F. KLEIN'S  
VORTRÄGE ÜBER AUSGEWÄHLTE FRAGEN DER ELEMENTARGEOMETRIE  
AUSGEARBEITET VON F. TÄGERT

BY

WOOSTER WOODRUFF BEMAN

PROFESSOR OF MATHEMATICS IN THE UNIVERSITY OF MICHIGAN

AND

DAVID EUGENE SMITH

PROFESSOR OF MATHEMATICS IN TEACHERS COLLEGE, COLUMBIA UNIVERSITY



GINN AND COMPANY

BOSTON · NEW YORK · CHICAGO · LONDON  
ATLANTA · DALLAS · COLUMBUS · SAN FRANCISCO

## PREFACE.

---

THE more precise definitions and more rigorous methods of demonstration developed by modern mathematics are looked upon by the mass of gymnasium professors as abstruse and excessively abstract, and accordingly as of importance only for the small circle of specialists. With a view to counteracting this tendency it gave me pleasure to set forth last summer in a brief course of lectures before a larger audience than usual what modern science has to say regarding the possibility of elementary geometric constructions. Some time before, I had had occasion to present a sketch of these lectures in an Easter vacation course at Göttingen. The audience seemed to take great interest in them, and this impression has been confirmed by the experience of the summer semester. I venture therefore to present a short exposition of my lectures to the Association for the Advancement of the Teaching of Mathematics and the Natural Sciences, for the meeting to be held at Göttingen. This exposition has been prepared by Oberlehrer Tagert, of Ems, who attended the vacation course just mentioned. He also had at his disposal the lecture notes written out under my supervision by several of my summer semester students. I hope that this unpretending little book may contribute to promote the useful work of the association.

GÖTTINGEN, Easter, 1895.

F. KLEIN.

## TRANSLATORS' PREFACE.

---

At the Göttingen meeting of the German Association for the Advancement of the Teaching of Mathematics and the Natural Sciences, Professor Felix Klein presented a discussion of the three famous geometric problems of antiquity, — the duplication of the cube, the trisection of an angle, and the quadrature of the circle, as viewed in the light of modern research.

This was done with the avowed purpose of bringing the study of mathematics in the university into closer touch with the work of the gymnasium. That Professor Klein is likely to succeed in this effort is shown by the favorable reception accorded his lectures by the association, the uniform commendation of the educational journals, and the fact that translations into French and Italian have already appeared.

The treatment of the subject is elementary, not even a knowledge of the differential and integral calculus being required. Among the questions answered are such as these: Under what circumstances is a geometric construction possible? By what means can it be effected? What are transcendental numbers? How can we prove that  $e$  and  $\pi$  are transcendental?

With the belief that an English presentation of so important a work would appeal to many unable to read the original,

Professor Klein's consent to a translation was sought and readily secured.

In its preparation the authors have also made free use of the French translation by Professor J. Griess, of Algiers, following its modifications where it seemed advisable.

They desire further to thank Professor Ziwet for assistance in improving the translation and in reading the proof-sheets.

August, 1897.

W. W. BEMAN.

D. E. SMITH.

## ANNOUNCEMENTS



---

---

# TEXTBOOKS IN MATHEMATICS

FOR HIGH SCHOOLS AND ACADEMIES

- Baker: Elements of Solid Geometry  
Barker: Computing Tables and Mathematical Formulas  
Beman and Smith: Higher Arithmetic  
Betz and Webb: Plane Geometry  
Betz and Webb: Plane and Solid Geometry  
Betz and Webb: Solid Geometry  
Breckenridge, Mersereau, and Moore: Shop Problems in Mathematics  
Cobb: Elements of Applied Mathematics  
Hawkes, Luby, and Touton: Complete School Algebra  
Hawkes, Luby, and Touton: First Course in Algebra (Rev.)  
Hawkes, Luby, and Touton: Second Course in Algebra (Revised Edition)  
Hawkes, Luby, and Touton: Plane Geometry  
Moore and Miner: Practical Business Arithmetic (Revised Edition)  
Moore and Miner: Concise Business Arithmetic  
Morrison: Geometry Notebook  
Powers and Loker: Practical Exercises in Rapid Calculation  
Robbins: Algebra Reviews  
Schorling and Reeve: General Mathematics  
Smith: Algebra for Beginners  
Wentworth: Advanced Arithmetic  
Wentworth: New School Algebra  
Wentworth and Hill: First Steps in Geometry  
Wentworth-Smith Mathematical Series  
    Junior High School Mathematics  
    Higher Arithmetic  
    Academic Algebra  
    School Algebra, Books I and II  
    Vocational Algebra  
    Commercial Algebra  
    Plane and Solid Geometry. Also in separate editions  
    Plane Trigonometry  
    Plane and Spherical Trigonometry

---

---

## WENTWORTH-SMITH MATHEMATICS

THE Wentworth-Smith mark on a textbook is known to represent the highest attainment in the art of textbook making. The ability and experience of the authors combined with their knowledge of school needs is advance assurance of excellence.

ACADEMIC ALGEBRA. One book

SCHOOL ALGEBRA. Two books

A one-book or two-book course, constructed on modern lines, always avoiding extremes.

PLANE AND SOLID GEOMETRY. One-book or two-book course

The best geometry in twenty years, revised by the two leading mathematical textbook writers of the day. Marked for its simplicity of treatment, clearness of expression, and symmetry and legibility of page.

TRIGONOMETRY. In four editions

PLANE TRIGONOMETRY. With or without tables

PLANE AND SPHERICAL TRIGONOMETRY. With or without tables

TRIGONOMETRIC AND LOGARITHMIC TABLES

A direct and progressive treatment of trigonometry which shows practical uses wherever possible before the theoretical part of the subject is introduced.

COMMERCIAL ALGEBRA. Two books

This series by Wentworth, Smith, and Schlauch retains the conventional topics in algebra only in so far as they contribute to the commercial interests of the students. Training and problems are based on the commercial needs of today.

HIGHER ARITHMETIC

A textbook which may be used in any year of the high school, but some knowledge of algebra is desirable. It is especially valuable to prospective teachers and commercial students.

JUNIOR HIGH SCHOOL MATHEMATICS Three books

Arithmetic and, in addition, the principles and everyday applications of algebra, geometry, and trigonometry introduced in an informal way on the basis of intuitional concepts already acquired.



COPYRIGHT, 1897, BY  
WOOSTER WOODRUFF BEMAN AND DAVID EUGENE SMITH

---

ALL RIGHTS RESERVED

621.4

The Athenæum Press  
GINN AND COMPANY · PRO-  
PRIETORS · BOSTON · U.S.A.

# CONTENTS.

## INTRODUCTION.

	PAGE
PRACTICAL AND THEORETICAL CONSTRUCTIONS . . . . .	2
STATEMENT OF THE PROBLEM IN ALGEBRAIC FORM . . . . .	3

## PART I.

### The Possibility of the Construction of Algebraic Expressions.

#### CHAPTER I. ALGEBRAIC EQUATIONS SOLVABLE BY SQUARE ROOTS.

1-4. Structure of the expression $x$ to be constructed . . . . .	5
5, 6. Normal form of $x$ . . . . .	6
7, 8. Conjugate values . . . . .	7
9. The corresponding equation $F(x) = 0$ . . . . .	8
10. Other rational equations $f(x) = 0$ . . . . .	8
11, 12. The irreducible equation $\phi(x) = 0$ . . . . .	10
13, 14. The degree of the irreducible equation a power of 2 . . . . .	11

#### CHAPTER II. THE DELIAN PROBLEM AND THE TRISECTION OF THE ANGLE.

1. The impossibility of solving the Delian problem with straight edge and compasses . . . . .	13
2. The general equation $x^3 = \lambda$ . . . . .	13
3. The impossibility of trisecting an angle with straight edge and compasses . . . . .	14

#### CHAPTER III. THE DIVISION OF THE CIRCLE INTO EQUAL PARTS.

1. History of the problem . . . . .	16
2-4. Gauss's prime numbers . . . . .	17
5. The cyclotomic equation . . . . .	19
6. Gauss's Lemma . . . . .	19
7, 8. The irreducibility of the cyclotomic equation . . . . .	21

# CHAPTER IV. THE CONSTRUCTION OF THE REGULAR POLYGON OF 17 SIDES.

	PAGE
1. Algebraic statement of the problem . . . . .	24
2-4. The periods formed from the roots . . . . .	25
5, 6. The quadratic equations satisfied by the periods . . . . .	27
7. Historical account of constructions with straight edge and compasses . . . . .	32
8, 9. Von Staudt's construction of the regular polygon of 17 sides . . . . .	34

# CHAPTER V. GENERAL CONSIDERATIONS ON ALGEBRAIC CONSTRUCTIONS.

1. Paper folding . . . . .	42
2. The conic sections . . . . .	42
3. The Cissoïd of Diocles . . . . .	44
4. The Conchoid of Nicomedes . . . . .	45
5. Mechanical devices . . . . .	47

## PART II.

### Transcendental Numbers and the Quadrature of the Circle.

#### CHAPTER I. CANTOR'S DEMONSTRATION OF THE EXISTENCE OF TRANSCENDENTAL NUMBERS.

1. Definition of algebraic and of transcendental numbers . . . . .	49
2. Arrangement of algebraic numbers according to <i>height</i> . . . . .	50
3. Demonstration of the existence of transcendental numbers . . . . .	53

#### CHAPTER II. HISTORICAL SURVEY OF THE ATTEMPTS AT THE COMPUTATION AND CONSTRUCTION OF $\pi$ .

1. The empirical stage . . . . .	56
2. The Greek mathematicians . . . . .	56
3. Modern analysis from 1670 to 1770 . . . . .	58
4, 5. Revival of critical rigor since 1770 . . . . .	59

#### CHAPTER III. THE TRANSCENDENCE OF THE NUMBER $e$ .

1. Outline of the demonstration . . . . .	61
2. The symbol $h_r$ and the function $\phi(x)$ . . . . .	62
3. Hermite's Theorem . . . . .	65

CHAPTER IV. THE TRANSCENDENCE OF THE NUMBER  $\pi$ .

	PAGE
1. Outline of the demonstration . . . . .	68
2. The function $\psi(x)$ . . . . .	70
3. Lindemann's Theorem . . . . .	73
4. Lindemann's Corollary . . . . .	74
5. The transcendence of $\pi$ . . . . .	76
6. The transcendence of $y = e^x$ . . . . .	77
7. The transcendence of $y = \sin^{-1}x$ . . . . .	77

CHAPTER V. THE INTEGRAPH AND THE GEOMETRIC CONSTRUCTION  
OF  $\pi$ .

1. The impossibility of the quadrature of the circle with straight edge and compasses . . . . .	78
2. Principle of the integraph . . . . .	78
3. Geometric construction of $\pi$ . . . . .	79

## INTRODUCTION.

---

THIS course of lectures is due to the desire on my part to bring the study of mathematics in the university into closer touch with the needs of the secondary schools. Still it is not intended for beginners, since the matters under discussion are treated from a higher standpoint than that of the schools. On the other hand, it presupposes but little preliminary work, only the elements of analysis being required, as, for example, in the development of the exponential function into a series.

We propose to treat of geometrical constructions, and our object will not be so much to find the solution suited to each case as to determine the *possibility* or *impossibility* of a solution.

Three problems, the object of much research in ancient times, will prove to be of special interest. They are

1. *The problem of the duplication of the cube* (also called the *Delian problem*).
2. *The trisection of an arbitrary angle.*
3. *The quadrature of the circle, i.e., the construction of  $\pi$ .*

In all these problems the ancients sought in vain for a solution with straight edge and compasses, and the celebrity of these problems is due chiefly to the fact that their solution seemed to demand the use of appliances of a higher order. In fact, we propose to show that a solution by the use of straight edge and compasses is impossible.

The impossibility of the solution of the third problem was demonstrated only very recently. That of the first and second is implicitly involved in the Galois theory as presented to-day in treatises on higher algebra. On the other hand, we find no explicit demonstration in elementary form unless it be in Petersen's text-books, works which are also noteworthy in other respects.

At the outset we must insist upon the difference between *practical* and *theoretical* constructions. For example, if we need a divided circle as a measuring instrument, we construct it simply on trial. Theoretically, in earlier times, it was possible (*i.e.*, by the use of straight edge and compasses) only to divide the circle into a number of parts represented by  $2^n$ , 3, and 5, and their products. Gauss added other cases by showing the possibility of the division into parts where  $p$  is a prime number of the form  $p = 2^{2^m} + 1$ , and the impossibility for all other numbers. No practical advantage is derived from these results; *the significance of Gauss's developments is purely theoretical*. The same is true of all the discussions of the present course.

Our fundamental problem may be stated: *What geometrical constructions are, and what are not, theoretically possible?* To define sharply the meaning of the word "construction," we must designate the instruments which we propose to use in each case. We shall consider

1. Straight edge and compasses,
2. Compasses alone,
3. Straight edge alone,
4. Other instruments used in connection with straight edge and compasses.

The singular thing is that elementary geometry furnishes no answer to the question. We must fall back upon algebra and the higher analysis. The question then arises: How



shall we use the language of these sciences to express the employment of straight edge and compasses? This new method of attack is rendered necessary because elementary geometry possesses no general method, no *algorithm*, as do the last two sciences.

In analysis we have first *rational* operations: addition, subtraction, multiplication, and division. These operations can be directly effected geometrically upon two given segments by the aid of proportions, if, in the case of multiplication and division, we introduce an auxiliary unit-segment.

Further, there are *irrational* operations, subdivided into *algebraic* and *transcendental*. The simplest algebraic operations are the extraction of square and higher roots, and the solution of algebraic equations not solvable by radicals, such as those of the fifth and higher degrees. As we know how to construct  $\sqrt{ab}$ , rational operations in general, and irrational operations involving only square roots, can be constructed. On the other hand, every *individual* geometrical construction which can be reduced to the intersection of two straight lines, a straight line and a circle, or two circles, is equivalent to a rational operation or the extraction of a square root. In the higher irrational operations the construction is therefore impossible, *unless we can find a way of effecting it by the aid of square roots*. In all these constructions it is obvious that the number of operations must be limited.

We may therefore state the following fundamental theorem: *The necessary and sufficient condition that an analytic expression can be constructed with straight edge and compasses is that it can be derived from the known quantities by a finite number of rational operations and square roots.*

Accordingly, if we wish to show that a quantity cannot be constructed with straight edge and compasses, we must prove that the corresponding equation is not solvable by a finite number of square roots.

*A fortiori* the solution is impossible when the problem has *no* corresponding algebraic equation. An expression which satisfies no algebraic equation is called a transcendental number. This case occurs, as we shall show, with the number  $\pi$ .

## PART I.

### THE POSSIBILITY OF THE CONSTRUCTION OF ALGEBRAIC EXPRESSIONS.



#### CHAPTER I.

##### Algebraic Equations Solvable by Square Roots.

The following propositions taken from the theory of algebraic equations are probably known to the reader, yet to secure greater clearness of view we shall give brief demonstrations.

*If  $x$ , the quantity to be constructed, depends only upon rational expressions and square roots, it is a root of an irreducible equation  $f(x) = 0$ , whose degree is always a power of 2.*

1. To get a clear idea of the structure of the quantity  $x$ , suppose it, *e.g.*, of the form

$$x = \frac{\sqrt{a + \sqrt{c + ef}} + \sqrt{d + \sqrt{b}}}{\sqrt{a} + \sqrt{b}} + \frac{p + \sqrt{q}}{\sqrt{r}},$$

where  $a, b, c, d, e, f, p, q, r$  are rational expressions.

2. The number of radicals one over another occurring in any term of  $x$  is called the *order of the term*; the preceding expression contains terms of orders 0, 1, 2.

3. Let  $\mu$  designate the *maximum order*, so that no term can have more than  $\mu$  radicals one over another.