

Diagnosis & Treatment of Lead Poisoning

Papers by

J. Julian Chisolm, Jr., Kathryn R. Mahaffey (Six),
Arthur L. Aronson et al.

MSS Information Corporation

655 Madison Avenue, New York, N.Y. 10021

Library of Congress Cataloging in Publication Data

Main entry under title:

Diagnosis and treatment of lead poisoning.

Collection of articles previously published in various journals.

Includes index.

CONTENTS: Toxicity of lead: Chisolm, J. J., Jr., Increased lead absorption: toxicological considerations. Mahaffey (Six), K.R., Goyer, R. A., and Haseman, J. K. Dose-response to lead ingestion in rats fed low dietary calcium. [etc.]

1. Lead-poisoning — Addresses, essays, lectures.

I. Chisolm, J. Julian. II. Hafaffey, Kathryn R., 1943- III. Aronson, Arthur L.

[DNLM: 1. Lead poisoning — Diagnosis — Collected works. 2. Lead poisoning — Therapy — Collected works. QV292 D536]

RA1231.L4D5

615.9'25'68808

74-26898

ISBN 0-8422-7262-3

Copyright © 1976
MSS Information Corporation
All Rights Reserved

TABLE OF CONTENTS

Credits and Acknowledgments	5
Preface	7
Toxicity of Lead	9
Increased Lead Absorption: Toxicological Considerations Chisolm	10
Dose-Response to Lead Ingestion in Rats Fed Low Dietary Calcium . . . Mahaffey(Six), Goyer and Haseman	14
Experimental Enhancement of Lead Toxicity by Low Dietary Calcium . . . Mahaffey(Six) and Goyer	23
The Influence of Iron Deficiency on Tissue Content and Toxicity of Ingested Lead in the Rat. Mahaffey(Six) and Goyer	33
Lead Poisoning in Cattle and Horses following Long-Term Exposure to Lead Aronson	42
Detection of Lead Toxicity	45
Tests for Lead Poisoning Deutsch	46
Screening for Lead Poisoning Lin-Fu	48
δ -Aminolevulinic Acid Dehydrase as a Measure of Lead Exposure Hernberg, Nikkanen, Mellin and Lilius	50
The Unsuitability of Random Urinary Aminolevulinic Acid Samples as a Screening Test for Lead Poisoning Specter, Guinee and Davidow	56
Screening Techniques for Undue Lead Exposure in Children: Biological and Practical Considerations Chisolm	62
Biochemical Tests for the Appraisal of Exposure to Lead Stanković	69
Toxic Effects of Lead on Kidney and Gastrointestinal Systems.	75
Lead and Protein Content of Isolated Intranuclear Inclusion Bodies from Kidneys of Lead-Poisoned Rats. Goyer, May, Cates and Krigman	76

The Devonshire Colic	Waldron	90
Lead Concentration Found in Human		
Blood in Association with Lead Colic	Beritić	123
Lead Poisoning	Guinee	126
Toxic Effects of Lead Poisoning		
on the Central Nervous System		141
Motor Neurone Disease and		
Exposure to Lead	Campbell, Williams and Barltrop	142
Effects of Lead on Blood Serum Components		
and the Reticuloendothelial System		151
Aminoaciduria in Experimental		
Lead Poisoning	Goyer, Leonard, Bream and Irons	152
Prevention and Treatment		157
Childhood Lead Intoxication:		
Diagnosis, Management and Prevention	Chisolm	158
Treatment of Acute Lead Intoxication —		
Choice of Chelating Agents and Supportive		
Therapeutic Measures	Chisolm	185
Prevention of Lead Paint —		
Or Prevention of Pica?	Lourie	199
Ambulatory Treatment of Lead		
Poisoning: Report of 1,155 Cases	Sachs, Blanksma, Murray and O'Connell	202

Diagnosis & Treatment of Lead Poisoning

Papers by

**J. Julian Chisolm, Jr., Kathryn R. Mahaffey (Six),
Arthur L. Aronson et al.**

MSS Information Corporation

655 Madison Avenue, New York, N.Y. 10021

Library of Congress Cataloging in Publication Data

Main entry under title:

Diagnosis and treatment of lead poisoning.

Collection of articles previously published in various journals.

Includes index.

CONTENTS: Toxicity of lead: Chisolm, J. J., Jr., Increased lead absorption: toxicological considerations. Mahaffey (Six), K.R., Goyer, R. A., and Haseman, J. K. Dose-response to lead ingestion in rats fed low dietary calcium. [etc.]

1. Lead-poisoning — Addresses, essays, lectures.

I. Chisolm, J. Julian. II. Hafaffey, Kathryn R., 1943- III. Aronson, Arthur L.

[DNLM: 1. Lead poisoning — Diagnosis — Collected works. 2. Lead poisoning — Therapy — Collected works. QV292 D536]

RA1231.L4D5

615.9'25'68808

74-26898

ISBN 0-8422-7262-3

Copyright © 1976
MSS Information Corporation
All Rights Reserved

TABLE OF CONTENTS

Credits and Acknowledgments	5
Preface	7
Toxicity of Lead.....	9
Increased Lead Absorption:	
Toxicological Considerations Chisolm	10
Dose-Response to Lead Ingestion in Rats Fed Low Dietary Calcium Mahaffey(Six), Goyer and Haseman	14
Experimental Enhancement of Lead Toxicity by Low Dietary Calcium ... Mahaffey(Six) and Goyer	23
The Influence of Iron Deficiency on Tissue Content and Toxicity of Ingested Lead in the Rat..... Mahaffey(Six) and Goyer	33
Lead Poisoning in Cattle and Horses following Long-Term Exposure to Lead Aronson	42
Detection of Lead Toxicity.....	45
Tests for Lead Poisoning..... Deutsch	46
Screening for Lead Poisoning Lin-Fu	48
δ -Aminolevulinic Acid Dehydrase as a Measure of Lead Exposure Hernberg, Nikkanen, Mellin and Lilius	50
The Unsuitability of Random Urinary Aminolevulinic Acid Samples as a Screening Test for Lead Poisoning Specter, Guinee and Davidow	56
Screening Techniques for Undue Lead Exposure in Children: Biological and Practical Considerations Chisolm	62
Biochemical Tests for the Appraisal of Exposure to Lead Stanković	69
Toxic Effects of Lead on Kidney and Gastrointestinal Systems.....	75
Lead and Protein Content of Isolated Intranuclear Inclusion Bodies from Kidneys of Lead-Poisoned Rats..... Goyer, May, Cates and Krigman	76

The Devonshire Colic	Waldron	90
Lead Concentration Found in Human		
Blood in Association with Lead Colic	Beritić	123
Lead Poisoning	Guinee	126
Toxic Effects of Lead Poisoning		
on the Central Nervous System		141
Motor Neurone Disease and		
Exposure to Lead	Campbell, Williams and Barltrop	142
Effects of Lead on Blood Serum Components		
and the Reticuloendothelial System		151
Aminoaciduria in Experimental		
Lead Poisoning	Goyer, Leonard, Bream and Irons	152
Prevention and Treatment		157
Childhood Lead Intoxication:		
Diagnosis, Management and Prevention	Chisolm	158
Treatment of Acute Lead Intoxication —		
Choice of Chelating Agents and Supportive		
Therapeutic Measures	Chisolm	185
Prevention of Lead Paint —		
Or Prevention of Pica?	Lourie	199
Ambulatory Treatment of Lead		
Poisoning: Report of 1,155 Cases	Sachs, Blanksma, Murray and O'Connell	202

CREDITS AND ACKNOWLEDGEMENTS

Aronson, Arthur L., "Lead Poisoning in Cattle and Horses following Long-Term Exposure to Lead," *American Journal of Veterinary Research*, 1972, 33:627-629.

Beritić, T., "Lead Concentration Found in Human Blood in Association with Lead Colic," *Archives of Environmental Health*, 1971, 23:289-291.

Campbell, A.M.G.; E.R. Williams; and D. Barltrop, "Motor Neurone Disease and Exposure to Lead," *Journal of Neurology, Neurosurgery and Psychiatry*, 1970, 33:877-885.

Chisolm, J. Julian, Jr., "Increased Lead Absorption: Toxicological Considerations," *Pediatrics*, 1971, 48:349-352.

Chisolm, J. Julian, Jr., "Screening Techniques for Undue Lead Exposure in Children: Biological and Practical Considerations," *The Journal of Pediatrics*, 1971, 79:799-804.

Chisolm, J. Julian, Jr., "Childhood Lead Intoxication: Diagnosis, Management and Prevention," *Medical Times*, 1970, 98:92-106.

Chisolm, J. Julian, Jr., "Treatment of Acute Lead Intoxication — Choice of Chelating Agents and Supportive Therapeutic Measures," *Clinical Toxicology*, 1970, 3:527-540.

Deutsch, Robyn H., "Tests for Lead Poisoning," *New England Journal of Medicine*, 1971, 285:1541.

Goyer, R.A.; D.L. Leonard; P.R. Bream; and T.G. Irons, "Aminoaciduria in Experimental Lead Poisoning," *Proceedings of the Society for Experimental Medicine and Biology*, 1970, 135:767-771.

Goyer, Robert A.; Phillip May; Maxine M. Cates; and Martin R. Krigman, "Lead and Protein Content of Isolated Intranuclear Inclusion Bodies from Kidneys of Lead-Poisoned Rats," *Laboratory Investigation*, 1970, 22:245-251.

Guinee, Vincent F., "Lead Poisoning," *The American Journal of Medicine*, 1972, 52:283-288.

Hernberg, Sven; Jorma Nikkanen; Guy Mellin; and Helena Lilius, "δ-Aminolevulinic Acid Dehydrase as a Measure of Lead Exposure," *Archives of Environmental Health*, 1970, 21:140-145.

Lin-Fu, Jane S., "Screening for Lead Poisoning," *Pediatrics*, 1970, 45:720-721.

Lourie, Reginald S., "Prevention of Lead Paint — Or Prevention of Pica?," *Pediatrics*, 1971, 48:490-491.

Mahaffey(Six), Kathryn R.; and Robert A. Goyer, "Experimental Enhancement of Lead Toxicity by Low Dietary Calcium," *The Journal of Laboratory and Clinical Medicine*, 1970, 76:933-942.

Mahaffey(Six), Kathryn R.; and Robert A. Goyer, "The Influence of Iron Deficiency on Tissue Content and Toxicity of Ingested Lead in the Rat," *The Journal of Laboratory and Clinical Medicine*, 1972, 79:128-136.

Mahaffey(Six), Kathryn R.; Robert Goyer; and Joseph K. Haseman, "Dose-Response to Lead Ingestion in Rats Fed Low Dietary Calcium," *The Journal of Laboratory and Clinical Medicine*, 1973, 82:92-100.

Sachs, Henrietta K.; Lorry A. Blanksma; Edward F. Murray; and Morgan J. O'Connell, "Ambulatory Treatment of Lead Poisoning: Report of 1,155 Cases," *Pediatrics*, 1970, 46:389-396.

Specter, Michael J.; Vincent F. Guinee; and Bernard Davidow, "The Unsuitability of Random Urinary Delta Aminolevulinic Acid Samples as a Screening Test for Lead Poisoning," *The Journal of Pediatrics*, 1971, 79:799-804.

Stanković, Milos K., "Biochemical Tests for the Appraisal of Exposure to Lead," *Archives of Environmental Health*, 1971, 23:265-269.

Waldron, H.A., "The Devonshire Colic," *Journal of the History of Medicine and Allied Sciences*, 1970, 25:383-413.

PREFACE

Recently, much attention has been directed to the study of lead as an environmental poison. No longer believed to be a problem confined to children eating paint, public health officials have expressed alarm at the increasing contamination of air with products of lead-using industries and automobile emissions from leaded gasolines. It is estimated that more than 200,000 tons of lead per year are exhausted from automobiles in North America alone. Overt lead intoxication manifests itself in the central nervous system, kidneys and bone. Chronic, diffuse renal disease, ataxia, acute lead encephalopathy, and mental retardation are only some of the pathological results of lead intoxication.

The present volume includes studies on the diagnosis, management and prevention of lead intoxication. Research on toxicity levels of lead and on various screening techniques for lead poisoning is presented. In addition, toxic effects of lead on the kidney, gastrointestinal system, central nervous system, blood serum components and on the reticuloendothelial system are considered.

Companion volumes, *Epidemiology of Lead Poisoning* and *Lead Poisoning in Animal Models*, are also available.

Toxicity of Lead

INCREASED LEAD ABSORPTION: TOXICOLOGICAL CONSIDERATIONS

If physicians are to deal effectively with the biologic effects of ubiquitous chemical agents such as lead, a working knowledge of the pertinent time-dose-response relationships is essential. Lead is currently considered a nonessential trace element and is detectable in minute amounts in food and water and in the blood, tissues, and excreta of virtually all healthy persons. Apparently, the biologic effects of the usual daily intake from uncontaminated food and water (approximately 0.2 to 0.3 mg lead/day in adults) is negligible, so that this small "dose" of lead has no known adverse effect on the health of people, but as the chronic "dose" of lead increases, adverse responses of increasing severity become evident. While initial increments in "dose" may be associated with undesirable subclinical effects for which the body may compensate, at least temporarily, further increments in "dose" (and absorption) produce obvious alterations in function and clinical symptoms of illness. Still higher "doses" will be fatal. Thus, as may seem self-evident, a continuum of increasingly severe adverse responses are apparently associated with increasing doses of lead. The time factor adds yet another important dimension to the problem.

Elsewhere in this issue, a statement by the U.S. Public Health Service entitled "Medical Aspects of Childhood Lead Poisoning,"¹ sets forth rather detailed guidelines for the management of children with increased lead absorption, with emphasis on preservation of health and prevention of overt plumbism. An examination of some of the risk factors involved and of our scanty knowledge of the dose-response relationships to lead in children may help to place

these rather detailed recommendations in perspective.

In the U.S. Public Health Service statement, great emphasis is placed on blood lead levels, which, in adults, reflect increases in the level, of current and recent exposure to inorganic lead salts. In particular, the work of Kehoe² in which human adult volunteers were fed supplemental amounts of lead (0.3, 1.0, 2.0 and 3.0 mg lead/day) has shown that blood lead levels increase in proportion to the dose and that the higher doses are associated with a more rapid rate of rise. While the relationship between blood lead levels and current exposure cannot be precisely defined, exponential increases in "dose" appear to be reflected by arithmetic increases in blood lead concentration.^{3,4} Clinical observations in children are consistent with this observation. The "dose" of lead associated with the repetitive ingestion of a few lead-containing paint chips (or of putty) or lead-contaminated acidic juices may contain 10 to 500 times or more the quantity of lead found in normal diet⁵⁻⁷—a truly enormous dose by comparison. Review of the basic data from previously published cases⁸⁻¹⁰ indicates that blood lead levels in children with pica for leaded paint may rise from the 40 to 60 μg lead range* to well above the 100 μg lead range within a period of 1 to 2 months. How frequently this occurs is, of course, unknown, but clearly it can occur. Finally, the blood lead level does not appear to be in equilibrium with the total body lead burden, most of which is apparently rather tightly bound in bone; rather, the limited data available suggest that blood lead levels provide an index of the small, but "mobile" pool of lead situated primarily in the soft tissues. It would further appear that this is the fraction of the

Preparation of this article was supported, in part, under Grant EC-00201-18, National Institute for Occupational Safety and Health, Public Health Service, Department of Health, Education and Welfare.

* It is understood that blood lead levels are expressed in terms of μg lead/100 gm whole blood throughout.

total body lead burden responsible for the known acute toxic effects of lead.¹¹

Both in children^{10,12} and adults^{13,14} with lead poisoning, important interrelationships have been found between blood lead levels, "chelatable" lead, and quantitative daily outputs of δ -aminolevulinic acid (ALA) and coproporphyrin (UCP) in urine. Again, arithmetic increases in blood lead levels appear to be associated with exponential increases in these other parameters. Increased excretion of ALA and UCP indicates interference in biosynthesis of heme. The "chelatable" lead (i.e., response to a standardized parenteral dose of CaEDTA†) correlates most closely with the daily output of ALA and UCP in urine.^{11,12} The curvilinear nature of these relationships is perhaps best seen in Figure 1 of the report of Selander and Cramer.¹⁴ Preliminary data in this laboratory indicate that similar relationships are also found in children.

Clinical risk factors may also be defined in terms of accurately determined blood lead levels. Epidemiologic surveys in "normal" adults indicate that mean blood lead levels are approximately 20 μg lead/100 gm whole blood (range 5 to 40 μg lead/100 gm whole blood) in populations *without* undue exposure to lead.^{15,16} Persons in close daily occupational contact with motor vehicular exhausts in confined spaces may exhibit blood lead levels in the 40 to 50 μg lead range, but rarely higher.¹⁵ Comparable data are not available in young children, since most of the published data are derived from populations in which pica, exposure to old housing and to air, dust, and dirt-borne lead cannot be excluded. Review of the files of previously reported children^{5,8-10} show that blood lead exceeded 100 μg in 136 of 139 cases of acute symptomatic lead poisoning. In 98 fatal cases of acute lead encephalopathy reported to the Baltimore City Health Department, the range of blood lead levels at the time of acute illness was 138 to 750 ($\bar{m} = 330$) μg /100 gm blood. The range in 46 children considered asymptomatic was 55 to 300 μg lead/100 gm. While

levels above 100 μg lead do not correlate well with the presence, absence, or severity of symptoms, virtually all such patients may be expected to show significant metabolic and functional abnormalities. Clearly, levels > 80 μg lead/100 gm whole blood in children signify a risk to health that is unacceptable. The biologic significance of sustained blood lead in the 50 to 80 μg lead/100 gm range is unclear: variations in metabolic responses are noted, but the usually mild and nonspecific nature of minimal symptoms precludes precise clinical diagnosis. Whether such children may sustain subtle, but significant impairment of function particularly in the nervous system is not known. Nevertheless, such children clearly have a potentially hazardous increase in soft tissue lead and continued excessive intake at a high dose level can quickly cause severe illness. Prompt termination of their abnormal intake is the paramount consideration.

The recommendations of the U.S. Public Health Service statement for pediatric care vary according to the estimate of current risk provided by blood lead levels. Levels < 40 μg lead are assumed to indicate a negligible risk with normal daily dietary intake probably accounting for most of the intake. (These assumptions remain to be documented satisfactorily in young children.) Blood lead levels > 40 μg lead signify some additional source of intake and hence "undue exposure to lead," which may result from airborne or dust- and dirt-borne lead, especially in congested urban areas.¹⁷ If so, current estimates indicate that blood lead levels, at least in adults, are not likely to rise much above the 50 μg lead range. These sources do, however, serve to increase the background level upon which further intake must be superimposed. If, on the other hand, the child's undue exposure to lead results from pica for high-dose environmental sources such as leaded paint, putty (? lead in dirt) or the use of improperly lead-glazed earthenware pottery, a rapid rise in blood and tissue lead levels may be expected. Only serial determina-

† Edethamil calcium disodium.

tions will indicate the trend and separate these two groups of children when the initial level lies in the 40 to 50 μg lead range. Levels from 50 to 79 μg lead indicate an accelerating risk, an increasing probability of demonstrable metabolic impairment in heme synthesis, and the possibility of ill-defined symptoms of illness. Levels above 80 μg lead appear to present an unacceptable risk, particularly if long sustained. Chelation therapy is clearly indicated in this group, but would be of progressively decreasing benefit at lower levels of lead in the soft tissues. Because of the persistent and often clandestine nature of pica in some children, and the prevalence of deteriorated housing, serial testing of preschool age children in identified high risk areas would appear to be an essential part of regular health care. The trend of blood lead levels will help to determine the need for ancillary tests and indicated medical care in a given child.

As noted in the accompanying Letter to the Editor,¹⁸ the recommendations concerning ancillary tests are deemed by some as unrealistic in that they call for timed quantitative urine collections. This is a serious deficiency of the current "state of the art." Qualitative tests in random samples of urine for aminolevulinic acid or urinary coproporphyrin (ALA or UCP) are apparently inadequate in children, as they do not discriminate between normal and subclinical increased lead absorption. Since they become reliable indicators only as blood lead exceeds 80 to 100 μg lead, their usefulness is limited to children already at high risk. When reproducible simplified procedures adapted to capillary blood samples can be developed, the wider use of meaningful biochemical data, so often essential to good medical care, will be possible.

Perhaps the most serious limitation imposed upon good medical management of children (and adults) is the very limited availability to the general public of reliable lead analyses in blood and urine. A recent interlaboratory study¹⁹ revealed wide discrepancies in measurements from a number

of laboratories. Reliable methods, quality control techniques, competent analysts, and continuous experience are essential if accurate and reproducible results are to be obtained in any trace metal analysis. The "U.S. Public Health Service method"²⁰ has served in recent years as a suitable reference method for other dithizone techniques. Criteria for evaluating other and newer ones may be found in the analytical literature.^{21,22}

In conclusion, some may deem the U.S. Public Health Service statement too idealistic. It is, however, aimed at the preservation of health with guidelines based on the limited data currently available. Careful attention to dose-response relationships in studies in experimental animal systems²³ and in clinical investigation may, in the future, help to define more closely some of the thresholds for the various significant adverse toxicological effects of excessive absorption of lead. At the biochemical level, the main adverse effects appear to be in the areas of heme synthesis, cellular respiration, and membrane function. At the action level, it is clear that sustained pediatric follow-up during the preschool years, coupled with better environmental control techniques, especially for the high-dose types of exposure, are essential to minimize this particular hazard to the health of children.

REFERENCES

1. Medical aspects of childhood lead poisoning; Statement approved by surgeon general, U.S.P.H.S. PEDIATRICS, 48:464, 1971.
2. Kehoe, R. A.: The Harben lectures, 1960. The metabolism of lead in man in health and disease. J. Roy. Inst. Public Health, 24:81, 1961.
3. Goldsmith, J. R., and Hexter, A. C.: Respiratory exposure to lead: Epidemiological and experimental dose-response relationships. Science, 158:132, 1967.