


SECOND EDITION

# CYTOPATHOLOGY OF THE HEAD AND NECK ULTRASOUND GUIDED FNAC



GABRIJELA KOCJAN



WILEY Blackwell

# Cytopathology of the Head and Neck

## Ultrasound Guided FNAC

---

### Second Edition

**Author: Gabrijela Kocjan, MBBS, Spec Clin Cyt, FRCPath**

Senior Lecturer/Honorary Consultant,  
Department of Cellular Pathology  
University College London/University College Hospitals London

**Contributor: Simon Morley, MA, BM, BCh, MRCP, FRCR**

Consultant Radiologist  
Department of Imaging  
University College Hospitals London

**WILEY** Blackwell



This edition first published 2017 © 2017 by John Wiley & Sons Ltd  
First edition published in 2001 by Greenwich Medical Media Limited (London)

*Registered Office*

John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

*Editorial Offices*

9600 Garsington Road, Oxford, OX4 2DQ, UK  
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK  
111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at [www.wiley.com/wiley-blackwell](http://www.wiley.com/wiley-blackwell)

The right of Gabrijela Kocjan to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

*Library of Congress Cataloging-in-Publication Data*

Names: Kocjan, Gabrijela, author.

Title: Cytopathology of the head and neck : ultrasound guided FNAC / Gabrijela Kocjan.

Other titles: Clinical cytopathology of the head and neck.

Description: Second edition. | Chichester, West Sussex ; Hoboken, NJ : John Wiley & Sons Inc., 2016. | Preceded by Clinical cytopathology of the head and neck : a text and atlas / Gabrijela Kocjan. 2001. | Includes bibliographical references and index.

Identifiers: LCCN 2016049896 | ISBN 9781118076026 (cloth) | ISBN 9781118560846 (Adobe PDF) | ISBN 9781118560792 (epub)

Subjects: | MESH: Head and Neck Neoplasms--diagnosis | Head and Neck Neoplasms--pathology | Biopsy, Fine-Needle--methods | Cyst Fluid--cytology | Cysts--pathology | Atlases

Classification: LCC RC280.H4 | NLM WE 17 | DDC 616.99/491--dc23

LC record available at <https://lcn.loc.gov/2016049896>

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover image: courtesy of the author

Cover design: Wiley

Set in 9/11pt Minion by SPi Global, Pondicherry, India

Printed and bound in Singapore by Markono Print Media Pte Ltd

# Cytopathology of the Head and Neck

Ultrasound Guided FNAC

Second Edition

Edited by Dr. Nijim Kocjan, MD, PhD, FRCR, FRCR Path

Dr. Nijim Kocjan, MD, PhD, FRCR, FRCR Path



MAXIMUM EFFECT WITH MINIMUM INTERVENTION

To all those striving to optimize diagnostic process and realize efficiency in the health service

# Preface to the first edition

The past century has seen Cytopathology as a discipline, and Fine Needle Aspiration Cytology as a method of obtaining material, become established. From the pioneering work of Martin and Ellis, through the enthusiasm of Zajicek and his colleagues, to the perseverance of numerous cytopathologists throughout the world, this simple technique has become part of the routine diagnostic investigations. To this end, this book summarises recent experiences with the role Cytopathology is playing in current clinical practice, particularly in relation to Head and Neck. This work is based on the experience drawn from a large referral practice from ear, nose and throat, maxillofacial and general surgeons, endocrinologists, oncologists and others. From sceptical beginnings the FNAC service has grown and developed, to the extent that it is now accepted as a routine investigation and cytopathologists are considered as the best people to deliver the service. Mutual understanding and trust between clinical colleagues and cytopathologists has led to the further development of skills and a desire for further improvement in respective disciplines. Clinically oriented

work is a valuable source for the education and training of laboratory and junior pathology staff. To a cytopathologist, who meets new patients daily, this work is enormously satisfying. In the new millennium, our efforts should not be spent any more at proving the validity of FNAC. Instead, with the advances in new technologies, our aim should be the refinement of morphological diagnoses in order to match the existing or improve future treatment options.

My sincere thanks for producing this book go, firstly, to the patients who posed for the, sometimes, unflattering photographs; to the medical and technical colleagues in the Cytopathology Laboratory, University College Hospital London, to eminent clinical colleagues for their advice and support and to Mr Paratian for processing the photographs. Lastly, I would like to thank Tony and Arabella for putting up with my absences during the long gestation of this book.

**Gabrijela Kocjan**  
London, May 2000

# Preface to the second edition

The idea for the second edition of this book arose through realization that the working practices of Head and Neck (HN) diagnostic and clinical teams have changed dramatically in the last 15 years, not only in terms of organization of health service with its aims for provision of HN cancer care but also in their diagnostic input. The publication of high profile professional guidance documents highlighted the importance of specialist Multidisciplinary Teams (MDT) with the intention that these should bring together all the services and organizations to provide high quality care. Written protocols, that specify investigations for each type of presentation of possible HN cancer as well as specific guidelines for investigation and diagnosis of each form of HN cancer, have emerged. The aim of reducing cancer waiting times meant that Rapid Access (One stop) diagnostic clinics have become a requirement not only in the base hospital of the specialist multidisciplinary teams (MDT) but also in many District General Hospitals and that these clinics are required to provide same day diagnosis by Ultrasound and Fine Needle Aspiration Cytology (FNAC), tissue/cell sampling thus becoming an essential function of these clinics. Pathologists within the HN networks now have to ensure that conditions under which FNAC and rapid diagnosis clinics services are provided follow the professional guidelines and are also part of the local network guidelines.

The initial experience with One stop Clinics found widespread diagnostic difficulties including a high non-diagnostic rate highlighting the need for a particularly high level of expertise required to achieve a precise and reliable diagnosis in HN through the involvement of specialist radiologists and cytopathologists. To achieve high levels of diagnostic accuracy, special training and commitment to cytopathology are required in addition to histopathology. There is a need for recognition of the new skills expected of practicing pathologists and a comprehensive approach to cytopathology training, to include performing FNAC, with or without ultrasound guidance and interpreting them on site, as is the case with frozen section specimen training in histopathology. Ancillary techniques that have become available in the past 10 years are now a mainstream requirement for diagnosis and sometimes prognosis of various conditions and can be applied to FNAC material. Trainee pathologists specializing in cytopathology require a secondment to centres where on site evaluation and rapid access clinics are in place and where molecular techniques are available. This may require pathologists to be absent from routine work at their institution in order to learn new skills and adopt different ways of working.

Ultrasound (US) guidance has emerged as an essential adjunct to either FNAC or needle core biopsy, and its use is expected to increase. US combined with US guided FNAC can be recommended as a method for evaluating regional metastases in HN patients, for both those with and those without palpable lumps. US and, if necessary, FNAC should continue to be the investigation method of

first choice for HN lesions. US-guided FNAC sessions benefit from attendance of cytopathology medical and non-medical staff to perform the procedure, assess adequacy of the samples and make decisions about collecting appropriate material for ancillary tests.

In our own practice, the emergence of MDT Meetings (MDM) where radiologists, oncologists, radiotherapists, surgeons, speech and language therapists, pathologists and other support staff meet regularly once a week and discuss individual cases in a formal meeting, meant a significant improvement in HN service. MDMs contributed to the understanding of the role each discipline plays in the clinical management and helped improve patient outcomes. This collaboration in a quest for successful outcomes has also helped drive the progress in using ancillary techniques in diagnosis thus enabling the so called personalised medicine. One stop HN clinics and MDMs are a model of service delivery that hopefully can be used as an example in successful health management.

My thanks for the publication of this book go primarily to all patients whose conditions served as an inspiration for education, training and research. I am extremely appreciative of all members of the MDT for their input, patience and support; our sessions were as much fun as they were informative. Thanks to my colleagues Simon Morley, a contributor to this book, and Timothy Beale, both radiologists, I managed to obtain a desk and a chair in the One Stop Clinic. Collaboration with surgeons, in particular Paul O'Flynn and Francis Vaz, went beyond the HN to tennis and golf tournaments. My gratitude goes to all my colleagues and staff in the Department of Cellular Pathology, who skeptically tolerated my indulgence in cytology, provided there was a Summer Party at the end. It is through cells that I met so many wonderful people, travelled around the world and made lasting friendships. It is a testament to Wiley editorial and production teams, headed by Claire Bonnet and Eswari Maruthu that this book is presented in such a clear and constructive manner which I am proud of and thankful for. Finally, my lasting devotion goes to my family who were a source of pride and encouragement throughout. I hope that this textbook justifies the sacrifice they made.

As I am approaching the end of my working life, this book represents forty years of experience working as a diagnostic cytopathologist in a prestigious institution, a tertiary referral and a Cancer Centre. As such, it is a summary of the most interesting clinical examples where FNAC made a real difference to the management. It is my life long ambition that this legacy continues and that, by using cells alone, maximum diagnostic effect is achieved with minimum of intervention. I believe that this is achievable in not too distant future.

**Gabrijela Kocjan**  
Cavtat, January 2017



## About the companion website

This book is accompanied by a companion website:

**[www.wiley.com/go/kocjan/clinical\\_cytopathology\\_head\\_neck2e](http://www.wiley.com/go/kocjan/clinical_cytopathology_head_neck2e)**

The website includes:

- Over 20 exclusive to website studies of head and neck ultrasound case histories, with description of essential diagnostic features and differential diagnosis, compiled by Dr Simon Morley.
- Powerpoints of all figures from the book

The password for the site is the last word in the caption for Figure 4.1.

# Contents

Preface to the first edition, ix

Preface to the second edition, x

About the companion website, xi

## 1 Introduction, 1

1.1 Introduction, 1

1.2 Fine needle aspiration cytology of the head and neck, 1

1.3 Ultrasound guided FNAC, 1

1.4 A Combined US/FNAC approach, 2

1.5 Sampling technique, 2

References, 8

## 2 Salivary gland, 9

2.1 Introduction, 9

2.1.1 Ultrasound guided FNAC, 10

2.1.2 Diagnostic accuracy, 10

2.1.3 Diagnostic pitfalls, 11

2.1.4 Ultrasound versus other imaging modalities, 12

2.1.5 FNAC versus frozen section and core biopsy, 12

2.1.6 Cost effectiveness, 12

2.2 Diagnostic imaging of salivary glands, 13

2.2.1 Normal Ultrasound appearance of salivary gland, 13

2.2.2 Imaging Pitfalls and scanning issues, 13

2.3 Cytology of the salivary gland, 14

2.3.1 Normal salivary gland cytology, 14

2.3.2 Sialadenosis, 14

2.3.3 Salivary gland cysts, 17

2.3.4 Sialadenitis, 21

2.3.5 Lymphoid proliferations of the salivary gland, 23

2.4 Salivary gland tumours, 29

2.4.1 Pleomorphic adenoma, 29

2.4.2 Adenolymphoma (Warthin's tumour), 35

2.4.3 Basal cell adenoma, 37

2.4.4 Oncocytoma, 39

2.4.5 Rare benign tumours, 40

2.5 Malignant tumours of the salivary gland, 42

2.5.1 Acinic cell carcinoma, 42

2.5.2 Mucoepidermoid carcinoma, 44

2.5.3 Adenoid cystic carcinoma, 48

2.5.4 Polymorphous low grade adenocarcinoma, 51

2.5.5 Epithelial myoepithelial carcinoma, 52

2.5.6 Basal cell adenocarcinoma, 53

2.5.7 Papillary cystadenocarcinoma, 54

2.5.8 Mucinous adenocarcinoma, 54

2.5.9 Oncocytic carcinoma, 54

2.5.10 Salivary duct carcinoma, 55

2.5.11 Adenocarcinoma (not otherwise specified), 57

2.5.12 Carcinoma ex PLA, 57

2.5.13 Primary squamous cell carcinoma of the salivary gland, 58

2.5.14 Small cell carcinoma, 58

2.5.15 Undifferentiated carcinoma of the salivary gland, 59

2.6 Miscellaneous tumours, 59

2.6.1 Lymphoepithelial carcinoma, 59

2.6.2 Mammary analogue secretory carcinoma, 59

2.6.3 Cribriform adenocarcinoma of minor salivary gland, 60

2.6.4 Soft tissue lesions, 61

2.6.5 Granulocytic sarcoma, 61

2.6.6 Paediatric lesions, 61

2.6.7 Lymphomas, 61

2.6.8 Primitive neuroectodermal tumour, 62

2.6.9 Metastatic tumours in salivary gland, 63

2.7 Clinical management of salivary gland lesions, 64

References, 65

## 3 Thyroid, 71

3.1 Introduction, 71

3.1.1 Ultrasound and FNAC procedure, 72

3.1.2 FNAC reporting categories, 74

3.1.3 Diagnostic accuracy, 74

3.1.4 Diagnostic pitfalls, 76

3.1.5 The role of FNAC thyroid in clinical management, 76

3.1.6 FNAC versus frozen section and core biopsy histology, 77

3.1.7 Ancillary techniques, 78

3.1.8 Complications of FNAC, 78

3.2 Non-neoplastic and inflammatory conditions, 79

3.2.1 Colloid goitre (non-toxic goitre, adenomatous hyperplasia, multinodular goitre), 79

3.2.2 Cysts, 81

3.2.3 Hyperactive goitre (toxic goitre, thyrotoxicosis, primary hyperthyroidism, Graves' disease), 83

3.2.4 Thyroiditis, 85

3.3 Indeterminate cytological findings: follicular lesions, 87

3.3.1 Atypia of uncertain significance (AUS)/Follicular lesion of uncertain significance (FLUS) (TBSRTC III, UK Thy 3a), 87

3.3.2 'Follicular lesions' (TBSTRC IV: Follicular neoplasm or suspicious for a follicular neoplasm) (UK: Thy 3f), 90

3.4 Malignant tumours, 94

3.4.1 Papillary carcinoma, 94

3.4.2 Follicular carcinoma, 101

3.4.3 Medullary carcinoma, 101

3.4.4 Anaplastic carcinoma, 103

- 3.4.5 Thyroid lymphoma, 103
- 3.4.6 Metastatic tumours, 105
- References, 106
- 4 Lymph nodes, 112**
  - 4.1 Introduction, 112
    - 4.1.1 Distribution of lymph node pathology, 112
    - 4.1.2 Diagnostic accuracy, 113
    - 4.1.3 Diagnostic pitfalls, 113
    - 4.1.4 Ancillary techniques, 113
    - 4.1.5 Core biopsy or FNAC?, 117
    - 4.1.6 Elastography or FNAC?, 117
  - 4.2 Non-neoplastic lymphoproliferative conditions, 118
    - 4.2.1 Follicular hyperplasia, 118
    - 4.2.2 Sinus histiocytosis with massive lymphadenopathy (Rosai Dorfman), 119
    - 4.2.3 Granulomatous lymphadenitis, 122
    - 4.2.4 Chronic lymphadenitis, 128
    - 4.2.5 Drug reactions, 131
    - 4.2.6 Miscellaneous lymphadenopathies, 131
  - 4.3 Hodgkin's lymphoma, 137
    - 4.3.1 Nodular Lymphocyte Predominant Hodgkin's lymphoma, 139
  - 4.4 Non-Hodgkin's lymphoma, 139
    - 4.4.1 Introduction, 139
    - 4.4.2 Obtaining appropriate material, 142
    - 4.4.3 Classification of Non-Hodgkin's lymphoma, 142
    - 4.4.4 Precursor lesions, 142
    - 4.4.5 B-cell lymphomas, 143
    - 4.4.6 Mantle zone lymphoma, 144
    - 4.4.7 Follicular lymphoma, 147
    - 4.4.8 Marginal zone lymphoma (MALT type), 148
    - 4.4.9 Diffuse large B-cell lymphoma, 150
    - 4.4.10 Primary effusion lymphoma, 151
    - 4.4.11 Burkitt's lymphoma, 152
    - 4.4.12 T/NK-cell lymphomas, 154
    - 4.4.13 Anaplastic large cell lymphoma, 154
  - 4.5 Metastatic carcinoma in lymph nodes, 160
  - References, 170
- 5 Miscellaneous lesions of the head and neck, 176**
  - 5.1 Introduction, 176
  - 5.2 Benign soft tissue lesions, 176
    - 5.2.1 Lipoma, 176
    - 5.2.2 Fibromatosis colli, 176
    - 5.2.3 Nodular fasciitis, 178
    - 5.2.4 Proliferative fasciitis and proliferative myositis, 178
    - 5.2.5 Benign nerve sheath tumour (neurilemmoma, schwannoma), 180
  - 5.3 Cysts of the head and neck, 181
    - 5.3.1 Thyroglossal cysts, 181
    - 5.3.2 Branchial cyst, 181
    - 5.3.3 Mucous retention cyst, 182
    - 5.3.4 Intraosseous cysts and other lesions, 183
    - 5.3.5 Rare cysts and differential diagnosis of cystic lesions of the head and neck, 183
  - 5.4 Small round cell tumours, 184
    - 5.4.1 Rhabdomyosarcoma, 185
    - 5.4.2 Ewing's sarcoma/peripheral neuroectodermal tumour/Askin tumour, 186
    - 5.4.3 Olfactory neuroblastoma, 187
    - 5.4.4 Lymphoma, 187
  - 5.5 Locally arising miscellaneous tumours, 189
    - 5.5.1 Carotid body tumours, 189
    - 5.5.2 Epithelioid sarcoma-like hemangioendothelioma, 190
    - 5.5.3 Meningioma, 190
    - 5.5.4 Ethmoid sinus intestinal type adenocarcinoma, 191
    - 5.5.5 Granular cell tumour, 192
  - References, 195
- Index, 198



## Chapter contents

- |   |                                    |
|---|------------------------------------|
| 1.1 Introduction, 1   | 1.4 A combined US/FNAC approach, 2 |
| 1.2 Fine needle aspiration cytology of the head and neck, 1 | 1.5 Sampling technique, 2          |
| 1.3 Ultrasound guided FNAC, 1                               | References, 8                      |

## 1.1 Introduction

The Head and Neck (HN) area is one of the most complex regions of the body because of its anatomical and functional diversity. Diseases of the HN, both primary and systemic, rarely go unnoticed; patients either notice changes themselves, or are alerted to them by the diagnostic investigations, often done for unrelated conditions.

HN cancer is the ninth most common cancer in the USA, accounting for 3.3 % of all cancers. The incidence of HN cancer has plateaued recently; however, morbidity and mortality continue to remain high. Despite the decline in overall mortality rates since 2001 a racial disparity between the whites and the African Americans, both in incidence and mortality, still exists [1].

Tobacco and alcohol use are the most important risk factors for most HN cancers. In addition, infection with certain types of human papillomavirus (HPV) is thought to be the cause of an escalating incidence of HPV-related oropharyngeal squamous cell carcinoma predominantly among middle-aged adults [2].

## 1.2 Fine needle aspiration cytology of the head and neck

Fine needle aspiration cytology (FNAC) has been recognised as one of the core activities for the management of HN disease [3–23]. Sites in the HN that are amenable to FNAC include the thyroid, cervical masses and nodules, salivary glands, intraoral lesions and lesions in the paraspinal area and base of skull [24].

FNAC has a high overall diagnostic accuracy: 85–95% for all HN masses, 95% for benign lesions, and 87% for malignant ones [25, 26]. Diagnostic accuracy is dependent on the site of aspiration as well as the skill of the individual performing and interpreting the FNAC [24]. Each site undergoing FNAC within the HN is associated with its own set of differential diagnoses and diagnostic challenges. There are virtually no contraindications, and complications are minimal [27].

FNAC allows an immediate diagnosis to be available to the clinician so that appropriate treatment can be discussed with the patient. It is recommended as a first line of investigation in palpable HN masses. FNAC is the preferred first-line pathological

investigation of salivary gland and thyroid lumps because of the risk of recurrence and complications, respectively, associated with tissue biopsies [28].

The majority of aspirates from the HN will be to confirm an otherwise suspected diagnosis, for example a reactive lymphadenopathy or to confirm clinical staging for a metastatic carcinoma. However, there are a number of occasions where an unsuspected condition may be revealed, such as lymphoma or a salivary gland tumour. Whilst the diagnosis of lymphoma may need further tissue work up, the diagnosis of salivary gland lesions is often definitive in that it guides the surgical or non-surgical management. FNAC can diagnose majority of thyroid enlargements and help reduce the rate of surgery for benign thyroid disease. Ancillary techniques, namely immunocytochemistry, flow cytometry and molecular techniques, can greatly broaden the diagnostic range and specificity of FNAC. They are particularly useful in the diagnosis of lymphoproliferative processes and in determining the precise nature of lesions as variable as rhabdomyosarcoma, olfactory neuroblastoma and granular cell tumour. The prudent use of these techniques can be cost-effective and avoid the need for more invasive diagnostic procedures [29].

## 1.3 Ultrasound guided FNAC

Ultrasound imaging is a dynamic and readily available technique that is particularly useful in the examination of superficial structures. Modern machines combined with high frequency linear probes (7.5–12 MHz) produce high definition images in multiple planes. The spatial resolution that is achieved surpasses that of both multislice computed tomography (CT) and magnetic resonance imaging (MRI). Images are rapidly acquired, artefacts are few, and the technique is highly acceptable to most patients. As an adjunct to structural imaging, colour (directional) and power Doppler (non-directional but more sensitive) are often used to assess blood flow and the vascularity of tissue. These techniques add value in detecting abnormal peripheral or chaotic flow patterns in malignant lymph nodes, in assessing the patency of normal vessels, and in the investigation of vascular and lymphatic malformations.



Ultrasound (US) guidance is a useful adjunct to either FNAC or needle core biopsy (CB), and its use is expected to increase. US combined with US guided FNAC, rather than a tissue biopsy, can be recommended as a method for evaluating possible regional metastases in HN cancer patients, for both those with and those without a known primary tumour [30, 31].

US and, if necessary, FNAC, should continue to be the investigation method of first choice for HN lesions. The main indication for CB is after repeated failures of FNAC to provide a diagnosis. It can also be performed in patients who are not surgical candidates or in those who refuse surgery. Kraft et al. found CB was superior to FNAC in providing a specific diagnosis (90 vs 66%), and achieved a higher accuracy in identifying true neoplasms (100 vs 93%) and detecting malignancy (99 vs 90%). However, the sensitivity and specificity did not differ significantly between the two methods [32]. Khalid et al. found that the use of US-guided FNAC as the initial modality for tissue sampling of a thyroid nodule is more effective than traditional FNAC at an additional cost of \$289 per additional correct diagnosis [33].

In our own experience, the adequacy rate of US guided FNAC is critical for the success of the service. In our institution the adequacy of US guided FNAC in the HN clinic is 97% [34]. Computed tomography and magnetic resonance imaging do not appear to add any advantage to FNAC in terms of specificity, sensitivity or accuracy of a malignant diagnosis [34]. As with rapid-diagnosis clinics, US-guided FNAC sessions benefit from attendance of cytopathology medical and non-medical staff to assess adequacy of the samples and make decisions about collecting appropriate material for ancillary tests.

#### 1.4 A Combined US/FNAC approach


Recently, some cytopathologists have learned to use ultrasound machines to assist them in performing FNAC procedures. The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and accuracy were 96, 50, 98, 33 and 94% in palpation guided (PGFNAC) versus 100, 86, 97, 100 and 97% in the US guided (USGFNAC) group, respectively. USGFNACs performed by a cytopathologist could significantly improve the specificity and NPV ( $P=0.04$ ) while preserving virtually the same excellent sensitivity and PPV as those of PGFNACs. With US guidance, a cytopathologist is able to perform FNACs in smaller, non-palpable lesions and target complex lesions with confidence and accuracy, thus achieving a better outcome [36].

In our experience, even better results are obtained when an experienced radiologist takes the FNAC and has a cytopathologist close by to process the material and interpret the slides (Fig. 1.1).

According to the British Society for Clinical Cytology (BSCC) Code of Practice, the combination of physical examination/clinical history, radiological assessment, careful needle sampling, appropriate cell preparation, subsequent interpretation and multidisciplinary clinical discussion are essential for a successful outcome [37]. The lack of skill, clinical information and communication can be detrimental to the result.

#### 1.5 Sampling technique

Sample collection is a major factor influencing both the adequacy and the accuracy of FNAC [38,39]. It is our experience and experience of others that a good sampling technique is essential for



(A)



(B)


**Figure 1.1** Ultrasound guided FNAC. (A) The radiologist, Dr Morley, uses the ultrasound probe in the left hand and injects the anesthetic into the lesion with the right hand. (B) Observing the monitor and using the ultrasound probe for guidance, the aspirator uses negative pressure to extract cystic fluid from the parotid gland lesion.

successful interpretation of FNAC. Comparing the material obtained by the cytopathologist with the material sent from various aspirators, Wu et al. found that the sensitivity of HN FNAC procedures is significantly better in the cytopathologist-performed group than in the non-cytopathologist-performed group (96 versus 67%) [40, 41]. Greater experience of the operator appears to improve the accuracy rate [42–44]. In experienced hands, palpation-guided FNAC is an excellent diagnostic tool. However, there is a movement towards using imaging guidance to target all masses [45].

The best results are obtained with a cytopathologist-led FNAC service, where the pathologist reviews the specimen immediately, in relation to the clinical context, thereby deciding on adequacy and the need for further sampling (Fig. 1.2) [46, 47].

With the FNAC procedure having been explained (Fig. 1.3), the patient is put in a supine position. The choice of whether to apply anaesthetic or not largely depends on the patient, the site involved and the extent of FNAC sampling planned. Since the average FNAC does not involve more than one pass with a 22+ G needle, most patients do not require local anaesthetic. However, if the patient is needle-phobic or a child, or if the site is particularly tender, for example, lip, nose, areola, or if it is expected that several passes will be necessary, a local anaesthetic is applied in the form of subcutaneous injection of 0.5 ml of 2% lignocaine. More recently we have






**Figure 1.2** Examination of the glass slides in the clinic. This gives an orientation of adequacy and indicates whether further samples need to be taken for special techniques and/or for microbiology cultures. Results are usually not discussed with the patient at this preliminary stage of the investigations.



**Figure 1.3** Clinical history and examination. The pathologist at the bedside examines the area referred to by the specialist and also asks the patient further relevant questions about the duration of the swelling, level of pain and any other associated systemic symptoms.

been using a needle free syringe where the pressurised air expels the anaesthetic, penetrating the skin, without the needle (Fig. 1.4A) [48]. Anaesthetic forms a small white ring through which the subsequent test needle is applied, once or more (Fig. 1.4B). Patients do not experience any pain on application of the anaesthetic and experience no or minimal pain at FNAC.



(A)




(B)


**Figure 1.4** (A) Needle free anaesthetic system. Designed for patients who need daily injections, e.g. insulin, but also applicable to the local anaesthesia. This is particularly useful in children, in needle-phobic patients, in sensitive sites and where multiple needle passes are anticipated. (B) Needle-free anaesthetic system. A pale ring indicates the area of subdermal infiltration. A test needle should be passed through this area.

The palpable area in question is cleaned with an antiseptic agent and fixed between the two fingers of the non-dominant hand. 22 G, 23G or smaller needle is then passed into the lump using a non-aspiration technique (capillary sampling) with the aid of a needle only (without the syringe attachment) (Fig. 1.5). In a meta analysis comprising over 2000 thyroid FNAC samples, there was no difference between the aspiration and non aspiration technique in assessing thyroid nodules [29]. In cases where a fluid aspirate is expected, a syringe and a syringe holder are attached to the needle to help aspiration (Fig. 1.6) [29, 49–54]. The needle is passed round in a fan-shaped manner several times in the cases of non-thyroid lumps. In the case of thyroid, several vertical movements in the same direction are usually sufficient to gain representative material. When exiting the lump, if using syringe attachment, it is important






**Figure 1.5** Free needle FNAC procedure. This, so called 'capillary technique', is particularly useful in very small, mobile lesions, e.g. lymph nodes. An aspirator has a much better feel of the tip of the needle and better control of the area sampled. It is not the method of choice for cystic or very sclerosed lesions.



**Figure 1.6** FNAC of fluids. A 20 ml syringe attached to a CAMECO (BELPRO MEDICAL, Canada), syringe holder A 23 G needle is used.

to release the negative pressure before exiting, otherwise the material is aspirated into the syringe and can only be retrieved by the aid of a needle wash.

Whilst adhering to the traditional technique of smearing the material ejected from the needle onto a slide and then either air drying or fixing it in alcohol, if necessary we also suspend the material from a separate needle pass in a liquid medium that can then be used for ancillary techniques including cell block



**Figure 1.7** Rapid staining and examination. Slides are stained with one of the rapid stains and examined under the microscope for cellularity. This gives a good orientation if more material is needed or if different cell preparation technique should be applied.

(CBL). Air dried smears can be stained by a rapid staining technique to assess material adequacy in the One Stop clinic (Fig. 1.7). CBL provide a method for immunocytochemistry (ICC) that has revolutionised cytopathology by making it possible to apply panels of antibodies to multiple sequential sections of aspirated or exfoliated cellular material [55]. CBL can be prepared from virtually all varieties of cytological samples. CBL sections offer advantages over conventional cytological smears with respect to cellular architecture and archival storage. They also provide several sections, which can be utilised to perform special stains, immunophenotypic analysis, ultrastructural studies and molecular tests, including cytogenetic and polymerase chain reaction (PCR)-based techniques [56–59]. In today's era of personalised medicine, the ability to perform these tests augment the utility of cytological samples in analysing the molecular alterations as effectively as surgical biopsies or resection specimens. With the availability of molecular targeted therapy for many cancers, a large number of recent studies have used cytological material or CBL for molecular characterisation. Jain et al. described various methods of preparations of CBL and their application in cytology. The advantages and disadvantages of various methods of cell preparation are outlined in Table 1.1 [50].

One of the easiest way to prepare a CBL is the so-called 'Poor Man's cell block'. This method should be available to any laboratory and is able to produce very good results (Fig. 1.8) [61].

The final cytopathology report should be clear, written with the knowledge of the ultrasound and clinical findings, morphology and ancillary techniques (Fig. 1.9). Difficult cases should be discussed at the intradepartmental and multidisciplinary meetings and are also an important source of education and training [62] (Fig. 1.10).



**Table 1.1** Comparison of different cytological preparation methods.

Direct smear <sup>13</sup>			Cytospin <sup>25</sup>		Cell block <sup>2-4</sup>		Liquid-based cytology <sup>26</sup>	
For	Against		For	Against	For	Against	For	Against
Diagnosis Fast			Optimal for cysts, urine and effusions	Cellular crowding	Additional to other slides	Needs time and histology skill	Easy to transport, collect and process	Low cellularity
Inexpensive	Needle handling		Multiple slides	Limited cellularity	Increased diagnostic yield		Reduced screening area	Cell shrinkage (methanol)
Routine process	Multiple slides		Air dried and/or alcohol fixed		More architecture		No air-drying artefacts	Less architecture
Permits rapid evaluation	Obscuring background				Archival storage		Clean background	Reduced or altered background material
Excellent cellular detail	Air-drying artefacts						Monolayer of cells	
ICC possible on fresh or destained slides	Adverse effect of stripped nuclei and cytoplasmic background		Routine laboratory method	Risk of FNs due to focal antigen expression	Easy to perform and compare with histology	Variable staining due to different fixatives	Equal or stronger than direct smears	Alcohol-based: may differ from formalin-fixed slides
Suitable for nuclear antibodies	Potential increased FNs				Routine histology controls Dual ICC possible		Equal distribution of staining	All antibodies not yet evaluated
							Clean background	
Cytogenetic and molecular testing Potentially effective for preparation and storage	May be diluted by normal cells		Suitable for FISH	Cell crowding may hamper nuclear signals	Optimal results with current methods	Depends on cellular yield and fixation methods	Suitable for FISH and molecular analysis	Limited studies so far available
Routinely available and cost-effective	Cells of interest may be enriched by microdissection		DNA extraction possible	May be diluted by normal cells	Cells of interest may be enriched by microdissection	Partial cells in sections	Avoids cross-linkage of formalin fixation	May be affected by alcohol fixation (e.g. PR)
Whole cells			Whole cells			Sections	Whole cells	

FISH, fluorescence in situ hybridization; FN, false negative; ICC, immunocytochemistry; PR, progesterone receptors.

Source: Jain D, Mathur SR, Iyer VK. Cell blocks in cytopathology: a review of preparative methods, utility in diagnosis and role in ancillary studies. *Cytopathology*. 2014 Dec; 25(6): 356–71.



# Cytology Report

CPA Accredited Laboratory

Department of Histopathology

University College London Medical School

Rockefeller Building, University St, London WC1E 6JJ

Lab No.: NG00-0404

Hospital No.: 0000000

Name : XXXXX,YYYYYY

Age/Sex : 18Y M

Cons/GP : Doctor X

Ward : Maxillofacial Unit,

SPECIMEN : Neck Fine needle aspirate

EXAMINED BY: DR T HATTER

SPECIMEN OBTAINED BY: DR G KOCJAN

## CLINICAL DATA:


Swelling left neck and post auricular region.  
Fluctuant below angle mandible.

## MATERIAL RECEIVED:

Nineteen slides from FNA 2 different sites in the  
left neck (see illustration).

## MICROSCOPIC DESCRIPTION:

Smears show numerous single and small  
aggregates of cells. The cells have large oval  
nuclei with fine chromatin pattern, indistinct  
nucleoli, and scanty basophilic cytoplasm with  
prominent vacuolation.



Immunocytochemistry: LCA and CD20 positive. Tdt, desmin, SMA, myoglobin, MYC2 negative.

## CYTOLOGICAL DIAGNOSIS:

FNA left neck. High grade B cell non Hodgkin's lymphoma (? Burkitt's). Biopsy is advised.

SNOMED : TY0600;M095903

Signed :

Pathologist : DR G KOCJAN


Date : 14/02/2000



PATHOLOGY DIRECTORATE

**Figure 1.8** Cytopathology report should contain patient's details, clinical history, number and position of the site(s) sampled (preferably illustrated by a photograph or a diagram), indicate the name of the aspirator, date of sample, number of slides made and/or other material obtained, microscopic description, special techniques used, cytological diagnosis, diagnostic code, pathologist's signature and date. This report is just an example for illustration. Today we would use flow cytometry and cell block immunocytochemistry to confirm and specify the diagnosis of lymphoma on FNAC.





**Figure 1.9** Images demonstrating the main steps in the preparation of a vapour fixed cell block. (A) The FNAC material from the thyroid is stained with May Grunwald Giemsa (MGG) and shows undifferentiated tumour cells. (B) The material is expelled from the needle into the well of the lid in a universal container. (C) It is left inverted for at least 6 h to allow the formalin vapours to fix it until solid after which it is handled as a histological sample. (D) afp negative, (E) calcitonin negative, (F) CD45 negative, (G) MNF 116 negative, (H) S100 negative and (I) Thyroglobulin positive. The conclusion was that this is anaplastic carcinoma arising from the thyroid.



**Figure 1.10** Cytology education and training. Most interesting cases are discussed weekly at the multiheaded microscope; once a year, a traditional Christmas quiz, in addition to the seasonal jollity, provides a reminder of the most difficult cases.