

Encyclopaedia of Geomechanics in Soil, Rock, and Environmental Engineering

Contributors | **Evgenii Sharkov, Wisley Moreira Farias, Tsuyoshi Ishikawa, and
Hayder Mohammed Salim Al-Maamori et al.**



Encyclopaedia of GEOMECHANICS IN SOIL, ROCK, AND ENVIRONMENTAL ENGINEERING

Volume I: Environmental Geomechanics

Contributors

**Evgenii Sharkov, Wisley Moreira Farias, Tsuyoshi Ishikawa, and Hayder
Mohammed Salim Al-Maamori et al.**



www.aurisreference.com

Encyclopaedia of Geomechanics in Soil, Rock, and Environmental Engineering

Volume I: Environmental Geomechanics

Contributors: Evgenii Sharkov, Wisley Moreira Farias, Tsuyoshi Ishikawa, and Hayder Mohammed Salim Al-Maamori et al.

Published by Auris Reference Limited

www.aurisreference.com

United Kingdom

Copyright 2016

The information in this book has been obtained from highly regarded resources. The copyrights for individual articles remain with the authors, as indicated. All chapters are distributed under the terms of the Creative Commons Attribution License, which permit unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Notice

Contributors, whose names have been given on the book cover, are not associated with the Publisher. The editors and the Publisher have attempted to trace the copyright holders of all material reproduced in this publication and apologise to copyright holders if permission has not been obtained. If any copyright holder has not been acknowledged, please write to us so we may rectify.

Reasonable efforts have been made to publish reliable data. The views articulated in the chapters are those of the individual contributors, and not necessarily those of the editors or the Publisher. Editors and/or the Publisher are not responsible for the accuracy of the information in the published chapters or consequences from their use. The Publisher accepts no responsibility for any damage or grievance to individual(s) or property arising out of the use of any material(s), instruction(s), methods or thoughts in the book.

Environmental Geomechanics

ISBN: 978-1-78154-835-6 (Set of 4 Volumes)

British Library Cataloguing in Publication Data

A CIP record for this book is available from the British Library

Printed in the United Kingdom

List of Abbreviations

ACF	Auto-Correlation Function
AWCP	Acoustic Water Column Profilers
CBM	Coal Bed Methane
CFE	Comisión Federal de Electricidad
CRP	Conservation Reserve Program
DEM	Digital Elevation Model
DITF	Drilling-Induced Tensile Fracture
DNRA	Dissimilatory Nitrate Reduction to Ammonium
ECD	Electron Capture Detector
FEM	Finite Element Method
GAM	Generalized Additive Model
GAMM	Generalized Additive Mixed Model
GLM	Generalized Linear Model
GLMM	Generalized Linear Mixed Model
LOT	Leak-Off Tests
LPB	La Popa Basin
MC	Multilayered Crust
MIMS	Membrane Inlet Mass Spectrometry
NOR	Nitric Oxide Reduction
PAL	Passive Aquatic Listener
REV	Representative Elementary Volume
SDR	Seaward-Dipping Reflector
SRV	Stimulated Reservoir Volume
TEB	Trans-Eurasian Belt

List of Contributors

E.V. Sharkov

Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry RAS, Moscow, Russia

V.B. Svalova

Institute of Geoecology RAS, Moscow, Russia

B.A. Dyachkov

East-Kazakhstan State Technical University, Kazakhstan

M.A. Mizernaya

East-Kazakhstan State Technical University, Kazakhstan

Nina Maiorova

East-Kazakhstan State Technical University, Kazakhstan

Zinaida Chernenko

East-Kazakhstan State Technical University, Kazakhstan

Victor Maiorov

East-Kazakhstan State Technical University, Kazakhstan

O.N. Kuzmina

East-Kazakhstan State Technical University, Kazakhstan

Vsevolod Yutsis

Universidad Autónoma de Nuevo León, Facultad de Ciencias de la Tierra, Linares, N.L., Mexico

Antonio Tamez Ponce

Posgrado en Ciencias Geológicas, Facultad de Ciencias de la Tierra, Universidad Autónoma de Nuevo León, Linares, N.L., Mexico

Konstantin Krivosheya

Universidad Autónoma de Nuevo León, Facultad de Ciencias de la Tierra, Linares, N.L., Mexico

Feng Gui

Baker Hughes, Perth, Australia

Khalil Rahman

Baker Hughes, Perth, Australia

Daniel Moos

Baker Hughes, Menlo Park, USA

George Vassilellis

Gaffney, Cline & Associates, Houston, USA

Chao Li

Gaffney, Cline & Associates, Houston, USA

Qing Liu

Baker Hughes, Beijing, China

Fuxiang Zhang

PetroChina Tarim Oil Company, Korla, China

Jianxin Peng

PetroChina Tarim Oil Company, Korla, China

Xuefang Yuan

PetroChina Tarim Oil Company, Korla, China

Guoqing Zou

PetroChina Tarim Oil Company, Korla, China

Nima Gholizadeh Doonechaly

School of Petroleum Engineering, University of New South Wales, Sydney, Australia

Sheik S. Rahman

School of Petroleum Engineering, University of New South Wales, Sydney, Australia

Andrei Kotousov

School of Mechanical Engineering, the University of Adelaide, South Australia, Australia

Víctor Manuel Arellano

Instituto de Investigaciones Eléctricas, Gerencia de Geotermia, Cuernavaca, México

Rosa María Barragán

Instituto de Investigaciones Eléctricas, Gerencia de Geotermia, Cuernavaca, México

Miguel Ramírez

Comisión Federal de Electricidad, Gerencia de Proyectos Geotermoelectricos, Morelia, México

Siomara López

Instituto de Investigaciones Eléctricas, Gerencia de Geotermia, Cuernavaca, México

Alfonso Aragón

Instituto de Investigaciones Eléctricas, Gerencia de Geotermia, Cuernavaca, México

Adriana Paredes

Instituto de Investigaciones Eléctricas, Gerencia de Geotermia, Cuernavaca, México

Emigdio Casimiro

Comisión Federal de Electricidad, Residencia Los Azufres, Campamento Agua Fría, México

Lisette Reyes

Comisión Federal de Electricidad, Residencia Los Azufres, Campamento Agua Fría, México

Marian Petre

Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry (IGEM), Russia

Sharkov

Russian Academy of Sciences, Moscow, Russia

Charles B. Moss

Food and Resource Economics Department, University of Florida, Gainesville, FL, USA

Andrew Schmitz

Food and Resource Economics Department, University of Florida, Gainesville, FL, USA

Yugui Yang

State Key Laboratory for Geomechanics and Deep Underground Engineering,
China University of Mining and Technology, Xuzhou, Jiangsu 221008, China
School of Mechanics and Civil Engineering, China University of Mining and
Technology, Jiangsu 221116, China

Feng Gao

School of Mechanics and Civil Engineering, China University of Mining and
Technology, Jiangsu 221116, China

Yuanming Lai

State Key Laboratory of Frozen Soil Engineering, Cold and Arid Regions
Environmental and Engineering Research Institute, Chinese Academy of Sci-
ences, Lanzhou 730000, China

Jatta Saarenheimo

Department of Biological and Environmental Science, University of Jyväskylä, 40014, Jyväskylä, Finland

Antti J. Rissanen

Department of Biological and Environmental Science, University of Jyväskylä, 40014, Jyväskylä, Finland

Lauri Arvola

Lammi Biological Station, University of Helsinki, 16900, Lammi, Finland

Hannu Nykänen

Department of Biological and Environmental Science, University of Jyväskylä, 40014, Jyväskylä, Finland

Moritz F. Lehmann

Department for Environmental Science, University of Basel, CH-4058, Basel, Switzerland

Marja Tiirola

Department of Biological and Environmental Science, University of Jyväskylä, 40014, Jyväskylä, Finland

Jennifer L. Miksis-Olds

Applied Research Laboratory, The Pennsylvania State University, State College, Pennsylvania, United States of America

Laura E. Madden

Applied Research Laboratory, The Pennsylvania State University, State College, Pennsylvania, United States of America

Preface

Encyclopaedia of Geomechanics in Soil, Rock, and Environmental Engineering deals with natural risk factors, of both natural and human origin, that should be borne in mind. *Environmental Geomechanics* covers a range of topics that are of increasing importance in engineering practice: natural hazards, pollution, and environmental protection through good practice. Transport of contaminants and other substances may occur in the fluids, e.g. water, water vapour and air, filling the pores of geomaterials as happens in waste disposal problems or durability problems. Mass transport also takes place in reservoir engineering problems, where the fluids involved are oil, water, and gas. First chapter deals with “alive” tectonomagmatic processes, which are reflected in neotectonics, topography, geophysical fields, and the present-day magmatic activity. Geotectonic position and metallogeny of the greater Altai geological structures in the system of the Central-Asian mobile belt is proposed in second chapter. Third chapter presents an approach on geophysical modeling of the surroundings of La Popa Basin, Ne Mexico, with gravity and magnetic data. Fourth chapter provides a comprehensive geomechanical study to optimize stimulation for a fractured tight gas reservoir in the northwest Tarim Basin. In fifth chapter, an innovative analytical approach based on the distributed dislocation technique is developed to simulate the roughness induced opening of fractures in the presence of compressive and shear stresses as well as fluid pressure inside the fracture. This provides fundamental basis for computation of aperture distribution for all parts of the fracture which can then be used in the next step of modeling fluid flow inside the fracture as a function of time. It also allows formulation of change in aperture due to thermal stresses. The objective of sixth chapter was to investigate the exploitation-related processes through the analysis of geochemical and production data of 39 wells. The goal of seventh chapter is to show that the present-day tectonomagmatic activity within the TEB can be interpreted that a new ocean has begun to open here. Eighth chapter examines the changes in land use between 1947 and 2007 focusing on the possibility that commercial uses generate significant environmental benefits. In ninth chapter, it is recognized experimentally that the compressibility of warm and ice-rich frozen soil is remarkable under loading, which will cause a significant deformation and affect the stability of infrastructure constructed in cold region. In tenth chapter, we studied potential links between environmental factors, nitrous oxide (N_2O) accumulation, and genetic indicators of nitrite and N_2O reducing bacteria in 12 boreal lakes. In last chapter, we identify specific environmental parameters, including components of the ambient background sound that are predictive of ice seal presence in the Bering Sea.

Contents

	<i>List of Abbreviations</i>	<i>vii</i>
	<i>List of Contributors.....</i>	<i>ix</i>
	<i>Preface.....</i>	<i>xv</i>
Chapter 1	Geological-Geomechanical Simulation of the Late Cenozoic Geodynamics in the Alpine-Mediterranean Mobile Belt.....	1
	E.V. Sharkov and V.B. Svalova	
Chapter 2	Geotectonic Position and Metallogeny of the Greater Altai Geological Structures in the System of the Central-Asian Mobile Belt	27
	B.A. Dyachkov, M.A. Mizernaya, Nina Maiorova, Zinaida Chernenko, Victor Maiorov and O.N. Kuzmina	
Chapter 3	Geophysical Modeling of the Surroundings of La Popa Basin, NE Mexico, with Gravity and Magnetic Data	51
	Vsevolod Yutsis, Antonio Tamez Ponce, and Konstantin Krivosheya	
Chapter 4	Optimizing Hydraulic Fracturing Treatment Integrating Geomechanical Analysis and Reservoir Simulation for a Fractured Tight Gas Reservoir, Tarim Basin, China	77
	Feng Gui, Khalil Rahman, Daniel Moos, George Vassilellis, Chao Li, Qing Liu, Fuxiang Zhang, Jianxin Peng, Xuefang Yuan, and Guoqing Zou	
Chapter 5	A New Approach to Hydraulic Stimulation of Geothermal Reservoirs by Roughness Induced Fracture Opening.....	107
	Nima Gholizadeh Doonechaly, Sheik S. Rahman and Andrei Kotousov	
Chapter 6	Reservoir Processes Related to Exploitation in Los Azufres (México) Geothermal Field Indicated by Geochemical and Production Monitoring Data.....	133
	Víctor Manuel Arellano, Rosa María Barragán, Miguel Ramírez, Siomara López, Alfonso Aragón, Adriana Paredes, Emigdio Casimiro, and Lisette Reyes	
Chapter 7	Does the Tethys Begin to Open Again? Late Cenozoic Tectonomagmatic Activation of the Eurasia from Petrological and Geomechanical Points of View	153
	Marian Petreand Sharkov	

Chapter 8 **Environmental Flows from Alternate Land Uses in the Delta, Pacific, and the Southeastern States: 1947-2007** 173
Charles B. Moss and Andrew Schmitz

Chapter 9 **Compressive Mechanical Properties and Micromechanical Characteristics of Warm and Ice-Rich Frozen Silt** 189
Yugui Yang, Feng Gao, and Yuanming Lai

Chapter 10 **Genetic and Environmental Controls on Nitrous Oxide Accumulation in Lakes**..... 205
Jatta Saarenheimo, Antti J. Rissanen, Lauri Arvola, Hannu Nykänen, Moritz F. Lehmann, and Marja Tirola

Chapter 11 **Environmental Predictors of Ice Seal Presence in the Bering Sea**..... 231
Jennifer L. Miksis-Olds and Laura E. Madden

Citations 299

Index..... 303

Chapter 1

GEOLOGICAL-GEOMECHANICAL SIMULATION OF THE LATE CENOZOIC GEODYNAMICS IN THE ALPINE- MEDITERRANEAN MOBILE BELT

E.V. Sharkov¹ and V.B. Svalova²

¹Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry
RAS, Moscow, Russia

²Institute of Geocology RAS, Moscow, Russia

INTRODUCTION

Alpine-Mediterranean Mobile Belt, which currently ongoing to develop, is one of the best sites for studying of geodynamic mechanisms for formation of such regions. In this case we are dealing with “alive” tectonomagmatic processes, which are reflected in neotectonics, topography, geophysical fields, and the present-day magmatic activity. Last circumstance allows independent control of petrological processes in the underlying mantle of the belt. Comparison of geological-geophysical data available with the results of mechanic and mathematic simulation allows us to establish the relationships of all these processes and the character of their manifestation on the Earth's surface. This is the purpose of our study.

GEOLOGY AND PETROLOGY OF ALPINE BELT

Alpine-Mediterranean Mobile Belt (Alpine Belt) represents the western part of the huge Alpine-Himalayan collision zone, which appeared in the late Cretaceous-early Paleogene after closure of the Tethys Ocean. The suture of this neotectonic zone is traced by chain of late Cenozoic andesite-latitude volcanism, stretching across Eurasia from Mediterranean to the Indonesian Island Arc and back-arc seas of the Western Pacific as well as areas of continental rifting and areas of intraplate basaltic volcanism.

The most complicated structure of this belt is in its west, in the Alpine segment (Fig. 1), where there is a whole system of mountain ridges, andesite-latite volcanic arcs and back-arc basins with thinned crust of intermediate to oceanic-type (Alboran, Tyrrhenian, Aegean Sea, and Pannonian Basin) occurs. Despite the differences in the morphology of these structures, they have several common features: along their periphery volcanic arcs and fold-thrust belts which form arc-shaped mountain ridges are developed. Among their thrust slices are often observed deep-water sedimentary rocks of Tethys, ophiolitic complexes, and sometimes blocks of the lower crust and upper mantle. In general, the situation in many aspects is similar to that which takes place on the active margins of continents and oceans. Such structures are characterized mainly for the West Mediterranean, while for the Eastern Mediterranean, as well as for the Black and Caspian seas typical passive margin. For this reason, we divide the Alpine Belt into two segments: the eastern, or the Aegean-Caucasian, and western, or proper Alpine, which will be considered separately.



Figure 1. Development of the Late Cenozoic igneous rocks within the Alpine Belt 1 – back-arc seas (A – Alboran, T – Tyrrhenian; Ae – Aegean) and “downfall” seas (B – Black, C – Caspian); 2 – back-arc sedimentary basins (P – Pannonian, Po – Po valley); 3 – Late Cenozoic andesite-latite volcanic arcs (in circles): 1 – Alboran, 2 – Cabil-Tell, 3 – Sardinian, 4 – South-Italian, 5 – Drava-Insubrian, 6 – Evganey, 7 – Carpatian, 8 – Balkanian, 9 – Aegean, 10-12 – Anatolia-Elbursian (10- Anatolia-Caucasian, 11 – zone of the Modern Caucasus volcanism, 12 – Caucasus-Elbursian); 4 – areas of flood basaltic volcanism (in square): 1 – South Spain and Portugal, 2 – Atlas, 3 – Eastern Spain, 4 – Central France massif, 5 – Rhine graben, 6 – Czech-Silesian, 7 – Pannonian, 8 – Western Turkey, 9 – northern Arabia; 5 – suture zones of major thrust structures interaction of a superplume head with mobile continental lithosphere. Good example of

such situation is the TEB (Sharkov, this book), where processes of collision continue now. The main feature of this belt is wide spread of the Late Cenozoic-derived volcanism, which has displayed practically coeval conditions on all its length, presuming existence of a superplume (or asthenospheric rise) beneath it. The belt has the most complicated structure within the Alpine segment, where a system of andesite-latite volcanic arcs and back-arc basin, bordering by nappe-folded mountain ridges were observed. In front of these ridges in Western Europe, north-west Africa and Arabia, coeval rift systems and flood basaltic volcanism often occur

Aegean-Caucasian Segment

Caucasian part of the segment is located to the north of the major suture zone traced by ophiolites of Cyprus, Syria, southeastern Turkey, Zagros, etc. (Periarabian ophiolitic arc). Late Cenozoic Anatolian-Elbursian andesite-latite volcanic arc is developed at its rear. It, in turn, is formed by two arcs – Anatolian-Caucasian and Caucasian-Elbursian, touching in the transverse (Transcaucasian) zone of the latest volcanism (Fig. 1). The Black and Caspian seas with oceanic crust, cut Pre-Pliocene structure of the Caucasus and the Kopet Dag; they are filled with Mesocenozoic sediments of 12-15 km thick. The nature of these deposits, until the late Miocene was generally similar to developed within the Caucasus and the Kopet Dag (Zonenshain, Le Pichon, 1986).

The Caucasus is located in the zone of the Arabian syntax (Burtman, 1989), where Arabian plate is subducted beneath Eurasian. Specific analog of deep-sea trench is represented by the Mesopotamian trough here, which, beginning from the Eocene, is experiencing an active submergence, due to thick molasse accumulated (Ponikarov et al, 1969). The northern part of the Arabian plate began to rise above sea level in the late Oligocene and early Miocene, about 26-25 Ma when there began the development of basaltic volcanism (Sharkov, 2000) and the Red Sea rift opened up in southwest, separating Arabian plate from Africa. Rate of ascending movements sharply increased to the Miocene-Pliocene boundary, when Gulf of Aqaba opened approximately 5 Ma and Arabian plate began quickly shifted to the north along a large Levant Fault (Dead Sea transform) (Kopp, Leonov, 2000; Prilepin et al, 2001; Sharkov, this book). However, this displacement is hardly manifested in the Greater Caucasus shift and, moreover, GPS data indicate that the width of the central Caucasus is not decreasing but increasing (Shevchenko et al, 1999)

Specific structure occurred at the northern side of the Black Sea. Judging from the geological and geophysical data along the profile of Tuapse-Armavir, the Black Sea microplate is separated from the Eurasian by narrow subvertical zone of strong positive gravitational anomalies (Shempelev and et al, 2001). It concerned to large blocks of deformed and metamorphic rocks, close on the

density to crust-mantle mixture. This zone can be traced to depth of 60-70 km and the Moho is not established here. The northern blocks move upward along their separating steeply-dipping faults, ensuring the existence of mountain relief of the Western Caucasus.

Formation of the Black Sea began, apparently, in the early Cretaceous, but significant deepening of the basin occurred at the Oligocene-Early Miocene boundary (Zonenshain, Le Pichon, 1986, Nikishin et al, 2001), followed in the Miocene by filling of the deep-water depressions by sediments and a gradual shallowing of the basin (Kazmin et al, 2000). Since the Pliocene-Quaternary new significant deepening of the Black Sea basin has occurred (Nikishin, Karataev, 2000), which occurred almost simultaneously with the uplift of the Caucasus and Crimea, which in the Oligocene-early Miocene were not expressed in the relief (Neotectonics..., 2000; Kostenko, Panina, 2001). The close sequence of events took place in the South Caspian Basin, which is a similar structure (Zonenshain, Le Pichon, 1986; Grachev, 2000).

Aegean part of the segment is characterized by island arc associated with subduction of the oceanic East-Mediterranean plate beneath the Eurasian (Papazachos et al., 1995). There are all typical structures of the active zones of transition from continent to ocean here: the deep-water Hellenic trough, Aegean volcanic arc and back-arc Aegean Sea with basaltic magmatism in its periphery (in the west of Asia Minor, near Izmir).

Numerous deformations of extension in subhorizontal submeridional direction (strike-slip and normal faults, nappe-thrusts, grabens, etc.) are known in the Aegean basin and in adjacent parts of Greece and Turkey (Prilepin et al, 2001). At that for Balkan Mountains in the north are characteristic north-vergentes imbricate nappes and thrusts whereas for Hellenides-Aegides-Taurides they are south-vergent. Judging from the seismic data, the stress state of the subhorizontal N-S stretching is characteristic of only for the upper 50-60 km of the Aegean basin lithosphere, leading to its expansion. At greater depths within the mantle beneath the basin both in south and north are fixed compression conditions. The regions of extension and compression are in direct contact by subhorizontal section at depth of 70-80 km; probably, it is the boundary of lithosphere and plastic material of extended plume head.

Aegean volcanic arc is a Pleistocene in age, but the development of the Aegean Sea, started earlier, about 12 Ma (Evsyukov, 1998). Apparently, earlier the main subduction zone located north, and its residues were survived in Dinarides and western Asia Minor (Western Anatolian: Fig. 1).

Descent of the Eastern Mediterranean (Ionian Sea, underwater Medina Ridge, Levant Basin) began in the late Miocene (Evsyukov, 1999).

Approximately at the middle Pliocene (about 3-3.5 Ma) processes reinforced on the east: judging by the results of the 5th Cruise of R/V "Akademik Strakhov", Sinai plate descended here beneath the sea level to depths of 2-2.5 km (Geological..., 1994). Fragments of this plate are preserved in the form of Eratosthenes Seamount and smaller rises.

The northern part of the Eastern Mediterranean is separated from the southern part by zone of powerful deformations, which runs along the base of the Cyprus arc (Zverev, 2002). As in the case of the northern side of the Black Sea, large sub-vertical faults and a sharp increase in the boundary velocities up to 7-7.6 km/sec is recorded here. In the northern zone is observed uplift of blocks of basement, represented by ophiolites of Cyprus. This deformation zone begins at the northern end of the Arabian plate, and following along Periarabian ophiolite belt, comes to Cyprus and then, continuing to the west, joins the Aegean subduction zone (see Figure 1).

The situation on the eastern passive margin of the Mediterranean Sea also looks like northern passive margin of the Black Sea. System of subparallel mountain ranges (Lebanon, Anti-Lebanon, Jabal al-Ansari, Amanus, etc.), separated by a system of steeply dipping faults, occur along the coast here. The largest of the latter is the aforementioned Levant (Dead Sea) Transform Fault. Parallel to it, already under water, there is a zone of Pelizium faults bounding the east Levant deep depression (Khair, Tsokas, 1999). This depression has the oceanic crust, overlapping by thick (approximately 10 km) sequence of Phanerozoic sediments, and in this respect no different from the Black and Caspian seas.

Origin of within-plate Palmirides folded-thrusted zone of deformed platform cover is related to development of Levant Fault. According to Kopp and Leonov (2000), the formation of this structure was caused by braking of the western edge of the Arabian plate in the zone above the bending S-like curve of this fault at its motion to the north during the Neogene-Quaternary. Palmirides were formed under compression of the crust by approximately 20-25 km, compensating the northern Arabian plate movement that started in the Middle-Late Miocene and continues today.

Areas of intraplate moderate alkaline basaltic volcanism are widely developed in the northern Arabian plate, indicating the presence of a mantle plume here. Judging from the isotopic dating, basalt eruption began at the Eocene-Miocene boundary, about 25-26 Ma, and almost without interruption continued until historic times. The most powerful eruption occurred in the late Miocene-early Pliocene and Late Pliocene-Quaternary (Sharkov, 2000; Lustrino, Sharkov, 2006).