

**TUMORS
OF THE
KIDNEY, RENAL PELVIS, AND URETER**

by

**JAMES L. BENNINGTON, M.D.
and
J. BRUCE BECKWITH, M.D.**

ATLAS OF TUMOR PATHOLOGY

Second Series

Fascicle 12

TUMORS OF THE KIDNEY, RENAL PELVIS, AND URETER

by

JAMES L. BENNINGTON, M. D.

Chairman

Department of Anatomic and Clinical Pathology

Children's Hospital of San Francisco

San Francisco, California 94119

and

Associate Clinical Professor of Pathology

University of California

San Francisco Medical Center

and

J. BRUCE BECKWITH, M. D.

Professor of Pathology and Pediatrics

University of Washington School of Medicine

and

Director of Laboratories

The Children's Orthopedic Hospital and Medical Center

Seattle, Washington 98105

Published by the

~~ARMED FORCES INSTITUTE OF PATHOLOGY~~

~~Washington, D. C.~~

~~Under the auspices of~~

UNIVERSITIES ASSOCIATED FOR RESEARCH AND EDUCATION IN PATHOLOGY, INC.

Bethesda, Maryland

1975

Accepted for Publication

1975

.....

For sale by the American Registry of Pathology

Armed Forces Institute of Pathology

Washington, D. C. 20306

ATLAS OF TUMOR PATHOLOGY

Sponsored and Supported by

AMERICAN CANCER SOCIETY*
NATIONAL CANCER INSTITUTE, NATIONAL INSTITUTES OF HEALTH*

and

ARMED FORCES INSTITUTE OF PATHOLOGY

EDITOR

HARLAN I. FIRMINER, M. D.

Professor, Department of Pathology
University of Maryland School of Medicine
Baltimore, Maryland 21201

and

Scientist Associate, Universities Associated for
Research and Education in Pathology, Inc.
9650 Rockville Pike
Bethesda, Maryland 20014

EDITORIAL ADVISORY COMMITTEE

Lauren V. Ackerman, M.D.	State University of New York Stony Brook, Long Island, New York
David G. Freiman, M.D.	Harvard Medical School Boston, Massachusetts
John K. Frost, M.D.	The Johns Hopkins School of Medicine Baltimore, Maryland
James C. Harkin, M.D.	Tulane University School of Medicine New Orleans, Louisiana
Dante G. Scarpelli, M.D.	University of Kansas Medical Center Kansas City, Kansas
Robert E. Scully, M.D.	The Massachusetts General Hospital Boston, Massachusetts
Philippe Shubik, M.D.	University of Nebraska College of Medicine Omaha, Nebraska
Harlan J. Sojut, M.D.	Baylor University College of Medicine Houston, Texas

*(Supported by Public Health Service grant no. CA08575 from the
National Cancer Institute and American Cancer Society grant no. MG-122)

EDITOR'S NOTE

The Atlas of Tumor Pathology was originated by the Committee on Pathology of the National Academy of Sciences—National Research Council in 1947. The form of the Atlas became the brainchild of the Subcommittee on Oncology and was shepherded by a succession of editors. It was supported by a long list of agencies; many of the illustrations were made by the Medical Illustration Service of the Armed Forces Institute of Pathology; the type was set by the Government Printing Office; and the final printing was made by the press at the Armed Forces Institute of Pathology. The American Registry of Pathology purchased the fascicles from the Government Printing Office and sold them at cost, plus a small handling and shipping charge. Over a period of 20 years, 15,000 copies each of 40 fascicles were produced. They provided a system of nomenclature and set standards for histologic diagnosis which received worldwide acclaim. Private contributions by almost 600 pathologists helped to finance the compilation of an index by The Williams & Wilkins Company to complete the original Atlas.

Following the preparation of the final fascicle of the first Atlas, the National Academy of Sciences—National Research Council handed over the task of further pursuit of the project to Universities Associated for Research and Education in Pathology, Inc. Grant support for a second series was generously made available by both the National Cancer Institute and the American Cancer Society. The Armed Forces Institute of Pathology has expanded and improved its press facilities to provide for a more rapid and efficient production of the new series. A new Editor and Editorial Advisory Committee were appointed, and the solicitation and preparation of manuscripts continues.

This second series of the Atlas of Tumor Pathology is not intended as a second edition of the first Atlas and, in general, there will be variation in authorship. The basic purpose remains unchanged in providing an Atlas setting standards of diagnosis and terminology. Throughout the rest of this new series, the term chosen for the World Health Organization's series "International Histological Classification of Tumours" (when available) is shown by an asterisk if it corresponds to the authors' choice, or as the first synonym in bold print if it differs from the authors' heading. Hematoxylin and eosin stained sections still represent the keystone of histologic diagnosis; therefore, most of the photomicrographs will be of sections stained by this technic, and only sections prepared by other technics will be specifically designated in the legends. It is hoped that in many of the new series a broader perspective of tumors may be offered by the inclusion of special stains, histochemical illustrations, electron micrographs, data on biologic behavior, and other pertinent information for better understanding of the disease.

The format of the new series is changed in order to allow better correlation of the illustrations with the text, and a more substantial cover is provided. An index will be included in each fascicle.

It is the hope of the Editor, the Editorial Advisory Committee, and the Sponsors that these changes will be welcomed by the readers. Constructive criticisms and suggestions will be appreciated.

Harlan I. Firminger, M. D.

PREFACE AND ACKNOWLEDGMENTS

In the nearly two decades since the publication of the First Series of the Fascicle on Tumors of the Kidney, Renal Pelvis, and Ureter, there has been a virtual explosion of new information on tumors of the upper urinary tract. Many significant contributions concerning nearly all of the tumors covered in the First Series Fascicle, as well as the recognition of important new entities, have been made during this time. The authors have attempted to incorporate into this text the new well established findings in the areas of epidemiology, histogenesis, histochemistry, morphology, classification, diagnosis, treatment, and prognosis, as well as our own views on unproved and controversial theories relating to this field. While this has been largely a joint effort, Dr. Beckwith was responsible primarily for those sections relating to embryonal tumors of the kidney and Dr. Bennington for the remainder.

We wish to acknowledge the excellent First Series Fascicle by Dr. Balduin Lucké and Dr. Hans G. Schlumberger, which was such a tremendous help in the development of our manuscript and the illustrations that we have been permitted to use. They are our current figure numbers 40, 57, 58, 67, 175, 176, 201, 206, 216–218, 247, 257, and 259.

The classification and nomenclature used in this fascicle are based on our understanding of the histogenetic origin of the various tumors of the kidney, renal pelvis, and ureter. This terminology, with minor exceptions, is in general use in English speaking countries and conforms to that proposed (but not yet adopted) by the World Health Organization (WHO).

The authors wish to express their gratitude for the support in part by the U. S. Public Health Service, Grant No. R10-CA 11722, of the National Wilms' Tumor Study.

The majority of the material used in the preparation of this text was gathered at the University of Washington Medical School, Seattle, Washington; Children's Orthopedic Hospital, Seattle, Washington; King County Hospital, Seattle, Washington; Kaiser Foundation Hospital, Oakland, California; Children's Hospital, San Francisco, California; and from the National Wilms' Tumor Registry. We are particularly grateful to Dr. Seth L. Haber, Santa Clara, California; Dr. George Farrow, Rochester, Minnesota; Dr. Roger Pugh and Dr. Noel Gowing, London, England for their kindness and generosity in providing much unusual and important material. We are also indebted to the following other contributors of materials and illustrations:

Figures 9, 10, 11 - Dr. Craig Tisher, Durham, N. C.

Figure 19 - Dr. Ruth E. Bulger, Baltimore, Md.

Figure 24 - Dr. Benjamin H. Landing, Los Angeles, Calif.

Figure 25 - Dr. Nathan B. Friedman, Los Angeles, Calif.

Figures 26, 27, 75–77, 185, 186, and Plates IV-A, B, and VII-B - Dr. Seth L. Haber, Santa Clara, Calif.

Figure 29 - Dr. Warren W. Johnson, Memphis, Tenn.

Figures 32, 33 - Dr. Howard Ricketts, St. Louis, Mo.

Figures 39, 220, 222, and Plates X-D, and XI-A, B - Armed Forces Institute of Pathology

Figure 67 - Dr. Sidney Farber, Boston, Mass.

Figures 68, 145 - Dr. Joseph B. Crawford, San Francisco, Calif.

Figure 83 - Dr. H. L. Ratcliffe, Philadelphia, Pa.

Figure 89 - Dr. Louis Komarmy, San Francisco, Calif.

Figures 90, 91, 92 - Dr. J. S. Ansell, Seattle, Wash.

Figures 94, 96, 188, 200 - Dr. Joachim Burhenne, San Francisco, Calif.

Figure 97 - Dr. George Annes, San Francisco, Calif.

Figure 98 - Dr. R. M. Shishido, San Diego, Calif.

Figure 101 - Dr. James L. McRea, Sydney, Australia

Figure 126 - Dr. M. D. Lagios, San Francisco, Calif.

Figure 133 - Dr. R. Seljelid, Tromso, Norway

Plate VI-A - Dr. Vivian Gildenhorn, Los Angeles, Calif.

Figure 138 - Dr. Bruce Mackay, Houston, Tex.

Figures 140, 157 - Dr. Noel Gowing, London, England

Figures 142, 203, 248 - Dr. Roger Pugh, London, England

Figure 143 - Dr. A. Lazar, Leonia, N. J.

Figure 144 - Dr. N. L. Morgenstern, Oakland, Calif.

Figure 152 - Dr. Averill A. Liebow, San Diego, Calif.

Figures 196, 221 - Dr. P. N. Cowen, Leeds, England

Figures 197, 215 - Dr. Bradford Young, San Francisco, Calif.

Figure 231 - Dr. Carlos Perez-Mesa, Columbia, Mo.

Figure 249 (lower) - Dr. P. W. Graff, Chicago, Ill.

Figures 251–253 - Dr. R. Kempson, Stanford, Calif.

Figures 257, 259 - Dr. Beecher-Smith, Columbus, Ohio

Figure 267 - Dr. B. L. Pear, Denver, Colo.

Plate XI-D - Dr. D. C. Schneiderman, Montreal, Canada

The following illustrations have also been used. They appeared in *Renal Carcinoma* by Drs. J. L. Bennington and R. Kradjian, published by W. B. Saunders Co. in 1967, and cover the following subjects:

Embryology and Anatomy of the Kidney: Our figures 1 and 4 (modified), 5, 9–11.

Histogenesis of Renal Carcinoma: Our figure 80.

Epidemiology and Etiology: Our figures 85, 86.

Morphology: Our figures 83, 87, 104–107, 109, 113, 116–118, 125, 129, 143, 144, 147, plates III-B and V, and tables V and VI (modified).

Distribution of Metastases from Renal Carcinoma: Our figures 139, 141, 155, and tables VII and VIII (modified).

Diagnostic Techniques: Our figures 90–92.

We gratefully acknowledge the secretarial assistance of Judy Fletcher and Nondis Barrett who have patiently and cheerfully carried the burden of preparing the manuscript. We also wish to thank Sheila Concannon for preparation of photographic illustrations, Dr. Michael Lagios for the original electron micrographs, Dr. John Azzopardi who kindly reviewed the manuscript and made many helpful suggestions, and the Armed Forces Institute of Pathology for the use of their facilities. Our special thanks go to Dr. Harlan I. Firminger, his editorial staff, and the reviewers chosen by the Editorial Advisory Committee for all their encouragement, kindness, and much needed patience.

Permission to use copyrighted illustrations has been granted by:
Copyright 1973 by: American Association for the Advancement of Science:
Science 179:393-395, 1973. For our figure 195.

American Cancer Society, Inc.:

Cancer 21:727-742, 1968. For our figure 81.
Cancer 22:545-550, 1968. For our figures 169, 171, 173, 177.
Cancer 22:556-563, 1968. For our figure 121.
Cancer 22:564-570, 1968. For our figure 161.
Cancer 24:535-542, 1969. For our figure 249.
Cancer 29:1597-1605, 1972. For our figure 34.
Cancer 32:1030-1042, 1973. For our figure 158.
Cancer 32:1078-1083, 1973. For our figures 202, 204.

American Medical Association:

Arch. Pathol. 76:277-289, 1963. For our figures 2, 3.
J.A.M.A. 204:753-757, 1968. For our figure 93.

British Medical Association: London

J. Clin. Pathol. 23:472-474, 1970. For our figure 78.
J. Clin. Pathol. 23:681-689, 1970. For our figures 260-263.

Churchill Livingstone (Longman Group, Ltd.): Edinburgh

Br. J. Urol. 44:517-527, 1972. For our figure 102.
Monograph on Neoplastic Diseases at Various Sites, Vol. V.
Tumors of the Kidney and Ureter, 1964. For our figure 35 and graph VII (modified).

H. K. Lewis & Co., Ltd.: London

Br. J. Cancer 12:507-516, 1958. For our table I (modified).

Harper & Row, Publishers, Inc.-Hoeber Medical Division:

Lab. Invest. 14:435-447, 1965. For our figure 132 A, B.
Lab. Invest. 15:1357-1394, 1966. For our figures 6, 7.

Lea & Febiger:

Renal Diseases, 2d Ed., 1950. For our graph V (data).

Mayo Clinic:

Mayo Clin. Proc. 45:161-169, 1970. For our figure 88.

The National Foundation—March of Dimes: (Ed.) Bergsma, D. Part II
Malformation Syndromes. Vol. V(2): White Plains, N. Y.

Birth Defects: Orig. Art. Ser. 191, 193, 1969. For our figures 28, 31.

New York Academy of Sciences:

Ann. N. Y. Acad. Sci. 126:188-203, 1965. For our figure 84.

- Oliver & Boyd, Ltd.: Edinburgh
J. Pathol. 87:424-425, 1964. For our figure 250.
- Penrose Cancer Hospital:
Penrose Cancer Seminar, Vol. IV, No. 5, 1971. For our figure 266.
- Prentice-Hall, Inc.: Sheldon C. Sommers, Englewood Cliffs, N. J.:
Pathol. Annu. 3:213-224, (c) 1968. For our figure 18.
- The Reuben H. Donnelley Corp.:
Am. J. Med. 43:963-976, 1967. For our figures 179, 180.
Am. J. Med. 55:86-92, 1973. For our plate IX-A, B, C.
- The Royal Veterinary College: London
J. Comp. Pathol. 78:335-339, 1968. For our figure 82.
- Springer-Verlag: Berlin-Heidelberg-New York
Virchows Arch. (Pathol. Anat.) 341:204-223, 1966. For our figure 134.
- W. B. Saunders Co.:
Renal Carcinoma, 1967. For our figures 1, 5, 9-11, 80, 83, 85-87, 90-92, 104-107, 109, 113, 116-118, 125, 129, 139, 141, 143, 144, 147, 155, plates III-B and V, and tables V, VI, VII, VIII (modified).
A Textbook of Histology, 8th Ed., 1962. For our figure 4 (modified).
- The Williams & Wilkins Co.:
J. Urol. 93:139-143, 1965. For our figure 17.
J. Urol. 94:356-361, 1965. For our figure 254.
J. Urol. 101:297-301, 1969. For our graph VI.
J. Urol. 102:291-293, 1969. For our figure 16.
J. Urol. 102:678-682, 1969. For our figure 258.
J. Urol. 104:528-531, 1970. For our figures 183, 184.
J. Urol. 106:515-517, 1971. For our figure 99.
J. Urol. 109:101-103, 1973. For our figure 74.
J. Urol. 109:567-568, 1973. For our figure 209.
- The Wistar Institute of Anatomy and Biology:
Am. J. Anat. 116:237-255, 1965. For our figure 8.

James L. Bennington, M. D.
J. Bruce Beckwith, M. D.

TUMORS OF THE KIDNEY, RENAL PELVIS, AND URETER

Contents

	Page No.
Embryology and Anatomy of the Upper Urinary Tract	1
Embryogenesis	1
Histologic Structure of the Nephron	4
Histologic Structure of the Renal Calix, Pelvis, and Ureter	13
Heterotopic Tissue in the Kidney	15
Endometrial Tissue in the Kidney, Renal Pelvis, and Ureter	16
Cartilage in Renal Stroma	19
Adipose Tissue and Smooth Muscle	20
Interstitium of the Kidney	22
Tumors of the Upper Urinary Tract	25
Embryonal Tumors	31
Nephroblastoma	31
Histogenesis	32
Comparative Pathology	44
Relation to Congenital Mesoblastic Nephroma	46
Epidemiology	46
Clinical Pathologic Features	52
Natural History and Patterns of Metastasis	74
Treatment	77
Prognosis	78
Congenital Mesoblastic Nephroma	79
Solitary Multilocular Cyst of the Kidney	81
Renal Adenocarcinoma	93
Pathogenesis	93
Comparative Pathology	99
Epidemiology	104
Etiologic Factors	107
Clinical Findings	108
Laboratory Findings	111
Diagnostic Technics	114
Pathologic Features	128
Differential Diagnosis from Other Primary Tumors	160
Distribution of Metastases from Renal Adenocarcinoma ...	168

Differential Diagnosis of Tumors in Other Sites	
from Metastatic Renal Adenocarcinoma	178
Prognosis	186
Gross Features	188
Microscopic Features	192
Mesenchymal Tumors of the Kidney	201
Introduction	201
Tumors of Muscle and Adipose Tissue	201
Leiomyoma and Lipoma	201
Angiomyolipoma	204
Leiomyosarcoma	213
Rhabdomyosarcoma	216
Liposarcoma	217
Tumors of Vascular Tissue	220
Hemangioma	220
Angiosarcoma	222
Hemangiopericytoma	222
Juxtaglomerular Tumor	224
Lymphangioma	227
Tumors of Fibrous Tissue	229
Fibroma	229
Fibrosarcoma	229
Fibroxanthosarcoma	229
Mesoblastic Nephroma in the Adult	231
Renomedullary Interstitial Cell Tumor	231
Tumors of Neurogenic Tissue	232
Tumors of Osteogenic Tissue	233
Tumors of Hematopoietic Tissue	234
Leukemia	234
Malignant Lymphoma	237
Multiple Myeloma	239
Tumors of the Renal Pelvis and Ureter	243
Epithelial Tumors	243
Histogenesis	243
Comparative Pathology	247
Epidemiology	248
Clinical Features	252
Laboratory Findings	253
Radiographic Findings	255
Pathologic Features	255

Papilloma	264
Pathologic Features	264
Carcinoma	266
Pathologic Features	266
Distribution of Metastases	309
Prognosis	309
Treatment	310
Mesodermal Tumors of the Renal Pelvis and Ureter	312
Tumors of Smooth Muscle	312
Leiomyoma	312
Leiomyosarcoma	312
Tumors of Neurogenic Tissue	312
Tumors of Vascular Tissue	312
Secondary Tumors Involving the Renal Pelvis and Ureter	319
Tumor-like Lesions of the Renal Pelvis and Ureter	320
Fibroepithelial Polyp	320
Malakoplakia	323
Retroperitoneal Fibrosis	323
Pathologic Features	327
Amyloid Tumor	327
Index	337



TUMORS OF THE KIDNEY, RENAL PELVIS, AND URETER

EMBRYOLOGY AND ANATOMY OF THE UPPER URINARY TRACT

EMBRYOGENESIS

The definitive or metanephric kidney in man has its beginnings in the second month of gestation. It is formed jointly from two different mesodermal structures; the ureteric bud, and the metanephric blastema (Hamilton et al.). The ureter, renal pelvis, renal calices, and the collecting tubules are derived from the ureteric bud, an out-

growth of the mesonephric (wolffian) duct. The nephron is composed of Bowman's capsule and glomerulus, the proximal and distal convoluted tubules and loop of Henle, all of which develop from a mesenchyme-like tissue, the metanephric blastema, located at the caudal end of the nephrogenic ridge (fig. 1).




Figure 1
DEVELOPMENT OF METANEPHRIC KIDNEY

The metanephric tissue (stipple) which originates from the caudal end of the nephrogenic ridge gives rise to the nephron. The ureter, renal pelvis, and calices develop from the metanephric duct which is an outgrowth of the mesonephric duct.

The ureteric bud stretches to reach the metanephric blastema and in the process forms the ureter. The cranial end of the bud penetrates the metanephric blastema and undergoes a succession of branching. At the origin of the first several branches, a coalescence forms the renal pelvis (fig. 2) while subsequent orders of branches form the calices and collecting tubules (fig. 3;

Osathanondh and Potter). Simultaneously, the metanephric blastema differentiates into two types of cells. The nephrogenic cells, characterized by scanty cytoplasm and prominent oval nuclei, orient themselves in compact masses around the growing ends of the collecting tubules, eventually differentiating into nephrons (fig. 4). Between the masses of nephrogenic cells




Figure 2
DEVELOPMENT OF RENAL PELVIS

Expansion of early generations of branches of the ureteral bud form the renal pelvis. The diagram represents coalescence of the third to fifth generations of branches (circled). (Fig. 9 from Osathanondh, V., and Potter, E. L. Development of human kidney as shown by microdissection. Arch. Pathol. 76:277-289, 1963.)




Figure 3
DEVELOPMENT OF RENAL CALICES AND PAPILLAE

Coalescence of the third to fifth generations of branches of the ureteral bud (circled) forms the primordial calix. (Fig. 10 from Osathanondh, V., and Potter, E. L. Development of human kidney as shown by microdissection. Arch. Pathol. 76:277-289, 1963.)

are widely spaced stromagenic cells with small naked-appearing nuclei, which will eventually give rise to the renal interstitial tissue. By the 36th week of gestation,

nephrogenesis is usually completed and the embryonic nephrogenic and stromagenic cells are no longer recognizable.




Figure 4

DIAGRAM OF DEVELOPMENT OF A METANEPHRIC KIDNEY

- a. Metanephrogenic tissue capping ampulla of collecting tubule.
- b. Enlarged blind end of ampulla.
- c. Primordium of uriniferous tubule just formed from metanephrogenic tissue.
- d. Vessel which forms the glomerulus.
- e. Bowman's capsule cut open.
- f. Uriniferous tubule in later stage of development.
- g. Collecting tubule formed from ureteric bud of the mesonephric duct.

h. Ampulla of collecting tubule cut open. (Fig. 28-22 from Bloom, W., and Fawcett, D. W. *A Textbook of Histology*, 8th ed. Philadelphia: W. B. Saunders Co., 1962. [Modified from Corning])

HISTOLOGIC STRUCTURE OF THE NEPHRON

LIGHT MICROSCOPIC FEATURES.

The glomerulus consists of a number of groups of capillary loops or lobules each arranged about a mesangial axis which merges with the media of the afferent arteriole at the hilum or vascular pole. Three cell types comprise the capillary loops: (1) A reflection of the visceral epithelium covers the urinary surface of the capillary loop and is arranged as short processes or podocytes which abut on the glomerular basement membrane; (2) a fenestrated endothelium which lines the vascular space; and (3) mesangial cells which are seen in the axial area where several capillaries are conjoined. The glomerular basement membrane consists of two immunochemically distinct basement membranes which are derived from the visceral epithelium and from the endothelium.

The juxtaglomerular apparatus (complex) consists of two elements: vascular and tubular (Barajas, 1970; 1971). The vascular component comprises a specialized secretory (endocrine) smooth muscle (the granular epithelioid cells) intercalated in the media of the afferent and occasionally efferent arterioles (Takeshita), and an extraglomerular mesangium, representing the polkissen of Zimmerman or lacis cells. The granular epithelioid cells have been shown by appropriate immunohistochemical technics to be the source of the protein, renin (Edelman and Hartroft; Hartroft et al.). The extraglomerular mesangium exhibits phagocytic activity similar to that of the glomerulus and in some species may contain granular epithelioid cells as well. It occupies the triangular space between the afferent and efferent

arterioles and the macula densa, a specialized segment of the distal convoluted tubule. The latter represents the tubular component of the juxtaglomerular apparatus (complex).

The granular epithelioid cells of the juxtaglomerular apparatus which synthesize the protein, renin, are thought to give rise to a distinctive renin-producing tumor of the kidney resembling hemangiopericytoma (Schambelan et al.; see mesenchymal tumors of the kidney on page 201). The visceral cells of Bowman's capsule have been shown recently by *in vitro* and immunohistochemical studies to be the source of erythropoietin (Busuttil et al., 1971, 1972; Burlington et al.) This new evidence suggesting synthesis of erythropoietin by the glomerular epithelial cells, developmentally a tubular portion of the nephron, may account for the production of erythropoietin by renal adenocarcinomas.

Where the proximal convoluted tubule emerges from the glomerulus, the tubular cells are similar to the parietal cells lining Bowman's capsule. In the remainder of its random tortuous course through the renal cortex, the proximal convoluted tubule is lined by a single layer of long, truncated pyramidal cells containing numerous mitochondria and abundant amounts of cytoplasm. The free surfaces of the proximal convoluted tubular cells are covered by elaborate microvilli which form the brush border seen in a light microscopic section. No such brush border is evident in the distal or collecting tubules.

The proximal convoluted tubules extend deep into the medulla and terminate as the narrowed loop of Henle. The epithelial cells of Henle's loop are squamoid in the descending portion and somewhat cuboidal in the ascending limb. After Henle's loop

returns to the cortex, the nephron continues as the distal convoluted tubule which in turn subsequently empties into one of the branches of the collecting tubules. A short specialized segment of the distal convoluted tubule comprises the macula densa, which is closely applied to the afferent arteriole and the extraglomerular mesangium. Its previously described close association with the afferent arteriole and granular epithelioid cells has recently been shown to be a less constant feature (Barajas, 1970;1971). Relative to the adjacent segments of the distal convoluted tubule, the cells of the macula densa exhibit poorly developed basal cisternae

and apical microvilli, contain fewer mitochondria, and show a more haphazard nuclear polarity. These features suggest that this specialized segment is not engaged in resorptive processes to the same degree as seen elsewhere in the distal tubule. There is a suggestion that the segment merely serves as an electrolyte leak facilitating monitoring of distal tubular sodium.

The coalescing peripheral tributaries of the collecting tubules join to form common straight collecting ducts which descend into the medulla and empty through the apex of a renal papilla into a renal calix. The microscopic appearance of the collecting tubular cells varies with the size of the tubule. In




Figure 5

GENERAL HISTOLOGIC FEATURES OF THE NEPHRON

Cross sections of the various segments of the tubule roughly indicate the cellular morphologic features and the relative size of cells and tubules at these sites. (Fig. 1-5 from Bennington, J. L., and Kradjian, R. *Renal Carcinoma*. Philadelphia: W. B. Saunders Co., 1967.)

general, the cells of the collecting tubules have sharp outlines with distinct hyperchromatic nuclei and clear pale cytoplasm. Cells of the smaller branching ducts are cuboidal, while those of the straight collecting tubules are more elongated. Surfaces of both are convex and bulge into the duct lumens (fig. 5).

ELECTRON MICROSCOPIC FEATURES. Cells of the proximal convoluted tubules have characteristic ultrastructural morphologic features which are distinctly different from those of the loop of Henle, distal convoluted tubule, and collecting ducts (Trump et al.). They are identified by their characteristic fine structure including: (1) The tall columnar shape; (2) elaborate tightly packed microvilli coated with glycocalyx; (3) pinocytotic apical vesicles, vacuoles, and tubules of characteristic structure in the apical cytoplasm; and (4) abundant elongated tortuous mitochondria intimately associated with basal and lateral cisternae (elaborate invaginations of the cell membrane) (figs. 6, 7). These features are demonstrated in the three dimensional diagrammatic reconstruction of a portion of the proximal convoluted tubule (fig. 8).

In contrast, the distal convoluted tubu-

lar cells are tall and cuboidal with scattered short microvilli (fig. 9). Extensive lateral and basal cisternae are present, but they enclose several mitochondrial profiles rather than the individual mitochondria seen in the proximal convoluted tubule cells. Apical pinocytotic vesicles and vacuoles are rare.

Cells of the loop of Henle tend to be squamoid with ovoid nuclei which have folded nuclear margins (fig. 10). Basal and lateral cell interdigitations are absent and cytoplasmic organelles are scanty.

Cells of the collecting ducts (fig. 11) are cuboidal, but become more elongated in the renal medulla. Sparse microvilli are seen on cell surfaces, but no brush border is present. The elaborate system of apical pinocytotic vesicles, tubules, and vacuoles, prominent features of proximal convoluted tubules, are absent. Mitochondria are shorter and have a more rounded configuration than those of the proximal convoluted tubular cells. They are rarely associated with the infrequent basal cisternae. Droplets of membrane-bound lipid (lipofuscin) are abundant in the basal cytoplasm.

Figure 6

PROXIMAL CONVOLUTED TUBULAR CELL

The apical part of the proximal convoluted tubular cell is covered by tightly packed microvilli forming the brush border (BB). In the apical cytoplasm, apical tubules (AT), vacuoles (V), and cytosomes (C) are evident. Mitochondria (M). Tubular Lumen (TL). Nucleus (N). X9300. (Fig. 5 from Tisher, C. C. Human renal ultrastructure. I. Proximal tubule of healthy individuals. Lab. Invest. 15:1357-1394, 1966.)