

HZ BOOKS

经 典 原 版 书 库

# 纯数学教程

(英文版·第10版)

## A COURSE OF PURE MATHEMATICS



G. H. Hardy

Cambridge Mathematical Library

(英) G. H. Hardy 著



机械工业出版社  
China Machine Press

经典原版书库

# 纯数学教程

(英文版·第10版)

A Course of Pure Mathematics  
(Tenth Edition)

(英) G. H. Hardy 著

 机械工业出版社  
China Machine Press

G. H. Hardy: *A Course of Pure Mathematics*, Tenth Edition (ISBN: 0-521-09227-2).

Originally published by Cambridge University Press in 2002.

This reprint edition is published with the permission of the Syndicate of the Press of the University of Cambridge, Cambridge, England.

Copyright © 2002 by Cambridge University Press.

This edition is licensed for distribution and sale in the People's Republic of China only, excluding Hong Kong, Taiwan and Macao and may not be distributed and sold elsewhere.

本书原版由剑桥大学出版社出版。

本书英文影印版由英国剑桥大学出版社授权出版。

此版本仅限在中华人民共和国境内（不包括中国香港、台湾、澳门地区）销售发行，未经授权的本书出口将被视为违反版权法的行为。

版权所有，侵权必究。

**本书版权登记号：图字：01-2004-0296**

### **图书在版编目（CIP）数据**

纯数学教程（英文版·第10版）/（英）哈代（Hardy, G. H.）著。—北京：机械工业出版社，2004.2

（经典原版书库）

书名原文：A Course of Pure Mathematics, Tenth Edition

ISBN 7-111-13785-X

I. 纯… II. 哈… III. 高等数学—教材—英文 IV. O13

中国版本图书馆CIP数据核字（2003）第126640号

机械工业出版社（北京市西城区百万庄大街22号 邮政编码 100037）

责任编辑：迟振春

北京瑞德印刷有限公司印刷·新华书店北京发行所发行

2004年2月第1版第1次印刷

787mm × 1092mm 1/16 · 32.5印张

印数：0 001-2 000册

定价：65.00元

凡购本书，如有倒页、脱页、缺页，由本社发行部调换

本社购书热线：(010) 68326294

## PREFACE TO THE TENTH EDITION

THE changes in the present edition are as follows:

1. An index has been added. Hardy had begun a revision of an index compiled by Professor S. Mitchell; this has been completed, as far as possible on Hardy's lines, by Dr T. M. Flett.
2. The original proof of the Heine-Borel Theorem (pp. 197-199) has been replaced by two alternative proofs due to Professor A. S. Besicovitch.
3. Example 24, p. 394 has been added to.

August, 1950

J. E. LITTLEWOOD

## PREFACE TO THE SEVENTH EDITION

THE changes in this edition are more important than in any since the second. The book has been reset, and this has given me the opportunity of altering it freely.

I have cancelled what was Appendix II (on the ' $O$ ,  $o$ ,  $\sim$ ' notation), and incorporated its contents in the appropriate places in the text. I have rewritten the parts of Chs. VI and VII which deal with the elementary properties of differential coefficients. Here I have found de la Vallée-Poussin's *Cours d'analyse* the best guide, and I am sure that this part of the book is much improved. These important changes have naturally involved many minor emendations.

I have inserted a large number of new examples from the papers for the Mathematical Tripos during the last twenty years, which should be useful to Cambridge students. These were collected for me by Mr E. R. Love, who has also read all the proofs and corrected many errors.

The general plan of the book is unchanged. I have often felt tempted, re-reading it in detail for the first time for twenty years, to make much more drastic changes both in substance and in style. It was written when analysis was neglected in Cambridge, and with an emphasis and enthusiasm which seem rather ridiculous now. If I were to rewrite it now I should not write (to use Prof. Littlewood's simile) like 'a missionary talking to cannibals', but with decent terseness and restraint; and, writing more shortly, I should be able to include a great deal more. The book would then be much more like a *Traité d'analyse* of the standard pattern.

It is perhaps fortunate that I have no time for such an undertaking, since I should probably end by writing a much better but much less individual book, and one less useful as an introduction to the books on analysis of which, even in England, there is now no lack.

November, 1937

G. H. H.

#### EXTRACT FROM THE PREFACE TO THE FIRST EDITION

THIS book has been designed primarily for the use of first year students at the Universities whose abilities reach or approach something like what is usually described as 'scholarship standard'. I hope that it may be useful to other classes of readers, but it is this class whose wants I have considered first. It is in any case a book for mathematicians: I have nowhere made any attempt to meet the needs of students of engineering or indeed any class of students whose interests are not primarily mathematical.

I regard the book as being really elementary. There are plenty of hard examples (mainly at the ends of the chapters): to these I have added, wherever space permitted, an outline of the solution. But I have done my best to avoid the inclusion of anything that involves really difficult ideas.

September, 1908

G. H. H.



# 华章教育 赤诚奉献

## 经典原版书库 (相关部分)

具体数学: 计算机科学基础 (第2版)	Graham/49.00
组合数学 (第3版)	Brualdi/35.00
离散数学及其应用 (第5版)	Rosen/79.00
离散数学及其应用 (第4版)	Rosen/59.00
高等微积分	Fitzpatrick/69.00
金融数学	Stampfli/35.00
数值分析 (第3版)	Kincaid/75.00
概率论及其在投资、保险、工程中的应用	Bean/55.00
代数	Isaacs/65.00
傅里叶分析与小波分析导论	Pinsky/49.00
数学建模 (第3版)	Giordano/59.00 (附光盘)
微分方程与边界值问题 (第5版)	Zill/69.00
应用回归分析和其他多元方法	Kleinbaum/88.00
概率统计	Stone/89.00
多元数据分析	Latin/69.00 (附光盘)
数理统计和数据分析	Rice/78.00
随机过程导论	Kao/49.00
预测与时间序列 (第3版)	Bowerman/89.00
线性代数	Jain/49.00 (附光盘)
复变函数及应用 (第7版)	Brown/42.00
实分析与复分析 (第3版)	Rudin/39.00
数学分析原理 (第3版)	Rudin/35.00
泛函分析 (第2版)	Rudin/42.00
复分析 (第3版)	Ahlfors/35.00
逼近论教程	Cheney/39.00

## 计算机科学丛书 (相关部分)

离散数学及其应用 (原书第4版)	Rosen/袁崇义/75.00
组合数学 (原书第3版)	Brualdi/冯舜玺/38.00

## 重点大学计算机教材 (相关部分)

数值方法	金一庆/25.00
------	-----------

# CONTENTS

*(Entries in small print at the end of the contents of each chapter refer to subjects discussed incidentally in the examples)*

## CHAPTER I

### REAL VARIABLES

SECT.		PAGE
1-2.	Rational numbers . . . . .	1
3-7.	Irrational numbers . . . . .	3
8.	Real numbers . . . . .	14
9.	Relations of magnitude between real numbers . . . . .	16
10-11.	Algebraical operations with real numbers . . . . .	17
12.	The number $\sqrt{2}$ . . . . .	20
13-14.	Quadratic surds . . . . .	20
15.	The continuum . . . . .	24
16.	The continuous real variable . . . . .	27
17.	Sections of the real numbers. Dedekind's theorem . . . . .	28
18.	Points of accumulation . . . . .	30
19.	Weierstrass's theorem . . . . .	31
	Miscellaneous examples . . . . .	32

Decimals, 1. Gauss's theorem, 7. Graphical solution of quadratic equations, 21. Important inequalities, 33. Arithmetical and geometrical means, 34. Cauchy's inequality, 34. Cubic and other surds, 36. Algebraical numbers, 38.

## CHAPTER II

### FUNCTIONS OF REAL VARIABLES

20.	The idea of a function . . . . .	40
21.	The graphical representation of functions. Coordinates . . . . .	43
22.	Polar coordinates . . . . .	45
23.	Polynomials . . . . .	46
24-25.	Rational functions . . . . .	49
26-27.	Algebraical functions . . . . .	52
28-29.	Transcendental functions . . . . .	55
30.	Graphical solution of equations . . . . .	60
31.	Functions of two variables and their graphical representation . . . . .	61
32.	Curves in a plane . . . . .	62
33.	Loci in space . . . . .	63
	Miscellaneous examples . . . . .	67

Trigonometrical functions, 55. Arithmetical functions, 58. Cylinders, 64. Contour maps, 64. Cones, 65. Surfaces of revolution, 65. Ruled surfaces, 66. Geometrical constructions for irrational numbers, 68. Quadrature of the circle, 70.

## CHAPTER III

## COMPLEX NUMBERS

SECT.		PAGE
34-38.	Displacements . . . . .	72
39-42.	Complex numbers . . . . .	80
43.	The quadratic equation with real coefficients . . . . .	84
44.	Argand's diagram . . . . .	87
45.	De Moivre's theorem . . . . .	88
46.	Rational functions of a complex variable . . . . .	90
47-49.	Roots of complex numbers . . . . .	101
	Miscellaneous examples . . . . .	104

Properties of a triangle, 92, 104. Equations with complex coefficients, 94. Coaxial circles, 96. Bilinear and other transformations, 97, 100, 107. Cross ratios, 99. Condition that four points should be concyclic, 100. Complex functions of a real variable, 100. Construction of regular polygons by Euclidean methods, 103. Imaginary points and lines, 106.

## CHAPTER IV

## LIMITS OF FUNCTIONS OF A POSITIVE INTEGRAL VARIABLE

50.	Functions of a positive integral variable . . . . .	110
51.	Interpolation . . . . .	111
52.	Finite and infinite classes . . . . .	112
53-57.	Properties possessed by a function of $n$ for large values of $n$ . . . . .	113
58-61.	Definition of a limit and other definitions . . . . .	120
62.	Oscillating functions . . . . .	126
63-68.	General theorems concerning limits . . . . .	129
69-70.	Steadily increasing or decreasing functions . . . . .	136
71.	Alternative proof of Weierstrass's theorem . . . . .	138
72.	The limit of $x^n$ . . . . .	139
73.	The limit of $\left(1 + \frac{1}{n}\right)^n$ . . . . .	142
74.	Some algebraical lemmas . . . . .	143
75.	The limit of $n(\sqrt[n]{x} - 1)$ . . . . .	144
76-77.	Infinite series . . . . .	145
78.	The infinite geometrical series . . . . .	149
79.	The representation of functions of a continuous real variable by means of limits . . . . .	153
80.	The bounds of a bounded aggregate . . . . .	155
81.	The bounds of a bounded function . . . . .	156
82.	The limits of indetermination of a bounded function . . . . .	156
83-84.	The general principle of convergence . . . . .	158
85-86.	Limits of complex functions and series of complex terms . . . . .	160



CONTENTS

vii

SECT.	PAGE
87-88. Applications to $z^n$ and the geometrical series . . . . .	162
89. The symbols $O, o, \sim$ . . . . .	164
Miscellaneous examples . . . . .	166
Oscillation of $\sin n\theta\pi$ , 125, 127, 158. Limits of $n^k x^n, \sqrt[n]{x}, \sqrt[n]{n}, \frac{x^n}{n!}, \sqrt[n]{(n!)}$ , $\binom{m}{n} x^n$ , 141, 144. Decimals, 149. Arithmetic series, 152. Harmonic series, 153. Equation $x_{n+1} = f(x_n)$ , 166. Limit of a mean value, 167. Expansions of rational functions, 170.	

CHAPTER V

LIMITS OF FUNCTIONS OF A CONTINUOUS VARIABLE. CONTINUOUS AND DISCONTINUOUS FUNCTIONS

90-92. Limits as $x \rightarrow \infty$ or $x \rightarrow -\infty$ . . . . .	172
93-97. Limits as $x \rightarrow a$ . . . . .	175
98. The symbols $O, o, \sim$ : orders of smallness and greatness	183
99-100. Continuous functions of a real variable . . . . .	185
101-105. Properties of continuous functions. Bounded functions. The oscillation of a function in an interval . . . . .	190
106-107. Sets of intervals on a line. The Heine-Borel theorem . . . . .	196
108. Continuous functions of several variables . . . . .	201
109-110. Implicit and inverse functions . . . . .	203
Miscellaneous examples . . . . .	206
Limits and continuity of polynomials and rational functions, 179, 187. Limit of $\frac{x^m - a^m}{x - a}$ , 181. Limit of $\frac{\sin x}{x}$ , 182. Infinity of a function, 188. Continuity of $\cos x$ and $\sin x$ , 188. Classification of discontinuities, 188. Semi-continuity, 209.	

CHAPTER VI

DERIVATIVES AND INTEGRALS

111-113. Derivatives . . . . .	210
114. General rules for differentiation . . . . .	216
115. Derivatives of complex functions . . . . .	218
116. The notation of the differential calculus . . . . .	218
117. Differentiation of polynomials . . . . .	220
118. Differentiation of rational functions . . . . .	223
119. Differentiation of algebraical functions . . . . .	224
120. Differentiation of transcendental functions . . . . .	225
121. Repeated differentiation . . . . .	228
122. General theorems concerning derivatives. Rolle's theorem . . . . .	231
123-125. Maxima and minima . . . . .	234
126-127. The mean value theorem . . . . .	242
128. Cauchy's mean value theorem . . . . .	244

SECT.	PAGE
129. A theorem of Darboux . . . . .	245
130–131. Integration. The logarithmic function . . . . .	245
132. Integration of polynomials . . . . .	249
133–134. Integration of rational functions . . . . .	250
135–142. Integration of algebraical functions. Integration by rationalisation. Integration by parts . . . . .	254
143–147. Integration of transcendental functions . . . . .	264
148. Areas of plane curves . . . . .	268
149. Lengths of plane curves . . . . .	270
Miscellaneous examples . . . . .	273
Derivative of $x^m$ , 214. Derivatives of $\cos x$ and $\sin x$ , 214. Tangent and normal to a curve, 214, 228. Multiple roots of equations, 221, 277. Rolle's theorem for polynomials, 222. Leibniz's theorem, 229. Maxima and minima of the quotient of two quadratics, 238, 277. Axes of a conic, 241. Lengths and areas in polar coordinates, 273. Differentiation of a determinant, 274. Formulae of reduction, 282.	

## CHAPTER VII

## ADDITIONAL THEOREMS IN THE DIFFERENTIAL AND INTEGRAL CALCULUS

150–151. Taylor's theorem . . . . .	285
152. Taylor's series . . . . .	291
153. Applications of Taylor's theorem to maxima and minima . . . . .	293
154. The calculation of certain limits . . . . .	293
155. The contact of plane curves . . . . .	296
156–158. Differentiation of functions of several variables . . . . .	300
159. The mean value theorem for functions of two variables . . . . .	305
160. Differentials . . . . .	307
161–162. Definite integrals . . . . .	311
163. The circular functions . . . . .	316
164. Calculation of the definite integral as the limit of a sum . . . . .	319
165. General properties of the definite integral . . . . .	320
166. Integration by parts and by substitution . . . . .	324
167. Alternative proof of Taylor's theorem . . . . .	327
168. Application to the binomial series . . . . .	328
169. Approximate formulæ for definite integrals. Simpson's rule . . . . .	328
170. Integrals of complex functions . . . . .	331
Miscellaneous examples . . . . .	332

Newton's method of approximation to the roots of equations, 288. Series for  $\cos x$  and  $\sin x$ , 292. Binomial series, 292. Tangent to a curve, 298, 310, 335. Points of inflexion, 298. Curvature, 299, 334. Osculating conics, 299, 334. Differentiation of implicit functions, 310. Maxima and minima of functions of two variables, 311. Fourier's integrals, 318, 323. The second mean value theorem, 325. Homogeneous functions, 334. Euler's theorem, 334. Jacobians, 335. Schwarz's inequality, 340.

## CHAPTER VIII

## THE CONVERGENCE OF INFINITE SERIES AND INFINITE INTEGRALS

SECT.	PAGE
171-174. Series of positive terms. Cauchy's and d'Alembert's tests of convergence . . . . .	341
175. Ratio tests . . . . .	343
176. Dirichlet's theorem . . . . .	347
177. Multiplication of series of positive terms . . . . .	347
178-180. Further tests for convergence. Abel's theorem. Mac-laurin's integral test . . . . .	349
181. The series $\sum n^{-s}$ . . . . .	352
182. Cauchy's condensation test . . . . .	354
183. Further ratio tests . . . . .	355
184-189. Infinite integrals . . . . .	356
190. Series of positive and negative terms . . . . .	371
191-192. Absolutely convergent series . . . . .	373
193-194. Conditionally convergent series . . . . .	375
195. Alternating series . . . . .	376
196. Abel's and Dirichlet's tests of convergence . . . . .	379
197. Series of complex terms . . . . .	381
198-201. Power series . . . . .	382
202. Multiplication of series . . . . .	386
203. Absolutely and conditionally convergent infinite integrals . . . . .	388
Miscellaneous examples . . . . .	390

The series  $\sum n^{k/r^n}$  and allied series, 345. Hypergeometric series, 355. Binomial series, 356, 386, 387. Transformation of infinite integrals by substitution and integration by parts, 361, 363, 369. The series  $\sum a_n \cos n\theta$ ,  $\sum a_n \sin n\theta$ , 374, 380, 381. Alteration of the sum of a series by rearrangement, 378. Logarithmic series, 385. Multiplication of conditionally convergent series, 388, 394. Recurring series, 392. Difference equations, 393. Definite integrals, 395.

## CHAPTER IX

## THE LOGARITHMIC, EXPONENTIAL, AND CIRCULAR FUNCTIONS OF A REAL VARIABLE

204-205. The logarithmic function . . . . .	398
206. The functional equation satisfied by $\log x$ . . . . .	401
207-209. The behaviour of $\log x$ as $x$ tends to infinity or to zero	402
210. The logarithmic scale of infinity . . . . .	403
211. The number $e$ . . . . .	405
212-213. The exponential function . . . . .	406
214. The general power $a^x$ . . . . .	409
215. The exponential limit . . . . .	410
216. The logarithmic limit . . . . .	411

SECT.		PAGE
217.	Common logarithms . . . . .	412
218.	Logarithmic tests of convergence . . . . .	417
219.	The exponential series . . . . .	422
220.	The logarithmic series . . . . .	425
221.	The series for arc tan $x$ . . . . .	426
222.	The binomial series . . . . .	429
223.	Alternative development of the theory . . . . .	431
224-226.	The analytical theory of the circular functions . . . . .	432
	Miscellaneous examples . . . . .	438

Integrals containing the exponential function, 413. The hyperbolic functions, 415. Integrals of certain algebraical functions, 416. Euler's constant, 420. Irrationality of  $e$ , 423. Approximation to surds by the binomial theorem, 430. Irrationality of  $\log_{10} n$ , 438. Definite integrals, 445, 446.

## CHAPTER X

### THE GENERAL THEORY OF THE LOGARITHMIC, EXPONENTIAL, AND CIRCULAR FUNCTIONS

227-228.	Functions of a complex variable . . . . .	447
229.	Curvilinear integrals . . . . .	448
230.	Definition of the logarithmic function . . . . .	449
231.	The values of the logarithmic function . . . . .	451
232-234.	The exponential function . . . . .	456
235-236.	The general power $a^x$ . . . . .	457
237-240.	The trigonometrical and hyperbolic functions . . . . .	462
241.	The connection between the logarithmic and inverse trigonometrical functions . . . . .	466
242.	The exponential series . . . . .	468
243.	The series for $\cos z$ and $\sin z$ . . . . .	469
244-245.	The logarithmic series . . . . .	471
246.	The exponential limit . . . . .	474
247.	The binomial series . . . . .	476
	Miscellaneous examples . . . . .	479

The functional equation satisfied by  $\text{Log } z$ , 454. The function  $e^z$ , 460. Logarithms to any base, 461. The inverse cosine, sine, and tangent of a complex number, 464. Trigonometrical series, 470, 472-474, 484, 485. Roots of transcendental equations, 479, 480. Transformations, 480-483. Stereographic projection, 482. Mercator's projection, 482. Level curves, 484-485. Definite integrals, 486.

APPENDIX I.	The proof that every equation has a root . . . . .	487
APPENDIX II.	A note on double limit problems . . . . .	493
APPENDIX III.	The infinite in analysis and geometry . . . . .	497
APPENDIX IV.	The infinite in analysis and geometry . . . . .	502
INDEX . . . . .		505

## CHAPTER I

### REAL VARIABLES

**1. Rational numbers.** A fraction  $r = p/q$ , where  $p$  and  $q$  are positive or negative integers, is called a *rational number*. We can suppose (i) that  $p$  and  $q$  have no common factor, since if they have a common factor we can divide each of them by it, and (ii) that  $q$  is positive, since

$$p/(-q) = (-p)/q, \quad (-p)/(-q) = p/q.$$

To the rational numbers thus defined we may add the 'rational number 0' obtained by taking  $p = 0$ .

We assume that the reader is familiar with the ordinary arithmetical rules for the manipulation of rational numbers. The examples which follow demand no knowledge beyond this.

**Examples I.** 1. If  $r$  and  $s$  are rational numbers, then  $r + s$ ,  $r - s$ ,  $rs$ , and  $r/s$  are rational numbers, unless in the last case  $s = 0$  (when  $r/s$  is of course meaningless).

2. If  $\lambda$ ,  $m$ , and  $n$  are positive rational numbers, and  $m > n$ , then  $\lambda(m^2 - n^2)$ ,  $2\lambda mn$ , and  $\lambda(m^2 + n^2)$  are positive rational numbers. Hence show how to determine any number of right-angled triangles the lengths of all of whose sides are rational.

3. Any terminated decimal represents a rational number whose denominator contains no factors other than 2 or 5. Conversely, any such rational number can be expressed, and in one way only, as a terminated decimal.

[The general theory of decimals will be considered in Ch. IV.]

4. The positive rational numbers may be arranged in the form of a simple series as follows:

$$\frac{1}{1}, \frac{2}{1}, \frac{1}{2}, \frac{3}{1}, \frac{2}{2}, \frac{1}{3}, \frac{4}{1}, \frac{3}{2}, \frac{2}{3}, \frac{1}{4}, \dots$$

Show that  $p/q$  is the  $[\frac{1}{2}(p+q-1)(p+q-2)+q]$ th term of the series.

[In this series every rational number is repeated indefinitely. Thus 1 occurs as  $\frac{1}{1}, \frac{2}{2}, \frac{3}{3}, \dots$ . We can of course avoid this by omitting every number

which has already occurred in a simpler form, but then the problem of determining the precise position of  $p/q$  becomes more complicated.]

**2. The representation of rational numbers by points on a line.** It is convenient, in many branches of mathematical analysis, to make a good deal of use of geometrical illustrations.

The use of geometrical illustrations in this way does not, of course, imply that analysis has any sort of dependence upon geometry: they are illustrations and nothing more, and are employed merely for the sake of clearness of exposition. This being so, it is not necessary that we should attempt any logical analysis of the ordinary notions of elementary geometry; we may be content to suppose, however far it may be from the truth, that we know what they mean.

Assuming, then, that we know what is meant by a *straight line*, a *segment* of a line, and the *length* of a segment, let us take a straight line  $A$ , produced indefinitely in both directions, and a segment  $A_0A_1$  of any length. We call  $A_0$  the *origin*, or *the point 0*, and  $A_1$  *the point 1*, and we regard these points as representing the numbers 0 and 1.

In order to obtain a point which shall represent a positive rational number  $r = p/q$ , we choose the point  $A_r$  such that

$$A_0A_r/A_0A_1 = r,$$

$A_0A_r$  being a stretch of the line extending in the same direction along the line as  $A_0A_1$ , a direction which we shall suppose to be from left to right when, as in Fig. 1, the line is drawn horizontally across the paper. In order to obtain a point to represent a



Fig. 1

negative rational number  $r = -s$ , it is natural to regard length as a magnitude capable of sign, positive if the length is measured in one direction (that of  $A_0A_1$ ), and negative if measured in the other, so that  $AB = -BA$ ; and to take as the point representing  $r$  the point  $A_{-s}$  such that

$$A_0A_{-s} = -A_{-s}A_0 = -A_0A_s.$$

We thus obtain a point  $A_r$  on the line corresponding to every rational value of  $r$ , positive or negative, and such that

$$A_0A_r = r \cdot A_0A_1;$$

and if, as is natural, we take  $A_0A_1$  as our unit of length, and write  $A_0A_1 = 1$ , then we have

$$A_0A_r = r.$$

We shall call the points  $A_r$  the *rational points* of the line.

**3. Irrational numbers.** If the reader will mark off on the line all the points corresponding to the rational numbers whose denominators are 1, 2, 3, ... in succession, he will readily convince himself that he can cover the line with rational points as closely as he likes. We can state this more precisely as follows: *if we take any segment  $BC$  on  $A$ , we can find as many rational points as we please on  $BC$ .*

Suppose, for example, that  $BC$  falls within the segment  $A_1A_2$ . It is evident that if we choose a positive integer  $k$  so that

$$k \cdot BC > 1 \quad \dots\dots\dots(1)*,$$

and divide  $A_1A_2$  into  $k$  equal parts, then at least one of the points of division (say  $P$ ) must fall inside  $BC$ , without coinciding with either  $B$  or  $C$ . For if this were not so,  $BC$  would be entirely included in one of the  $k$  parts into which  $A_1A_2$  has been divided, which contradicts the supposition (1). But  $P$  obviously corresponds to a rational number whose denominator is  $k$ . Thus at least one rational point  $P$  lies between  $B$  and  $C$ . But then we can find another such point  $Q$  between  $B$  and  $P$ , another between  $B$  and  $Q$ , and so on indefinitely; i.e., as we asserted above, we can find as many as we please. We may express this by saying that  $BC$  includes *infinitely many* rational points.

The meaning of such phrases as '*infinitely many*' or '*an infinity of*', in such sentences as ' $BC$  includes infinitely many rational points' or 'there are an infinity of rational points on  $BC$ ' or 'there are an infinity of positive integers', will be considered more closely in Ch. IV. The assertion 'there are an infinity of positive integers' means 'given any positive integer  $n$ ,

\* The assumption that this is possible is equivalent to the assumption of what is known as the axiom of Archimedes.

however large, we can find more than  $n$  positive integers'. This is plainly true whatever  $n$  may be, e.g. for  $n = 100,000$  or  $100,000,000$ . The assertion means exactly the same as 'we can find *as many positive integers as we please*'.

The reader will easily convince himself of the truth of the following assertion, which is substantially equivalent to what was proved in the second paragraph of this section: given any rational number  $r$ , and any positive integer  $n$ , we can find another rational number lying on either side of  $r$  and differing from  $r$  by less than  $1/n$ . It is merely to express this differently to say that we can find a rational number lying on either side of  $r$  and differing from  $r$  by *as little as we please*. Again, given any two rational numbers  $r$  and  $s$ , we can interpolate between them a chain of rational numbers in which any two consecutive terms differ by as little as we please, that is to say by less than  $1/n$ , where  $n$  is any positive integer assigned beforehand.

From these considerations the reader might be tempted to infer that an adequate view of the nature of the line could be obtained by imagining it to be formed simply by the rational points which lie on it. And it is certainly the case that if we imagine the line to be made up solely of the rational points, and all other points (if there are any such) to be eliminated, the figure which remained would possess most of the properties which common sense attributes to the straight line, and would, to put the matter roughly, look and behave very much like a line.

A little further consideration, however, shows that this view would involve us in serious difficulties.

Let us look at the matter for a moment with the eye of common sense, and consider some of the properties which we may reasonably expect a straight line to possess if it is to satisfy the idea which we have formed of it in elementary geometry.

The straight line must be composed of points, and any segment of it by all the points which lie between its end points. With any such segment must be associated a certain entity called its *length*, which must be a *quantity* capable of *numerical measurement* in terms of any standard or unit length, and these lengths must be capable of combination with one another, according to the ordinary rules of algebra, by means of addition or multiplication.



Again, it must be possible to construct a line whose length is the sum or product of any two given lengths. If the length  $PQ$ , along a given line, is  $a$ , and the length  $QR$ , along the same straight line, is  $b$ , the length  $PR$  must be  $a + b$ . Moreover, if the lengths  $OP$ ,  $OQ$ , along one straight line, are 1 and  $a$ , and the length  $OR$  along another straight line is  $b$ , and if we determine the length  $OS$  by Euclid's construction (Euc. VI. 12) for a fourth proportional to the lines  $OP$ ,  $OQ$ ,  $OR$ , this length must be  $ab$ , the algebraical fourth proportional to 1,  $a$ ,  $b$ . And it is hardly necessary to remark that the sums and products thus defined must obey the ordinary 'laws of algebra'; viz.

$$a + b = b + a, \quad a + (b + c) = (a + b) + c,$$

$$ab = ba, \quad a(bc) = (ab)c, \quad a(b + c) = ab + ac.$$

The lengths of our lines must also obey a number of obvious laws concerning inequalities as well as equalities: thus if  $A$ ,  $B$ ,  $C$  are three points lying along  $\Lambda$  from left to right, we must have  $AB < AC$ , and so on. Moreover it must be possible, on our fundamental line  $\Lambda$ , to find a point  $P$  such that  $A_0P$  is equal to any segment whatever taken along  $\Lambda$  or along any other straight line. All these properties of a line, and more, are involved in the presuppositions of our elementary geometry.

Now it is very easy to see that the idea of a straight line as composed of a series of points, each corresponding to a rational number, cannot possibly satisfy all these requirements. There




Fig. 2

are various elementary geometrical constructions, for example, which purport to construct a length  $x$  such that  $x^2 = 2$ . For instance, we may construct an isosceles right-angled triangle