

Tomoko M. Nakanishi · Keitaro Tanoi
Editors

Agricultural Implications of the Fukushima Nuclear Accident

The First Three Years



 Springer Open


Tomoko M. Nakanishi • Keitaro Tanoi
Editors

Agricultural Implications of the Fukushima Nuclear Accident

The First Three Years



Springer Open



Editors

Tomoko M. Nakanishi
Graduate School of Agricultural
and Life Sciences
The University of Tokyo
Bunkyo-ku, Tokyo
Japan

Keitaro Tanoi
Graduate School of Agricultural
and Life Sciences
The University of Tokyo
Bunkyo-ku, Tokyo
Japan

ISBN 978-4-431-55826-2
DOI 10.1007/978-4-431-55828-6

ISBN 978-4-431-55828-6 (eBook)

Library of Congress Control Number: 2015960944

Springer Tokyo Heidelberg New York Dordrecht London

© The Editor(s) (if applicable) and The Author(s) 2016. The book is published with open access at SpringerLink.com

Open Access This book is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and sources are credited.

All commercial rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer Japan KK is part of Springer Science+Business Media (www.springer.com)

Agricultural Implications of the Fukushima Nuclear Accident

Foreword

Four years have passed since the Tohoku earthquake struck on March 11, 2011. With a focus on Fukushima Prefecture, the recovery of regions that incurred damage from the nuclear power plant accident is continuing, and citizens are returning to their homes. For the agriculture, forestry, and fishing industries in Fukushima Prefecture, the nation and the municipality are utilizing methods that assist exclusion of radioactive materials, e.g., potassium fertilization. Moreover, there are adequate countermeasures against radiation exposure in Fukushima Prefecture; for example, entire crops of rice are inspected prior to shipment. Through these countermeasures, all of the products that are introduced in the market are managed so that they are below the standard limits for radionuclides. However, it seems that it will take a considerable amount of time for decontamination, for the decay of radioactive material, and for consumer anxiety about agricultural, forestry, and fishing products to be dispelled.

Immediately after the accident, the Graduate School of Agricultural and Life Sciences at The University of Tokyo created an independent team of specialists in a wide variety of areas including soil, vegetation, animal life, fishing, and forestry. This team entered the affected areas right after the accident and proceeded with research and studies. In addition, it was important to the Graduate School of Agricultural and Life Sciences that the results of this research and these studies should be useful to the recovery of the affected area; therefore, we have worked to officially announce these results. For example, there have been 11 meetings to report research results since November 2011. The objective of these meetings was to provide a simple explanation of the results of the studies and research so that the general public could understand.

Right after the first book about the research in Fukushima was published by Springer, an easy-to-understand book was published in Japanese. That book was published to allow a wide range of ordinary people to correctly understand the

impact of radioactive material on agricultural, forestry, and fishing products and the countermeasures taken against radiation exposure.


It appears that it will still take quite some time for the agricultural, forestry, and fishing industries in Fukushima Prefecture to recover, but the Graduate School of Agricultural and Life Sciences at The University of Tokyo will continue to support the recovery of these industries in the future.

Professor
Dean, Graduate School of Agricultural and Life Sciences
The University of Tokyo
Bunkyo-ku, Tokyo
Japan

Ken Furuya

Preface

More than 4 years have passed since the Fukushima Daiichi nuclear power plant accident. Even now, some 110,000 people who were evacuated from the area are in locations away from where they used to live. Right after the nuclear accident, 40–50 academic staff members of the Agricultural Department of The University of Tokyo began to study the movement of radioactive materials emitted from the nuclear reactor, because most of the contaminated area in Fukushima is related to agriculture. Researchers are still continuing their work to find out what effects the accident has had on agricultural fields. Our Graduate School of Agricultural and Life Sciences (The University of Tokyo) comprises many research areas and there are many facilities attached to the School, such as meadows, experimental forests, and farm fields. Together with these facilities, many on-site research studies have been conducted in Fukushima. The projects now ongoing can be illustrated as follows.



Through our activities many scientific findings have been collected. Based on these scientific data we started lecture classes for students as well as for the general public to explain the actual movement of radioactivity in nature, along with basic knowledge about radioactivity. So that students can experience radioecology in nature for themselves, we are periodically holding open lectures in the contaminated fields or mountains.



The first collection of papers on this subject was published by Springer Japan in 2013. It was made available as an open access book, free to download, so that the results of the research and studies could be widely shared with foreign and domestic researchers. We received an extremely large amount of feedback from that effort. There were many queries from researchers both inside and outside the country, as well as words of encouragement.

We are continuing with our studies, and we have decided to publish the present volume as a second collection of papers summarizing the subsequent research results. This collection will show more detailed results about the transfer of radioactive material into agricultural products and the movement of radioactive materials within environments like forests. We hope that this will be of use to everyone concerned in the same way that the first collection of papers was.

Tokyo, Japan

Tomoko M. Nakanishi

Contents

1	An Overview of Our Research	1
	Tomoko M. Nakanishi	
2	Monitoring Inspection for Radioactive Substances in Agricultural, Livestock, Forest and Fishery Products in Fukushima Prefecture	11
	Naoto Nihei	
3	Rice Inspections in Fukushima Prefecture	23
	Naoto Nihei	
4	Cesium Accumulation in Paddy Field Rice Grown in Fukushima from 2011 to 2013: Cultivars and Fertilization	33
	Yoshihiro Ohmori, Nobuhiro Tanaka, and Toru Fujiwara	
5	Physiological Verification of the Effect of Potassium Supply on the Reduction of Radiocesium Content in Rice Grain	45
	Natsuko I. Kobayashi	
6	Consecutive Field Trials of Rice Cultivation in Partially Decontaminated Paddy Fields to Reduce Radiocesium Absorption in the Iitate Village in Fukushima Prefecture	55
	Ichio Ii and Keitaro Tanoi	
7	Effects of “Clean Feeding” Management on Livestock Products Contaminated with Radioactive Cesium Due to the Fukushima Daiichi Nuclear Power Plant Accident	77
	Noboru Manabe, Tomotsugu Takahashi, Maiko Endo, Chunxiang Piao, Junyou Li, Hiroshi Kokado, Minoru Ohta, Keitaro Tanoi, and Tomoko M. Nakanishi	

8	Adverse Effects of Radiocesium on the Promotion of Sustainable Circular Agriculture Including Livestock Due to the Fukushima Daiichi Nuclear Power Plant Accident	91
	Noboru Manabe, Tomotsugu Takahashi, Chunxiang Piao, Junyou Li, Keitaro Tanoi, and Tomoko M. Nakanishi	
9	Wild Boars in Fukushima After the Nuclear Power Plant Accident: Distribution of Radiocesium	99
	Keitaro Tanoi	
10	Contamination of Wild Animals: Microhabitat Heterogeneity and Ecological Factors of Radioactive Cesium Exposure in Fukushima	107
	Ken Ishida	
11	Translocation of Radiocesium in Fruit Trees	119
	Daisuke Takata	
12	The Effects of Radioactive Contamination on the Forestry Industry and Commercial Mushroom-Log Production in Fukushima, Japan	145
	Satoru Miura	
13	Radiocesium in Timber of Japanese Cedar and Japanese Red Pine, in the Forests of Minamisoma, Fukushima	161
	Masaya Masumori, Norio Nogawa, Shin Sugiura, and Takeshi Tange	
14	Ecosystem Monitoring of Radiocesium Redistribution Dynamics in a Forested Catchment in Fukushima After the Nuclear Power Plant Accident in March 2011	175
	Nobuhito Ohte, Masashi Murakami, Izuki Endo, Mizue Ohashi, Kohei Iseda, Takahiro Suzuki, Tomoki Oda, Norifumi Hotta, Keitaro Tanoi, Natsuko I. Kobayashi, and Nobuyoshi Ishii	
15	Reduction of Air Radiation Dose by Ponding Paddy Fields	189
	Naritaka Kubo, Toshiaki Iida, and Masaru Mizoguchi	
16	Collaboration Structure for the Resurrection of Iitate Village, Fukushima: A Case Study of a Nonprofitable Organization	205
	Hanae Yokokawa and Masaru Mizoguchi	
17	Impacts of the Nuclear Power Plant Accident and the Start of Trial Operations in Fukushima Fisheries	217
	Nobuyuki Yagi	

18 Consumer Evaluation of Foods from the Disaster Affected Area: Change in 3 Years 229
Hiromi Hosono, Yuko Kumagai, Mami Iwabuchi,
and Tsutomu Sekizaki

19 Imaging Techniques for Radiocesium in Soil and Plants 247
Ryohei Sugita, Atsushi Hirose, Natsuko I. Kobayashi,
Keitaro Tanoi, and Tomoko M. Nakanishi

Chapter 1

An Overview of Our Research

Tomoko M. Nakanishi

Abstract The movement of radioactive Cs emitted from the Fukushima Nuclear accident has been studied by the academic staff of the Graduate School of Agricultural and Life Sciences, The University of Tokyo. The targeted items for research ranged widely, including soils, plants, animals, fish, mountains, water, etc. The relevant feature, with regard to the fallout, is that the radioactive Cs has remained at the initial contact sites and has hardly moved since. However, in the case of living individuals, such as animals, the amount of radioactivity has decreased with time at a much faster rate than the physiological half-life because of their metabolic activities. The biological half-life in animals was estimated to be within a period of 100 days. Soil plays a major role in fixing fallout. When fallout nuclides are adsorbed into the soil, plants growing there can absorb little of the radioactive Cs. In the mountains, radioactive Cs was gradually transferred from litter to soil and moved little even when washed with heavy rains. The method of contamination by radioactive nuclides is completely different from that of heavy metals.

Keywords Fukushima nuclear accident • Fallout • Radioactive Cs movement • Soil • Plant • Forest • The method of contamination

1.1 Research Project

Immediately after the Fukushima Nuclear accident, the academic staff at the Graduate School of Agricultural and Life Sciences, The University of Tokyo, organized several groups to research into the behavior of radioactive materials in the Fukushima prefecture. The researchers were divided into the following six large groups and they still continue their work today:

1. Crop plants and soils
2. Livestock and dairy products

T.M. Nakanishi (✉)

Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

e-mail: atomoko@mail.ecc.u-tokyo.ac.jp

3. Fishery
4. Environment, including wild life and forest
5. Radioactivity measurement
6. Science communication

Rice is one of the important cereal crops in Japan; however, most of the rice plants growing in the contaminated soil showed very low radioactivity in the grain, <500 Bq/kg, which was the initial regulation value. Adding K to soil proved to be an excellent and most effective way of preventing radioactive Cs uptake by plants. However, there were some exceptional cases where radioactivity in the rice grain was >100 Bq/kg, which has become the current revised regulation value for foods. Such cases are rare, and they have been studied by Prof. Keisuke Nemoto in an attempt to analyze the origin and chemical form of the radioactive Cs that is easily absorbed by rice plants in a paddy field. Rice is grown and harvested once a year; therefore, only one set of data is being collected per year. Thus, only a few data sets have been accumulated since the accident. This makes it difficult to estimate future contamination of the cereal crop until more data sets are available. In the case of crop radioactivity inspection, the Fukushima prefecture has established a system to measure the radioactivity of all the rice grains before its transfer to market. More than 10 million rice bags, each containing 30 kg of rice grain produced in Fukushima, were measured every year. The contaminated rice grain was not sold.

To measure radioactive Cs, we use pure Ge counters and Na(Tl)I counters in our radioisotope lab. An enormous number of samples of various types, mainly collected by our colleagues, were brought to this lab, and their radioactivity was measured by professional employees. Over 10,000 samples were measured per year.

Within a few months after the Fukushima Nuclear accident, most of the radioactive nuclides measured were radioactive Cs because other radioactive nuclides, such as ^{131}I , had decayed out because of their relatively short half-lives. Figure 1.1 is an example of measurements using a pure Ge counter showing the gamma-ray spectrum of rice grains. The gamma-ray energy used to calculate the nuclides ^{134}Cs , ^{137}Cs , and ^{40}K was 604.7, 661.7, and 1460.8 keV, respectively. The respective detection limits of the nuclides ^{134}Cs , ^{137}Cs , and ^{40}K were 0.7, 0.8, and 23.8 Bq/kg.

In our previous book (Nakanishi and Tanoi 2013), we reported that soil and plant contamination are different from that of animals. The features of these contaminations are summarized briefly below, with new findings that were not previously included.




Fig. 1.1 An example of gamma-ray spectrum of rice grains

1.2 Fallout on Soil, Crops, and Trees

1.2.1 Soil

The role of the soil was the most important consideration in the movement of radioactive Cs which is adsorbed into the very fine clay and organic matter. When soil from the contaminated fields was collected and radioactivity images of the fallout were taken, they showed spot-like distribution even after several months. In addition, the fallout showed no movement. The radioactive Cs was difficult to separate from the soil by chemical treatment and could only be washed from the soil during the first few weeks after fallout; the adsorption of radioactive nuclides became stronger with time, thereby making them more difficult to remove.

Prof. Sho Shiozawa has been measuring the radioactivity of the soil along with the depth. He set several pipes vertically in the soil and periodically measured the radioactivity in the pipes along with the depth of the soil. He found that the downward movement of radioactive fallout is now about 1–2 mm/year, whereas in the first 3 months after the accident it moved approximately 20 mm/3 months and then, for the following 3 months, it moved approximately 6 mm/3 months. The speed of the movement is now much slower.

Prof. Shiozawa also measured the radioactivity at the surface of the basement soil of ponds, under the water, using a special waterproof survey meter prepared by himself. The radioactivity at the surface of the soil showed little downward movement with time, similar to land soil. There are two main radioactive nuclides detectable now: ^{137}Cs and ^{134}Cs . In most cases, the radioactivity of soil under

water decreased with time, especially because of the relatively short half-life of ^{134}Cs , i.e., 2 years. The ponds he selected were collecting water from the mountains, and he recorded that little radioactive Cs was flowing into the ponds even after heavy rains.

He found only one pond where the radioactivity of the soil surface under the water was not decreased. He examined the location and environment of the pond, especially the surrounding upper mountainous area, and he found that there was a small village in close proximity to the pond where the people were always washing roads, houses, etc. using water. As a result of this activity, it seems that some of the radioactive Cs was removed from the concrete surfaces and flowed into the pond. In a few years, he will summarize and report these findings.

1.2.2 *Crops*

The Fukushima Nuclear accident occurred in March, and 2 months later, the ears of wheat crops developed and were ready for harvest. When the distribution of the radioactivity in the wheat was measured, to our great surprise, we found that it was more concentrated in the old leaves, which were exposed to the air at the time of the accident. The radioactivity of the leaves or ears that developed after the accident was comparatively very low. The fallout nuclides had hardly moved from the place where they had first landed, even after a few months.

When the radioactivity image on the leaves was magnified, the shape was still spot-like. If the radioactive Cs was incorporated into the leaves and had moved along the phloem or xylem, the vein should have been visible in the leaves. The behavior of the radioactive Cs emitted from the nuclear accident was different from the so-called macroscopic Cs chemistry we know. Because the amount of Cs deposited on leaves was so small, and most of them were carrier-free, the nuclides seem to behave like radio-colloids, or as if they were electronically adsorbed onto the tissue.

The radioactive Cs was adsorbed into the soil; therefore, it was unavailable for plant absorption. The real-time moving pictures taken by Dr. Natsuko I. Kobayashi were very convincing to many people. She grew rice plants both in water culture solution and in paddy soil which had been collected from Fukushima. Subsequently, ^{137}Cs was supplied to both the water culture solution and the paddy soil, and a comparison was made between the plants' absorption of ^{137}Cs . In water culture, the plants absorbed high amounts of ^{137}Cs within hours, and it was possible to trace its progress. Because in water culture, ^{137}Cs dissolved as an ion, it was easy for plants to absorb it. However, in the paddy soil, ^{137}Cs was trapped firmly by the soil and was unavailable to the plants. Using both ^{137}Cs and ^{42}K tracers, Dr. Kobayashi noted the effect of K on Cs uptake and the manner of translocation of Cs in the presence of K.

1.2.3 Trees

In the forests, leaves became highly contaminated at the time of the accident, and after falling to the ground and being decomposed by microorganisms, the radioactive Cs that was initially absorbed into the leaves became available to be absorbed by soil. It was found that radioactive Cs adsorption has been moving from the leaves to the soil in the forest. The radioactivity in the forest is also decreasing, along with the decay of ^{134}Cs , which has a half-life of 2 years.

However, as shown in our previous book, the radioactivity of mushrooms growing in the forest is not decreasing. Some of the mushrooms are still accumulating radioactive Cs that originated from the fallout of the nuclear test bomb during the 1960s.

In the case of fruit trees, the first question concerns the origin of the radioactive Cs in the fruit. It was taken for granted that nutrients are absorbed by roots and delivered to the whole tree. However, the fallout remained at the surface of the soil, away from the roots. Despite the active root of the peach tree being about 30 cm below the surface of the soil, where radioactive nuclides do not exist, the fruits still accumulated radioactive Cs and were slightly contaminated. Few people had considered that radioactive Cs moves into the bark from the trunk surface and could then be transferred to the fruits. Dr. Daisuke Takada performed numerous experiments to determine how radioactive Cs moves within trees, and now some of the new findings are described in this book.

1.2.4 Summary of Soil, Crop, and Tree Contamination

Most of the radioactive Cs remains adsorbed on the surface of the substances it first contacted, and the radioactivity image of radioactive Cs still maintains a spot-like shape, indicating its presence. It is very difficult to remove radioactive Cs from soil or plants (Fig. 1.2).

Below is a summary of the features of the fallout, i.e., information in the previous book:

1. The contamination in soil, crops, and trees were found as spots.
2. Emitted nuclides stay where they first landed.
3. They rarely move and stay as spots even after a few years.
4. Only a small portion of the fallout was dissolved in solution and moved.
5. The adsorption manner became stronger with time.
6. Supplying K as a fertilizer is the most effective and efficient way to prevent Cs uptake by plants.




Fig. 1.2 Features of fallout for soil, crops, and trees. Radioactive Cs was adsorbed in spot-like distribution at the surface and rarely moved with time

1.3 Fallout on Birds, Fish, and Livestock

1.3.1 Birds

In the case of birds, it was difficult to identify their point of contamination because they can fly long distances. Associate Prof. Ken Ishida caught several birds in Fukushima and compared the contamination of their feathers to those caught in other areas. A radiograph of the feathers revealed the spot-like distribution of the radioactivity. This distribution pattern was similar to that found in soil or plants. Radioactive Cs was not removed by washing with chemicals. In the following year, the same species of birds were caught and examined; however, no radioactivity was measured in their feathers. The contaminated feathers seemed to have been renewed or replaced as contaminated birds were only found in the year of the accident.