

I. S. GRADSHTEYN

I. M. RYZHIK




CD INCLUD

TABLE OF INTEGRALS, SERIES, AND PRODUCTS

积分、级数和乘积表

第 7 版

SEVENTH EDITION



Edited by Alan Jeffrey and Daniel Zwillinger



Elsevier (Singapore) Pte Ltd.

世界图书出版公司

www.wpcbj.com.cn

Table of Integrals, Series, and Products

Seventh Edition

I.S. Gradshteyn and I.M. Ryzhik

Alan Jeffrey, Editor

University of Newcastle upon Tyne, England

Daniel Zwillinger, Editor

Rensselaer Polytechnic Institute, USA

Translated from Russian by Scripta Technica, Inc.



AMSTERDAM • BOSTON • HEIDELBERG • LONDON
NEW YORK • OXFORD • PARIS • SAN DIEGO
SAN FRANCISCO • SINGAPORE • SYDNEY • TOKYO

Academic Press is an imprint of Elsevier



图书在版编目 (C I P) 数据

积分、级数和乘积表: 英文 / (美) 格里兹特恩
(Gradshteyn, I. S.), (美) 赖兹克 (Ryzhik, I. M.)
编译. —北京: 世界图书出版公司北京公司, 2007. 5
书名原文: Table of Integrals, Eries, and Products
ISBN 978-7-5062-8235-2

I. 积… II. ①格…②赖… III. ①积分—图表—英文
②级数—图表—英文③无穷乘积—图表—英文 IV.
0172.2-64 0173-64

中国版本图书馆CIP数据核字 (2007) 第060865号

书 名: Table of Integrals, Series, and Products 7th ed.

作 者: I. S. Gradshteyn, I. M. Ryzhik

中 译 名: 积分, 级数和乘积表 第7版

责任编辑: 焦小浣

出 版 者: 世界图书出版公司北京公司

印 刷 者: 北京世图印刷厂

发 行 者: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64015659, 64038348

电子信箱: kjsk@vip.sina.com

开 本: 24 开

印 张: 51

版 次: 2007 年 5 月第 1 次印刷

版权登记: 图字:01-2007-1490

书 号: 978-7-5062-8235-2 / O · 611

定 价: 178.00 元

世界图书出版公司北京公司已获得 Elsevier (Singapore) Pte Ltd. 授权在中国大陆独家重印发行

Academic Press is an imprint of Elsevier
30 Corporate Drive, Suite 400, Burlington, MA 01803, USA
525 B Street, Suite 1900, San Diego, California 92101-4495, USA
84 Theobald's Road, London WC1X 8RR, UK

This book is printed on acid-free paper. ☺

Copyright © 2007, Elsevier Inc. All rights reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Permissions may be sought directly from Elsevier's Science & Technology Rights Department in Oxford, UK: phone: (+44) 1865 843830, fax: (+44) 1865 853333, E-mail: permissions@elsevier.com. You may also complete "Support & Contact" then "Copyright and Permission" and then "Obtaining Permissions."

Library of Congress Cataloging-in-Publication Data

Application submitted

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

ISBN-13: 978-0-12-373637-6

ISBN-10: 0-12-373637-4

For information on all Elsevier Academic Press publications
visit our Web site at www.books.elsevier.com

Printed in the United States of America

07 08 09 10 11 9 8 7 6 5 4 3 2 1

Working together to grow
libraries in developing countries

www.elsevier.com | www.bookaid.org | www.sabre.org

ELSEVIER

BOOK AID
International

Sabre Foundation

Preface to the Seventh Edition

Since the publication in 2000 of the completely reset sixth edition of Gradshteyn and Ryzhik, users of the reference work have continued to submit corrections, new results that extend the work, and suggestions for changes that improve the presentation of existing entries. It is a matter of regret to us that the structure of the book makes it impossible to acknowledge these individual contributions, so, as usual, the names of the many new contributors have been added to the acknowledgment list at the front of the book.

This seventh edition contains the corrections received since the publication of the sixth edition in 2000, together with a considerable amount of new material acquired from isolated sources. Following our previous conventions, an amended entry has a superscript "11" added to its entry reference number, where the equivalent superscript number for the sixth edition was "10." Similarly, an asterisk on an entry's reference number indicates a new result. When, for technical reasons, an entry in a previous edition has been removed, to preserve the continuity of numbering between the new and older editions the subsequent entries have not been renumbered, so the numbering will jump.

We wish to express our gratitude to all who have been in contact with us with the object of improving and extending the book, and we want to give special thanks to Dr. Victor H. Moll for his interest in the book and for the many contributions he has made over an extended period of time. We also wish to acknowledge the contributions made by Dr. Francis J. O'Brien Jr. of the Naval Station in Newport, in particular for results involving integrands where exponentials are combined with algebraic functions.

Experience over many years has shown that each new edition of Gradshteyn and Ryzhik generates a fresh supply of suggestions for new entries, and for the improvement of the presentation of existing entries and errata. In view of this, we do not expect this new edition to be free from errors, so all users of this reference work who identify errors, or who wish to propose new entries, are invited to contact the authors, whose email addresses are listed below. Corrections will be posted on the web site www.az-tec.com/gr/errata.

Alan Jeffrey
Alan.Jeffrey@newcastle.ac.uk

Daniel Zwillinger
zwillinger@alum.mit.edu

Acknowledgments

The publisher and editors would like to take this opportunity to express their gratitude to the following users of the *Table of Integrals, Series, and Products* who, either directly or through errata published in *Mathematics of Computation*, have generously contributed corrections and addenda to the original printing.

Dr. A. Abbas
Dr. P. B. Abraham
Dr. Ari Abramson
Dr. Jose Adachi
Dr. R. J. Adler
Dr. N. Agmon
Dr. M. Ahmad
Dr. S. A. Ahmad
Dr. Luis Alvarez-Ruso
Dr. Maarten H P Ambaum
Dr. R. K. Amiet
Dr. L. U. Ancarani
Dr. M. Antoine
Dr. C. R. Appledorn
Dr. D. R. Appleton
Dr. Mitsuhiro Arikawa
Dr. P. Ashoshauvati
Dr. C. L. Axness
Dr. E. Badralexe
Dr. S. B. Bagchi
Dr. L. J. Baker
Dr. R. Ball
Dr. M. P. Barnett
Dr. Florian Baumann
Dr. Norman C. Beaulieu
Dr. Jerome Benoit
Mr. V. Bentley
Dr. Laurent Berger
Dr. M. van den Berg
Dr. N. F. Berk
Dr. C. A. Bertulani

Dr. J. Betancort-Rijo
Dr. P. Bickerstaff
Dr. Iwo Bialynicki-Birula
Dr. Chris Bidinosti
Dr. G. R. Bigg
Dr. Ian Bindloss
Dr. L. Blanchet
Dr. Mike Blaskiewicz
Dr. R. D. Blevins
Dr. Anders Blom
Dr. L. M. Blumberg
Dr. R. Blumel
Dr. S. E. Bodner
Dr. M. Bonsager
Dr. George Boros
Dr. S. Bosanac
Dr. B. Van den Bossche
Dr. A. Boström
Dr. J. E. Bowcock
Dr. T. H. Boyer
Dr. K. M. Briggs
Dr. D. J. Broadhurst
Dr. Chris Van Den Broeck
Dr. W. B. Brower
Dr. H. N. Browne
Dr. Christoph Bruegger
Dr. William J. Bruno
Dr. Vladimir Bubanja
Dr. D. J. Buch
Dr. D. J. Bukman
Dr. F. M. Burrows

Dr. R. Caboz
Dr. T. Calloway
Dr. F. Calogero
Dr. D. Dal Cappello
Dr. David Cardon
Dr. J. A. Carlson Gallos
Dr. B. Carrascal
Dr. A. R. Carr
Dr. S. Carter
Dr. G. Cavalleri
Mr. W. H. L. Cawthorne
Dr. A. Cecchini
Dr. B. Chan
Dr. M. A. Chaudhry
Dr. Sabino Chavez-Cerda
Dr. Julian Cheng
Dr. H. W. Chew
Dr. D. Chin
Dr. Young-seek Chung
Dr. S. Ciccariello
Dr. N. S. Clarke
Dr. R. W. Cleary
Dr. A. Clement
Dr. P. Cochrane
Dr. D. K. Cohoon
Dr. L. Cole
Dr. Filippo Colomo
Dr. J. R. D. Copley
Dr. D. Cox
Dr. J. Cox
Dr. J. W. Criss

-
- | | | |
|--------------------------|-----------------------------|-----------------------------|
| Dr. A. E. Curzon | Dr. J. France | Dr. D. L. Gunter |
| Dr. D. Dadyburjor | Dr. B. Frank | Dr. Julio C. Gutiérrez-Vega |
| Dr. D. Dajaputra | Dr. S. Frasier | Dr. Roger Haagmans |
| Dr. C. Dal Cappello | Dr. Stefan Fredenhagen | Dr. H. van Haeringen |
| Dr. P. Daly | Dr. A. J. Freeman | Dr. B. Hafizi |
| Dr. S. Dasgupta | Dr. A. Frink | Dr. Bahman Hafizi |
| Dr. John Davies | Dr. Jason M. Gallaspy | Dr. T. Hagfors |
| Dr. C. L. Davis | Dr. J. A. C. Gallas | Dr. M. J. Haggerty |
| Dr. A. Degasperis | Dr. J. A. Carlson Gallas | Dr. Timo Hakulinen |
| Dr. B. C. Denardo | Dr. G. R. Gamertsfelder | Dr. Einar Halvorsen |
| Dr. R. W. Dent | Dr. T. Garavaglia | Dr. S. E. Hammel |
| Dr. E. Deutsch | Dr. Jaime Zaratiegui Garcia | Dr. E. Hansen |
| Dr. D. deVries | Dr. C. G. Gardner | Dr. Wes Harker |
| Dr. P. Dita | Dr. D. Garfinkle | Dr. T. Harrett |
| Dr. P. J. de Doelder | Dr. P. N. Garner | Dr. D. O. Harris |
| Dr. Misha Dohler | Dr. F. Gasser | Dr. Frank Harris |
| Dr. G. Dôme | Dr. E. Gath | Mr. Mazen D. Hasna |
| Dr. Shi-Hai Dong | Dr. P. Gatt | Dr. Joel G. Heinrich |
| Dr. Balazs Dora | Dr. D. Gay | Dr. Sten Herlitz |
| Dr. M. R. D'Orsogna | Dr. M. P. Gelfand | Dr. Chris Herzog |
| Dr. Adrian A. Dragulescu | Dr. M. R. Geller | Dr. A. Higuchi |
| Dr. Eduardo Duenez | Dr. Ali I. Genc | Dr. R. E. Hise |
| Mr. Tommi J. Dufva | Dr. Vincent Genot | Dr. Henrik Holm |
| Dr. E. B. Dussan, V | Dr. M. F. George | Dr. Helmut Hölzler |
| Dr. C. A. Ebner | Dr. P. Germain | Dr. N. Holte |
| Dr. M. van der Ende | Dr. Ing. Christoph Gierull | Dr. R. W. Hopper |
| Dr. Jonathan Engle | Dr. S. P. Gill | Dr. P. N. Houle |
| Dr. G. Eng | Dr. Federico Giroi | Dr. C. J. Howard |
| Dr. E. S. Erck | Dr. E. A. Gislason | Dr. J. H. Hubbell |
| Dr. Jan Erkelens | Dr. M. I. Glasser | Dr. J. R. Hull |
| Dr. Olivier Espinosa | Dr. P. A. Glendinning | Dr. W. Humphries |
| Dr. G. A. Estévez | Dr. L. I. Goldfischer | Dr. Jean-Marc Huré |
| Dr. K. Evans | Dr. Denis Golosov | Dr. Ben Yu-Kuang Hu |
| Dr. G. Evendon | Dr. I. J. Good | Dr. Y. Iksbe |
| Dr. V. I. Fabrikant | Dr. J. Good | Dr. Philip Ingenhoven |
| Dr. L. A. Falkovsky | Mr. L. Gorin | Mr. L. Iossif |
| Dr. K. Farahmand | Dr. Martin Götz | Dr. Sean A. Irvine |
| Dr. Richard J. Fateman | Dr. R. Govindaraj | Dr. Öttar Ísberg |
| Dr. G. Fedele | Dr. M. De Grauf | Dr. Cyril-Daniel Iskander |
| Dr. A. R. Ferchmin | Dr. L. Green | Dr. S. A. Jackson |
| Dr. P. Ferrant | Mr. Leslie O. Green | Dr. John David Jackson |
| Dr. H. E. Fettis | Dr. R. Greenwell | Dr. Francois Jaclot |
| Dr. W. B. Fichter | Dr. K. D. Grimsley | Dr. B. Jacobs |
| Dr. George Fikioris | Dr. Albert Groenenboom | Dr. E. C. James |
| Mr. J. C. S. S. Filho | Dr. V. Gudmundsson | Dr. B. Jancovici |
| Dr. L. Ford | Dr. J. Guillera | Dr. D. J. Jeffrey |
| Dr. Nicolao Fornengo | Dr. K. Gunn | Dr. H. J. Jensen |

-
- | | | |
|---------------------------|-------------------------|----------------------------|
| Dr. Edwin F. Johnson | Dr. Todd Lee | Dr. D. L. Miller |
| Dr. I. R. Johnson | Dr. J. Legg | Dr. Steve Miller |
| Dr. Steven Johnson | Dr. Armando Lemus | Dr. P. C. D. Milly |
| Dr. I. Johnstone | Dr. S. L. Levie | Dr. S. P. Mitra |
| Dr. Y. P. Joshi | Dr. D. Levi | Dr. K. Miura |
| Dr. Jae-Hun Jung | Dr. Michael Lexa | Dr. N. Mohankumar |
| Dr. Damir Juric | Dr. Kuo Kan Liang | Dr. M. Moll |
| Dr. Florian Kaempfer | Dr. B. Linet | Dr. Victor H. Moll |
| Dr. S. Kanmani | Dr. M. A. Lisa | Dr. D. Monowalow |
| Dr. Z. Kapal | Dr. Donald Livesay | Mr. Tony Montagnese |
| Dr. Dave Kasper | Dr. H. Li | Dr. Jim Morehead |
| Dr. M. Kaufman | Dr. Georg Lohoefer | Dr. J. Morice |
| Dr. B. Kay | Dr. I. M. Longman | Dr. W. Mueck |
| Dr. Avinash Khare | Dr. D. Long | Dr. C. Muhlhausen |
| Dr. Ilki Kim | Dr. Sylvie Lorthois | Dr. S. Mukherjee |
| Dr. Youngsun Kim | Dr. Y. L. Luke | Dr. R. R. Müller |
| Dr. S. Klama | Dr. W. Lukosz | Dr. Pablo Parmezani Munhoz |
| Dr. L. Klingen | Dr. T. Lundgren | Dr. Paul Nanninga |
| Dr. C. Knessl | Dr. E. A. Luraev | Dr. A. Natarajan |
| Dr. M. J. Knight | Dr. R. Lynch | Dr. Stefan Neumeier |
| Dr. Mel Knight | Dr. R. Mahurin | Dr. C. T. Nguyen |
| Dr. Yannis Kohninos | Dr. R. Mallier | Dr. A. C. Nicol |
| Dr. D. Koks | Dr. G. A. Mamon | Dr. M. M. Nieto |
| Dr. L. P. Kok | Dr. A. Mangiarotti | Dr. P. Noerdlinger |
| Dr. K. S. Kölbig | Dr. I. Manning | Dr. A. N. Norris |
| Dr. Y. Komninos | Dr. J. Marmur | Dr. K. H. Norwich |
| Dr. D. D. Konowalow | Dr. A. Martin | Dr. A. H. Nuttall |
| Dr. Z. Kopal | Sr. Yuzo Maruyama | Dr. Frank O'Brien |
| Dr. I. Kostyukov | Dr. David J. Masiello | Dr. R. P. O'Keeffe |
| Dr. R. A. Krajcik | Dr. Richard Marthar | Dr. A. Ojo |
| Dr. Vincent Krakoviack | Dr. H. A. Mavromatis | Dr. P. Olsson |
| Dr. Stefan Kramer | Dr. M. Mazzoni | Dr. M. Ortner |
| Dr. Tobias Kramer | Dr. K. B. Ma | Dr. S. Ostlund |
| Dr. Hermann Krebs | Dr. P. McCullagh | Dr. J. Overduin |
| Dr. J. W. Krozel | Dr. J. H. McDonnell | Dr. J. Pachner |
| Dr. E. D. Krupnikov | Dr. J. R. McGregor | Dr. John D. Paden |
| Dr. Kun-Lin Kuo | Dr. Kim McInturff | Mr. Robert A. Padgug |
| Dr. E. A. Kuraev | Dr. N. McKinney | Dr. D. Papadopoulos |
| Dr. Konstantinos Kyritsis | Dr. David McA McKirdy | Dr. F. J. Papp |
| Dr. Velimir Labinac | Dr. Rami Mehrem | Mr. Man Sik Park |
| Dr. A. D. J. Lambert | Dr. W. N. Mei | Dr. Jong-Do Park |
| Dr. A. Lambert | Dr. Angelo Melino | Dr. B. Patterson |
| Dr. A. Larraza | Mr. José Ricardo Mendes | Dr. R. F. Pawula |
| Dr. K. D. Lee | Dr. Andy Mennim | Dr. D. W. Peaceman |
| Dr. M. Howard Lee | Dr. J. P. Meunier | Dr. D. Pelat |
| Dr. M. K. Lee | Dr. Gerard P. Michon | Dr. L. Peliti |
| Dr. P. A. Lee | Dr. D. F. R. Mildner | Dr. Y. P. Pellegrini |

-
- | | | |
|------------------------------|-----------------------------|--------------------------|
| Dr. G. J. Pert | Dr. Kazuhiko Seki | Dr. Ming Tsai |
| Dr. Nicola Pessina | Dr. B. Seshadri | Dr. N. Turkkan |
| Dr. J. B. Peterson | Dr. A. Shapiro | Dr. Sandeep Tyagi |
| Dr. Rickard Petersson | Dr. Masaki Shigemori | Dr. J. J. Tyson |
| Dr. Andrew Plumb | Dr. J. S. Sheng | Dr. S. Uehara |
| Dr. Dror Porat | Dr. Kenneth Ing Shing | Dr. M. Vadacchino |
| Dr. E. A. Power | Dr. Tomohiro Shirai | Dr. O. T. Valls |
| Dr. E. Predazzi | Dr. S. Shlomo | Dr. D. Vandeth |
| Dr. William S. Price | Dr. D. Siegel | Mr. Andras Vanyolos |
| Dr. Paul Radmore | Dr. Matthew Stapleton | Dr. D. Veitch |
| Dr. F. Raynal | Dr. Steven H. Simon | Mr. Jose Lopez Vicario |
| Dr. X. R. Resende | Dr. Ashok Kumar Singal | Dr. K. Vogel |
| Dr. J. M. Riedler | Dr. C. Smith | Dr. J. M. M. J. Vogels |
| Dr. Thomas Richard | Dr. G. C. C. Smith | Dr. Alexis De Vos |
| Dr. E. Ringel | Dr. Stefan Llewellyn Smith | Dr. Stuart Walsh |
| Dr. T. M. Roberts | Dr. S. Smith | Dr. Reinhold Wannemacher |
| Dr. N. I. Robinson | Dr. G. Solt | Dr. S. Wanzura |
| Dr. P. A. Robinson | Dr. J. Sondow | Dr. J. Ward |
| Dr. D. M. Rosenblum | Dr. A. Sørenssen | Dr. S. I. Warshaw |
| Dr. R. A. Rosthal | Dr. Marcus Spradlin | Dr. R. Weber |
| Dr. J. R. Roth | Dr. Andrzej Staruszkiewicz | Dr. Wei Qian |
| Dr. Klaus Rottbrand | Dr. Philip C. L. Stephenson | Dr. D. H. Werner |
| Dr. D. Roy | Dr. Edgardo Stockmeyer | Dr. E. Wetzel |
| Dr. E. Royer | Dr. J. C. Straton | Dr. Robert Whittaker |
| Dr. D. Rudermann | Mr. H. Suraweera | Dr. D. T. Wilton |
| Dr. Sanjib Sabhapandit | Dr. N. F. Svaiter | Dr. C. Wiuf |
| Dr. C. T. Sachradja | Dr. V. Svaiter | Dr. K. T. Wong |
| Dr. J. Sadiku | Dr. R. Szymtkowski | Mr. J. N. Wright |
| Dr. A. Sadiq | Dr. S. Tabachnik | Dr. J. D. Wright |
| Dr. Motohiko Saitoh | Dr. Erik Talvila | Dr. D. Wright |
| Dr. Naoki Saito | Dr. G. Tanaka | Dr. D. Wu |
| Dr. A. Salim | Dr. C. Tanguy | Dr. Michel Daoud Yacoub |
| Dr. J. H. Samson | Dr. G. K. Tannahill | Dr. Yu S. Yakovlev |
| Dr. Miguel A. Sanchis-Lozano | Dr. B. T. Tan | Dr. H.-C. Yang |
| Dr. J. A. Sanders | Dr. C. Tavad | Dr. J. J. Yang |
| Dr. M. A. F. Sanjun | Dr. Gonçalo Tavares | Dr. Z. J. Yang |
| Dr. P. Sarquiz | Dr. Aba Teleki | Dr. J. J. Wang |
| Dr. Avadh Saxena | Dr. Arash Dahi Taleghani | Dr. Peter Widerin |
| Dr. Vito Scarola | Dr. D. Temperley | Mr. Chun Kin Au Yeung |
| Dr. O. Schärpf | Dr. A. J. Tervoort | Dr. Kazuya Yuasa |
| Dr. A. Scherzinger | Dr. Theodoros Theodoulidis | Dr. S. P. Yukon |
| Dr. B. Schizer | Dr. D. J. Thomas | Dr. B. Zhang |
| Dr. Martin Schmid | Dr. Michael Thorwart | Dr. Y. C. Zhang |
| Dr. J. Scholes | Dr. S. T. Thynell | Dr. Y. Zhao |
| Dr. Mel Schopper | Dr. D. C. Torney | Dr. Ralf Zimmer |
| Dr. H. J. Schulz | Dr. R. Tough | |
| Dr. G. J. Sears | Dr. B. F. Treadway | |

The Order of Presentation of the Formulas

The question of the most expedient order in which to give the formulas, in particular, in what division to include particular formulas such as the definite integrals, turned out to be quite complicated. The thought naturally occurs to set up an order analogous to that of a dictionary. However, it is almost impossible to create such a system for the formulas of integral calculus. Indeed, in an arbitrary formula of the form

$$\int_a^b f(x) dx = A$$

one may make a large number of substitutions of the form $x = \varphi(t)$ and thus obtain a number of "synonyms" of the given formula. We must point out that the table of definite integrals by Bierens de Haan and the earlier editions of the present reference both sin in the plethora of such "synonyms" and formulas of complicated form. In the present edition, we have tried to keep only the simplest of the "synonym" formulas. Basically, we judged the simplicity of a formula from the standpoint of the simplicity of the arguments of the "outer" functions that appear in the integrand. Where possible, we have replaced a complicated formula with a simpler one. Sometimes, several complicated formulas were thereby reduced to a single, simpler one. We then kept only the simplest formula. As a result of such substitutions, we sometimes obtained an integral that could be evaluated by use of the formulas of Chapter Two and the Newton-Leibniz formula, or to an integral of the form

$$\int_{-a}^a f(x) dx,$$

where $f(x)$ is an odd function. In such cases, the complicated integrals have been omitted.

Let us give an example using the expression

$$\int_0^{\pi/4} \frac{(\cot x - 1)^{p-1}}{\sin^2 x} \ln \tan x dx = -\frac{\pi}{p} \operatorname{cosec} p\pi. \quad (0.1)$$

By making the natural substitution $u = \cot x - 1$, we obtain

$$\int_0^\infty u^{p-1} \ln(1+u) du = \frac{\pi}{p} \operatorname{cosec} p\pi. \quad (0.2)$$

Integrals similar to formula (0.1) are omitted in this new edition. Instead, we have formula (0.2).

As a second example, let us take

$$I = \int_0^{\pi/2} \ln(\tan^p x + \cot^p x) \ln \tan x \, dx = 0.$$

The substitution $u = \tan x$ yields

$$I = \int_0^\infty \frac{\ln(u^p + u^{-p}) \ln u}{1 + u^2} \, du.$$

If we now set $v = \ln u$, we obtain

$$I = \int_{-\infty}^\infty \frac{ve^v}{1 + e^{2v}} \ln(e^{pv} + e^{-pv}) \, dv = \int_{-\infty}^\infty v \frac{\ln(2 \cosh pv)}{2 \cosh v} \, dv.$$

The integrand is odd, and, consequently, the integral is equal to 0.

Thus, before looking for an integral in the tables, the user should simplify as much as possible the arguments (the "inner" functions) of the functions in the integrand.

The functions are ordered as follows: First we have the elementary functions:

1. The function $f(x) = x$.
2. The exponential function.
3. The hyperbolic functions.
4. The trigonometric functions.
5. The logarithmic function.
6. The inverse hyperbolic functions. (These are replaced with the corresponding logarithms in the formulas containing definite integrals.)
7. The inverse trigonometric functions.

Then follow the special functions:

8. Elliptic integrals.
9. Elliptic functions.
10. The logarithm integral, the exponential integral, the sine integral, and the cosine integral functions.
11. Probability integrals and Fresnel's integrals.
12. The gamma function and related functions.
13. Bessel functions.
14. Mathieu functions.
15. Legendre functions.
16. Orthogonal polynomials.
17. Hypergeometric functions.
18. Degenerate hypergeometric functions.
19. Parabolic cylinder functions.
20. Meijer's and MacRobert's functions.
21. Riemann's zeta function.

The integrals are arranged in order of outer function according to the above scheme: the farther down in the list a function occurs, (i.e., the more complex it is) the later will the corresponding formula appear

in the tables. Suppose that several expressions have the same outer function. For example, consider $\sin e^x$, $\sin x$, $\sin \ln x$. Here, the outer function is the sine function in all three cases. Such expressions are then arranged in order of the inner function. In the present work, these functions are therefore arranged in the following order: $\sin x$, $\sin e^x$, $\sin \ln x$.

Our list does not include polynomials, rational functions, powers, or other algebraic functions. An algebraic function that is included in tables of definite integrals can usually be reduced to a finite combination of roots of rational power. Therefore, for classifying our formulas, we can conditionally treat a power function as a generalization of an algebraic and, consequently, of a rational function.* We shall distinguish between all these functions and those listed above, and we shall treat them as operators. Thus, in the expression $\sin^2 e^x$, we shall think of the squaring operator as applied to the outer function, namely, the sine. In the expression $\frac{\sin x + \cos x}{\sin x - \cos x}$, we shall think of the rational operator as applied to the trigonometric functions sine and cosine. We shall arrange the operators according to the following order:

1. Polynomials (listed in order of their degree).
2. Rational operators.
3. Algebraic operators (expressions of the form $A^{p/q}$, where q and p are rational, and $q > 0$; these are listed according to the size of q).
4. Power operators.

Expressions with the same outer and inner functions are arranged in the order of complexity of the operators. For example, the following functions [whose outer functions are all trigonometric, and whose inner functions are all $f(x) = x$] are arranged in the order shown:

$$\sin x, \quad \sin x \cos x, \quad \frac{1}{\sin x} = \operatorname{cosec} x, \quad \frac{\sin x}{\cos x} = \tan x, \quad \frac{\sin x + \cos x}{\sin x - \cos x}, \quad \sin^m x, \quad \sin^m x \cos x.$$

Furthermore, if two outer functions $\varphi_1(x)$ and $\varphi_2(x)$, where $\varphi_1(x)$ is more complex than $\varphi_2(x)$, appear in an integrand and if any of the operations mentioned are performed on them, the corresponding integral will appear [in the order determined by the position of $\varphi_2(x)$ in the list] after all integrals containing only the function $\varphi_1(x)$. Thus, following the trigonometric functions are the trigonometric and power functions [that is, $\varphi_2(x) = x$]. Then come

- combinations of trigonometric and exponential functions,
- combinations of trigonometric functions, exponential functions, and powers, etc.,
- combinations of trigonometric and hyperbolic functions, etc.

Integrals containing two functions $\varphi_1(x)$ and $\varphi_2(x)$ are located in the division and order corresponding to the more complicated function of the two. However, if the positions of several integrals coincide because they contain the same complicated function, these integrals are put in the position defined by the complexity of the second function.

To these rules of a general nature, we need to add certain particular considerations that will be easily understood from the tables. For example, according to the above remarks, the function $e^{\frac{1}{x}}$ comes after e^x as regards complexity, but $\ln x$ and $\ln \frac{1}{x}$ are equally complex since $\ln \frac{1}{x} = -\ln x$. In the section on "powers and algebraic functions," polynomials, rational functions, and powers of powers are formed from power functions of the form $(a + bx)^n$ and $(\alpha + \beta x)^\nu$.

*For any natural number n , the involution $(a + bx)^n$ of the binomial $a + bx$ is a polynomial. If n is a negative integer, $(a + bx)^n$ is a rational function. If n is irrational, the function $(a + bx)^n$ is not even an algebraic function.

Use of the Tables*

For the effective use of the tables contained in this book, it is necessary that the user should first become familiar with the classification system for integrals devised by the authors Ryzhik and Gradshteyn. This classification is described in detail in the section entitled *The Order of Presentation of the Formulas* (see page xxvii) and essentially involves the separation of the integrand into *inner* and *outer* functions. The principal function involved in the integrand is called the *outer* function; and its argument, which is itself usually another function, is called the *inner* function. Thus, if the integrand comprised the expression $\ln \sin x$, the *outer* function would be the logarithmic function while its argument, the *inner* function, would be the trigonometric function $\sin x$. The desired integral would then be found in the section dealing with logarithmic functions, its position within that section being determined by the position of the *inner* function (here a trigonometric function) in Gradshteyn and Ryzhik's list of functional forms.

It is inevitable that some duplication of symbols will occur within such a large collection of integrals, and this happens most frequently in the first part of the book dealing with algebraic and trigonometric integrands. The symbols most frequently involved are α , β , γ , δ , t , u , z , z_k , and Δ . The expressions associated with these symbols are used consistently within each section and are defined at the start of each new section in which they occur. Consequently, reference should be made to the beginning of the section being used in order to verify the meaning of the substitutions involved.

Integrals of algebraic functions are expressed as combinations of roots with rational power indices, and definite integrals of such functions are frequently expressed in terms of the Legendre elliptic integrals $F(\phi, k)$, $E(\phi, k)$ and $\Pi(\phi, n, k)$, respectively, of the first, second, and third kinds.

The four inverse hyperbolic functions $\operatorname{arcsinh} z$, $\operatorname{arcosh} z$, $\operatorname{arctanh} z$, and $\operatorname{arcoth} z$ are introduced through the definitions

$$\begin{aligned}\operatorname{arcsin} z &= \frac{1}{i} \operatorname{arcsinh}(iz) \\ \operatorname{arccos} z &= \frac{1}{i} \operatorname{arccosh}(z) \\ \operatorname{arctan} z &= \frac{1}{i} \operatorname{arctanh}(iz) \\ \operatorname{arccot} z &= i \operatorname{arcoth}(iz)\end{aligned}$$

*Prepared by Alan Jeffrey for the English language edition.

or

$$\begin{aligned}\operatorname{arcsinh} z &= \frac{1}{i} \operatorname{arcsin}(iz) \\ \operatorname{arccosh} z &= i \operatorname{arccos} z \\ \operatorname{arctanh} z &= \frac{1}{i} \operatorname{arctan}(iz) \\ \operatorname{arccoth} z &= \frac{1}{i} \operatorname{arccot}(-iz)\end{aligned}$$

The numerical constants C and G which often appear in the definite integrals denote Euler's constant and Catalan's constant, respectively. Euler's constant C is defined by the limit

$$C = \lim_{s \rightarrow \infty} \left(\sum_{n=1}^s \frac{1}{n} - \ln s \right) = 0.577215 \dots$$

On occasion, other writers denote Euler's constant by the symbol γ , but this is also often used instead to denote the constant

$$\gamma = e^C = 1.781072 \dots$$

Catalan's constant G is related to the complete elliptic integral

$$K \equiv K(k) \equiv \int_0^{\pi/2} \frac{dx}{\sqrt{1 - k^2 \sin^2 x}}$$

by the expression

$$G = \frac{1}{2} \int_0^1 K dk = \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)^2} = 0.915965 \dots$$

Since the notations and definitions for higher transcendental functions that are used by different authors are by no means uniform, it is advisable to check the definitions of the functions that occur in these tables. This can be done by identifying the required function by symbol and name in the *Index of Special Functions and Notation* on page xxxix, and by then referring to the defining formula or section number listed there. We now present a brief discussion of some of the most commonly used alternative notations and definitions for higher transcendental functions.

Bernoulli and Euler Polynomials and Numbers

Extensive use is made throughout the book of the Bernoulli and Euler numbers B_n and E_n that are defined in terms of the Bernoulli and Euler polynomials of order n , $B_n(x)$ and $E_n(x)$, respectively. These polynomials are defined by the generating functions

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad \text{for } |t| < 2\pi$$

and

$$\frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \quad \text{for } |t| < \pi.$$

The Bernoulli numbers are always denoted by B_n and are defined by the relation

$$B_n = B_n(0) \quad \text{for } n = 0, 1, \dots,$$

when

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \dots$$

The Euler numbers E_n are defined by setting

$$E_n = 2^n E_n \left(\frac{1}{2} \right) \quad \text{for } n = 0, 1, \dots$$

The E_n are all integral, and $E_0 = 1$, $E_2 = -1$, $E_4 = 5$, $E_6 = -61$,

An alternative definition of Bernoulli numbers, which we shall denote by the symbol B_n^* , uses the same generating function but identifies the B_n^* differently in the following manner:

$$\frac{t}{e^t - 1} = 1 - \frac{1}{2}t + B_1^* \frac{t^2}{2!} - B_2^* \frac{t^4}{4!} + \dots$$

This definition then gives rise to the alternative set of Bernoulli numbers

$$\begin{aligned} B_1^* &= 1/6, & B_2^* &= 1/30, & B_3^* &= 1/42, & B_4^* &= 1/30, & B_5^* &= 5/66, \\ B_6^* &= 691/2730, & B_7^* &= 7/6, & B_8^* &= 3617/510, & \dots \end{aligned}$$

These differences in notation must also be taken into account when using the following relationships that exist between the Bernoulli and Euler polynomials:

$$B_n(x) = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} B_{n-k} E_k(2x) \quad n = 0, 1, \dots$$

$$E_{n-1}(x) = \frac{2^n}{n} \left\{ B_n \left(\frac{x+1}{2} \right) - B_n \left(\frac{x}{2} \right) \right\}$$

or

$$E_{n-1}(x) = \frac{2}{n} \left\{ B_n(x) - 2^n B_n \left(\frac{x}{2} \right) \right\} \quad n = 1, 2, \dots$$

and

$$E_{n-2}(x) = 2 \binom{n}{2}^{-1} \sum_{k=0}^{n-2} \binom{n}{k} (2^{n-k} - 1) B_{n-k} B_n(x) \quad n = 2, 3, \dots$$

There are also alternative definitions of the Euler polynomial of order n , and it should be noted that some authors, using a modification of the third expression above, call

$$\left(\frac{2}{n+1} \right) \left\{ B_n(x) - 2^n B_n \left(\frac{x}{2} \right) \right\}$$

the Euler polynomial of order n .

Elliptic Functions and Elliptic Integrals

The following notations are often used in connection with the inverse elliptic functions $\operatorname{sn} u$, $\operatorname{cn} u$, and $\operatorname{dn} u$:

$$\operatorname{ns} u = \frac{1}{\operatorname{sn} u}$$

$$\operatorname{sc} u = \frac{\operatorname{sn} u}{\operatorname{cn} u}$$

$$\operatorname{sd} u = \frac{\operatorname{sn} u}{\operatorname{dn} u}$$

$$\operatorname{nc} u = \frac{1}{\operatorname{cn} u}$$

$$\operatorname{cs} u = \frac{\operatorname{cn} u}{\operatorname{sn} u}$$

$$\operatorname{cd} u = \frac{\operatorname{cn} u}{\operatorname{dn} u}$$

$$\operatorname{nd} u = \frac{1}{\operatorname{dn} u}$$

$$\operatorname{ds} u = \frac{\operatorname{dn} u}{\operatorname{sn} u}$$

$$\operatorname{dc} u = \frac{\operatorname{dn} u}{\operatorname{cn} u}$$

The elliptic integral of the third kind is defined by Gradshteyn and Ryzhik to be

$$\begin{aligned}\Pi(\varphi, n^2, k) &= \int_0^\varphi \frac{da}{(1 - n^2 \sin^2 a) \sqrt{1 - k^2 \sin^2 a}} \\ &= \int_0^{\sin \varphi} \frac{dx}{(1 - n^2 x^2) \sqrt{(1 - x^2)(1 - k^2 x^2)}}\end{aligned}\quad (-\infty < n^2 < \infty)$$

The Jacobi Zeta Function and Theta Functions

The Jacobi zeta function $\text{zn}(u, k)$, frequently written $Z(u)$, is defined by the relation

$$\text{zn}(u, k) = Z(u) = \int_0^u \left\{ \text{dn}^2 v - \frac{E}{K} \right\} dv = E(u) - \frac{E}{K} u.$$

This is related to the theta functions by the relationship

$$\text{zn}(u, k) = \frac{\partial}{\partial u} \ln \Theta(u)$$

giving

$$\begin{aligned}\text{(i).} \quad \text{zn}(u, k) &= \frac{\pi}{2K} \frac{\vartheta'_1\left(\frac{\pi u}{2K}\right)}{\vartheta_1\left(\frac{\pi u}{2K}\right)} - \frac{\text{cn } u \, \text{dn } u}{\text{sn } u} \\ \text{(ii).} \quad \text{zn}(u, k) &= \frac{\pi}{2K} \frac{\vartheta'_2\left(\frac{\pi u}{2K}\right)}{\vartheta_2\left(\frac{\pi u}{2K}\right)} - \frac{\text{dn } u \, \text{sn } u}{\text{cn } u} \\ \text{(iii).} \quad \text{zn}(u, k) &= \frac{\pi}{2K} \frac{\vartheta'_3\left(\frac{\pi u}{2K}\right)}{\vartheta_3\left(\frac{\pi u}{2K}\right)} - k^2 \frac{\text{sn } u \, \text{cn } u}{\text{dn } u} \\ \text{(iv).} \quad \text{zn}(u, k) &= \frac{\pi}{2K} \frac{\vartheta'_4\left(\frac{\pi u}{2K}\right)}{\vartheta_4\left(\frac{\pi u}{2K}\right)}\end{aligned}$$

Many different notations for the theta function are in current use. The most common variants are the replacement of the argument u by the argument u/π and, occasionally, a permutation of the identification of the functions ϑ_1 to ϑ_4 with the function ϑ_4 replaced by ϑ .

The Factorial (Gamma) Function

In older reference texts, the gamma function $\Gamma(z)$, defined by the Euler integral

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt,$$

is sometimes expressed in the alternative notation

$$\Gamma(1+z) = z! = \Pi(z).$$

On occasions, the related derivative of the logarithmic factorial function $\Psi(z)$ is used where

$$\frac{d(\ln z!)}{dz} = \frac{(z!)' }{z!} = \Psi(z).$$

This function satisfies the recurrence relation

$$\Psi(z) = \Psi(z-1) + \frac{1}{z-1}$$

and is defined by the series

$$\Psi(z) = -C + \sum_{n=0}^{\infty} \left(\frac{1}{n+1} - \frac{1}{z+n} \right).$$

The derivative $\Psi'(z)$ satisfies the recurrence relation

$$\Psi'(z+1) = \Psi'(z) - \frac{1}{z^2}$$

and is defined by the series

$$\Psi'(z) = \sum_{n=0}^{\infty} \frac{1}{(z+n)^2}.$$

Exponential and Related Integrals

The exponential integrals $E_n(z)$ have been defined by Schloemilch using the integral

$$E_n(z) = \int_1^{\infty} e^{-zt} t^{-n} dt \quad (n = 0, 1, \dots, \operatorname{Re} z > 0).$$

They should not be confused with the Euler polynomials already mentioned. The function $E_1(z)$ is related to the exponential integral $\operatorname{Ei}(z)$ through the expressions

$$E_1(z) = -\operatorname{Ei}(-z) = \int_z^{\infty} e^{-t} t^{-1} dt$$

and

$$\operatorname{li}(z) = \int_0^z \frac{dt}{\ln t} = \operatorname{Ei}(\ln z) \quad \{z > 1\}.$$

The functions $E_n(z)$ satisfy the recurrence relations

$$E_n(z) = \frac{1}{n-1} \{e^{-z} - z E_{n-1}(z)\} \quad [n > 1]$$

and

$$E'_n(z) = -E_{n-1}(z)$$

with

$$E_0(z) = e^{-z}/z.$$

The function $E_n(z)$ has the asymptotic expansion

$$E_n(z) \sim \frac{e^{-z}}{z} \left\{ 1 - \frac{n}{z} + \frac{n(n+1)}{z^2} - \frac{n(n+1)(n+2)}{z^3} + \dots \right\} \quad \left[\arg z < \frac{3\pi}{2} \right]$$

while for large n ,

$$E_n(x) = \frac{e^{-x}}{x+n} \left\{ 1 + \frac{n}{(x+n)^2} + \frac{n(n-2x)}{(x+n)^4} + \frac{n(6x^2 - 8nx + n^2)}{(x+n)^6} + R(n, x) \right\},$$

where

$$-0.36n^{-4} \leq R(n, x) \leq \left(1 + \frac{1}{x+n-1} \right) n^{-4} \quad [x > 0].$$

The sine and cosine integrals $\operatorname{si}(x)$ and $\operatorname{ci}(x)$ are related to the functions $\operatorname{Si}(x)$ and $\operatorname{Ci}(x)$ by the integrals

$$\operatorname{Si}(x) = \int_0^x \frac{\sin t}{t} dt = \operatorname{si}(x) + \frac{\pi}{2}$$

and