

CLASSICS IN MATHEMATICS

Robert M. Switzer

**Algebraic Topology
Homology and
Homotopy**

代数拓扑

Springer-Verlag

世界图书出版公司

Robert M. Switzer

Algebraic Topology – Homotopy and Homology

Reprint of the 1975 Edition

Springer

世界图书出版公司

Robert M. Switzer
Georg-August-Universität
Mathematisches Institut
Bunsenstrasse 3-5
37073 Göttingen
Germany
e-mail: switzer@math.uni-goettingen.de

Originally published as Vol. 212 of the
Grundlehren der mathematischen Wissenschaften

Cataloging-in-Publication Data applied for

Die Deutsche Bibliothek - CIP-Einheitsaufnahme
Switzer, Robert M.:

Algebraic topology : homotopy and homology / Robert M. Switzer. - Reprint of the 1975 ed.. - Berlin;
Heidelberg; New York; Barcelona; Hong Kong; London; Milan; Paris; Tokyo: Springer, 2002
(Classics in mathematics)

ISBN 3-540-42750-3

Mathematics Subject Classification (2000): Primary 55-02;
Secondary 57A55, 57A65, 57D75

ISSN 1431-0821

ISBN 3-540-42750-3 Springer-Verlag Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

Springer-Verlag Berlin Heidelberg New York
a member of BertelsmannSpringer Science+Business Media GmbH

<http://www.springer.de>

© Springer-Verlag Berlin Heidelberg 2002

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the People's Republic of China only and not for export therefrom.

Classics in Mathematics

Robert M. Switzer

**Algebraic Topology—
Homotopy and Homology**

Springer

Berlin

Heidelberg

New York

Barcelona

Hong Kong

London

Milan

Paris

Tokyo

Robert M. Switzer

Algebraic Topology— Homotopy and Homology

Springer-Verlag
Berlin Heidelberg New York 1975

Introduction

This book is the result of lecture courses on algebraic topology given by the author at the University of Manchester in 1967–1970, at Cornell University in 1970–1971 and at the Georg August University, Göttingen, in 1971–1972. The level of the material is more advanced than that of a first-year graduate course in algebraic topology; it is assumed that the student has already had a course on basic algebraic topology which included singular homology, the fundamental group and covering spaces. Moreover, a student who has never encountered differentiable manifolds will probably have difficulty with Chapter 12. On the other hand no knowledge of homotopy theory beyond the fundamental group is assumed.

The last few years have seen the publication of several excellent textbooks on basic algebraic topology, most notably the book by Spanier [80], which I suggest as a companion volume to this one. There is a certain overlap between Spanier's book and this text—particularly in Chapters 0–6, 14 and 15—but the present book goes considerably further and has as its goal that the reader should be brought to a point from which he could begin research in certain areas of algebraic topology: stable homotopy theory, K -theory, cobordism theories.

Despite the title “Algebraic Topology” this book does not (and could not) pretend to achieve the same very advanced level in all areas of this subject. The choice of topics to be emphasized is, of course, heavily influenced by the research interests of the author. Thus, for example, unstable homotopy theory is only developed to the point at which it really begins to be interesting and is then dropped in favor of stable homotopy theory. The reader who finds that his appetite for unstable homotopy theory has been whetted is advised to follow the signposts set up by Adams in [10]. Another important branch of algebraic topology which is omitted is obstruction theory—partly due to lack of space, partly because one could scarcely give a better introduction than Thomas' *Seminar on fibre spaces* [86].

The following basic idea occurs repeatedly as a leitmotiv in this text: the majority of the problems which have been solved by means of algebraic

topology have first been reduced to the question of the existence or non-existence of a continuous function $f: X \rightarrow Y$ between two given topological spaces X, Y . One tries to prove the non-existence, for example, by finding an appropriate functor F from the category of topological spaces to some algebraic category—that is for every space X we are given $F(X)$, a group, ring, module, ..., and for every continuous function $f: X \rightarrow Y$ we have $F(f): F(X) \rightarrow F(Y)$, a homomorphism preserving the given algebraic structure. Then one seeks to demonstrate that the algebraic map $F(f): F(X) \rightarrow F(Y)$ cannot possibly exist. (Proofs of existence are in general more difficult to handle). From this point of view the richer the natural algebraic structure on $F(X)$ the better: if $F(X), F(Y)$ have a very complex algebraic structure, then there will not be many homomorphisms $\phi: F(X) \rightarrow F(Y)$ preserving this structure, and thus the chances of showing $F(f)$ cannot exist are good. At several points, then, (cf. Chapters 2, 13, 17, 18) we shall strive to enrich the natural algebraic structure available on our functors. In Chapter 19 we make the happy discovery that we have a sufficiently complex natural algebraic structure on $F(X)$ that we can (under favorable circumstances) say precisely which algebraic maps $\phi: F(X) \rightarrow F(Y)$ are of the form $\phi = F(f)$ for some continuous function $f: X \rightarrow Y$. At this point existence proofs become possible.

Chapters 0 and 1 contain respectively certain results from set-theoretic topology which are repeatedly used in the text and the basic definitions of category theory; both chapters should be in the nature of a review for the reader. Chapter 2 takes up the sets $[X, Y]$ of homotopy classes of maps $f: X \rightarrow Y$ and deals with such questions as: under what conditions on X or Y is $[X, Y]$ a group, when is a sequence

$$[X, W] \xleftarrow{f^*} [Y, W] \xleftarrow{g^*} [Z, W]$$

exact, etc. (enrichment of structure!). In Chapter 3 we then specialize to $X = S^n$ and consider $\pi_n(Y, y_0) = [S^n, s_0; Y, y_0]$, which is always a group of $n \geq 1$ —the n th homotopy group of Y . The more elementary properties of these groups are demonstrated in this chapter. In Chapter 4 we define the notions of *fibration* and *weak fibration* and show that for a weak fibration $p: E \rightarrow B$ with fibre $F = p^{-1}(b_0)$ there is an exact sequence

$$\cdots \longrightarrow \pi_n(F, e_0) \xrightarrow{i_*} \pi_n(E, e_0) \xrightarrow{p_*} \pi_n(B, b_0) \xrightarrow{\partial} \pi_{n-1}(F, e_0) \longrightarrow \cdots$$

We define the geometrically very important notion of a *fibre bundle* and show that every fibre bundle is a weak fibration. There follow some important examples of fibre bundles:

$$O(n) \rightarrow O(n)/O(n-k), \quad O(n)/O(n-k) \rightarrow O(n)/O(n-k) \times O(k)$$

and others. The chapter is concluded by remarking that a covering $p: X' \rightarrow X$ is a fibre bundle with discrete fibre and by using this remark to compute the homotopy groups of S^1 and $T^n = S^1 \times S^1 \times \cdots \times S^1$.

Proofs of deeper results for arbitrary topological spaces X, Y are difficult; it is not easy to demonstrate, for example, the existence of continuous functions $f: X \rightarrow Y$. We turn therefore to the smaller category of CW-complexes. Since CW-complexes are built up by glueing cells D^n together, it is possible to construct maps and homotopies cell by cell. This property permits strong statements about $[X, Y]$ when X or Y is a CW-complex. In Chapter 5 we define CW-complexes and prove some straightforward properties. Chapter 6 contains some deeper homotopy results, such as: $\pi_n(X, x_0)$ depends only on the cells of dimension at most $n + 1$; the suspension homomorphism

$$\Sigma: \pi_q(X, x_0) \rightarrow \pi_{q+1}(SX, *) \quad ([f] \mapsto [1 \wedge f])$$

is an isomorphism for $q < 2n + 1$ if X is an n -connected CW-complex; a map $f: X \rightarrow Y$ between CW-complexes induces an isomorphism $f_*: \pi_q(X, x_0) \rightarrow \pi_q(Y, y_0)$, $q \geq 0$, if and only if f is a homotopy equivalence.

At this point we turn from homotopy theory to homology and cohomology theories. A *generalized homology theory* is a family of functors $\{h_n: n \in \mathbb{Z}\}$ from the category of topological spaces to the category of abelian groups which satisfies the first six of the seven Eilenberg-Steenrod axioms (i.e. the "Dimension Axiom" is not necessarily satisfied). Chapter 7 is an investigation of the properties of such theories which follow directly from the axioms; this amounts to carrying out such parts of the program of Eilenberg and Steenrod [40] as still go through without the seventh axiom.

Chapter 8 contains a construction of Boardman's stable category of spectra and demonstrations of some of its most important properties—in particular that $\Sigma: [E, F] \rightarrow [E \wedge S^1, F \wedge S^1]$ is a bijection for all spectra E, F . We show how to construct a homology theory E_* and a cohomology theory E^* for every spectrum E . Then in Chapter 9 we prove that we have already constructed all possible cohomology theories in Chapter 8 (Brown's representation theorem). Then come the three most important known examples of homology and cohomology theories: ordinary homology (Ch. 10), K -theory (Ch. 11) and bordism (Ch. 12). In Chapter 10 we also show how to compute the singular homology groups of an arbitrary CW-complex and prove the Hurewicz isomorphism theorem. Chapter 11 contains the computation of the homotopy groups of the stable groups O, U, Sp .

Next comes a chapter on products in homology and cohomology. Chapter 13 begins with the universal coefficient theorems (how does one express $H_*(X; G)$ and $H^*(X; G)$ in terms of $H_*(X; \mathbb{Z})$ and G ?) and the

Künneth theorem (how does one express $H_*(X \times Y)$ in terms of $H_*(X)$, $H_*(Y)$?). Next the \times -, \cup - and \cap -products for singular homology and cohomology are briefly discussed. Then we make a digression in order to construct the smash product of two spectra, whereupon we can describe products in the generalized homology and cohomology theories E_* , E^* associated to a *ring spectrum* E . We describe explicitly the products for ordinary homology, K -theory and bordism.

Chapter 14 then applies what we have learned about products to investigations of duality (Alexander, Lefschetz, Poincaré for manifolds; Spanier-Whitehead for finite spectra) and to questions of orientability of manifolds with respect to generalized cohomology theories. Spanier-Whitehead duality also permits a proof of a representation theorem for homology theories similar to the one for cohomology theories in Chapter 9.

In Chapter 15 the level of difficulty increases with the introduction of spectral sequences. Everyone finds spectral sequences baffling at the first encounter. Experience, however, shows that spectral sequences are among the topologist's most effective tools, so that the effort required to master their use is well worth while. We develop the Atiyah-Hirzebruch-Whitehead and Leray-Serre spectral sequences and make some important applications of them: Gysin, Wang and Serre exact sequences, Leray-Hirsch theorem, Thom isomorphism theorem.

Chapter 16 is concerned with the calculation of the homology and cohomology rings $E_*(BG)$, $E^*(BG)$ of the stable classifying spaces BG , $G = O$, U and Sp . Here the Atiyah-Hirzebruch-Whitehead spectral sequence proves very useful. In the process we construct certain classes $c_i \in E^*(BG)$ with whose help we can form invariants $c_i(\xi) \in E^*(X)$ of isomorphism classes of G -vector bundles $\xi \rightarrow X$ —the so-called *characteristic* classes. The chapter ends with a proof of the Bott periodicity theorem $BU \simeq \Omega_0^2 BU$.

Chapter 17 represents a large step in our natural-algebraic-structure-enrichment program: we show how to make $E^*(X)$ into a module over an algebra $A^*(E) = E^*(E)$ and $E_*(X)$ into a comodule over a Hopf algebra $A_*(E) = E_*(E)$ in a natural way (under favorable conditions on E). We compute the Hopf algebras $A_*(E)$ for $E = MU$, MSp , K and KO and find they have satisfyingly complex algebraic structures (recall our leitmotiv). As an application we find bounds on the image of the Hurewicz homomorphisms

$$\begin{aligned} h_K : \pi_q(MU) &\rightarrow K_q(MU) \\ h_{KO} : \pi_q(MSp) &\rightarrow KO_q(MSp) \end{aligned}$$

for small values of q .

Chapter 18 contains a determination of the algebras $A^*(H(\mathbb{Z}_2))$ and $A_*(H(\mathbb{Z}_2))$ including the construction of certain elements $Sq^i \in A^*(H(\mathbb{Z}_2))$. The analogous results for $A_*(H(\mathbb{Z}_p))$, $A^*(H(\mathbb{Z}_p))$ are stated without proof.

In Chapter 19 we enjoy the triumph of our natural-algebraic-structure-enrichment program: using the comodule structure of $E_*(X)$ over $E_*(E)$ constructed in Chapter 17 we can build a spectral sequence (the *Adams spectral sequence*) $\{E_r^s, d_r\}$ whose E_2 -term is the purely algebraic construct

$$\text{Ext}_{E_*(E)}^{s,t}(\tilde{E}_*(S^0), E_*(X))$$

($\text{Ext}_C^{s,t}(-, -)$ is a functor derived from $\text{Hom}_C^s(-, -)$, C a coalgebra, and thus has to do with algebraic maps) and which converges (for connected E) to $\pi_*^S(X)/D_*$, the stable homotopy of X modulo a certain subgroup D_* which depends on E . For $E = MU$ or MSp we show that $D_* = 0$; for $E = H(\mathbb{Z}_p)$ D_* is the subgroup of elements of finite order prime to p . Thus in cases where the spectral sequence proves manageable one can start from a knowledge of $E_*(X)$ as $E_*(E)$ -comodule and compute $\pi_*^S(X)/D_*$. We show how the Adams spectral sequence has permitted the determination of $\pi_q^S(S^0)$ for small q . We then turn to a consideration of the spectral sequence for $E = K, KO$. Here E is not connected, so the spectral sequence may not converge, but it still provides homomorphisms

$$\begin{aligned} e_C: \ker k_K &\rightarrow \text{Ext}_{K_*(K)}^{1, q+1}(\tilde{K}_*(S^0), K_*(X)) \\ e_R: \ker h_{KO} &\rightarrow \text{Ext}_{KO_*(KO)}^{1, q+1}(\tilde{KO}_*(S^0), KO_*(X)). \end{aligned}$$

The Ext -groups for $X = S^0$ are computed and the result is used to localize a non-trivial direct summand in $\pi_*^S(S^0)$ whose order is related to the Bernoulli numbers.

Chapter 20 then represents an extended application of the Adams spectral sequence. The cases $E = H(\mathbb{Z}_p)$ and $X = MG$, $G = O, SO$, or U are, fortunately, of the sort in which the Adams spectral sequence is manageable and permits a complete determination of $\pi_*(X) = \pi_*(MG) \cong \Omega_*^G$, the G -cobordism ring. We also prove the theorem of Hattori and Stong, which describes the image of

$$h_K: \pi_*(MU) \rightarrow K_*(MU).$$

After reading Chapter 20 the student should be able to understand without undue difficulty the papers [13, 14] of Anderson, Brown and Peterson in which Ω_*^{SU} and Ω_*^{Spin} are determined.

This summary undoubtedly makes clear that the level of mathematical expertise demanded of the reader rises rather markedly from Chapter 2, say, to Chapter 20. The student who begins with the minimal prerequisites described in the first paragraph will not acquire the facility needed for understanding the later chapters merely by reading straight through. He must try to master the material in each chapter to such an extent that he can apply it to the solution of problems other than those worked out in the text. In some cases he will find it valuable to seek further applications in the books and articles listed at the end of each chapter.

The bibliography included here does not attempt to be comprehensive. Steenrod's valuable compendium of all mathematical reviews having to do with topology makes such a comprehensive bibliography unnecessary. Instead this bibliography has two goals: (1) to suggest to the student where he might begin to pursue a given topic further and (2) to acknowledge the sources from which much of the material in this text is drawn.

In addition, however, I wish to acknowledge quite explicitly and with gratitude my debt to Frank Adams. It is no exaggeration to say that most of what I know about algebraic topology I learned from him. Anyone familiar with his work will recognize the influence of his way of looking at algebraic topology on the presentation of the subject given here. Moreover, on certain topics (e.g. Chapter 8, Chapter 9 in the case of finite CW-complexes, the construction of $E \wedge F$) I have largely reproduced his presentation of the topic with only small alterations (which may not have been for the better).

I further wish to express my deep gratitude to Egbert Brieskorn for his encouragement and helpful suggestions. My thanks also go to Fräulein Ingrid Sochaczewsky and Frau Christiane Preywisch for help in typing the manuscript.

The manuscript of this book was submitted to the Georg August University, Göttingen, in January 1973 as Habilitationsschrift.

Göttingen, November, 1973

ROBERT M. SWITZER

Table of Contents

Chapter 0	Some Facts from General Topology	1
Chapter 1	Categories, Functors and Natural Transformations	6
Chapter 2	Homotopy Sets and Groups	11
Chapter 3	Properties of the Homotopy Groups	36
Chapter 4	Fibrations	52
Chapter 5	CW-Complexes	64
Chapter 6	Homotopy Properties of CW-Complexes	74
Chapter 7	Homology and Cohomology Theories	99
Chapter 8	Spectra	133
Chapter 9	Representation Theorems	152
Chapter 10	Ordinary Homology Theory	167
Chapter 11	Vector Bundles and K -Theory	190
Chapter 12	Manifolds and Bordism	218
Chapter 13	Products	233
Chapter 14	Orientation and Duality	306
Chapter 15	Spectral Sequences	336
Chapter 16	Characteristic Classes	375
Chapter 17	Cohomology Operations and Homology Cooperations	411
Chapter 18	The Steenrod Algebra and its Dual	440
Chapter 19	The Adams Spectral Sequence and the e -Invariant	458
Chapter 20	Calculation of the Cobordism Groups	490
	Bibliography.	518
	Subject Index	522

Chapter 0

Some Facts from General Topology

It is assumed that the reader is familiar with the elements of general topology—e.g. the most important properties of continuous functions, compact sets, connected sets, etc. Nevertheless, certain general topological results which will be used repeatedly in this book are assembled here for the reader's convenience.

0.1. Let X be a topological space and A_1, A_2, \dots, A_n be closed subspaces such that $X = \bigcup_{i=1}^n A_i$.

Suppose $f_i: A_i \rightarrow Y$ is a function, $1 \leq i \leq n$; there is a function $f: X \rightarrow Y$ such that $f|_{A_i} = f_i$, $1 \leq i \leq n$, if and only if $f_i|_{A_i \cap A_j} = f_j|_{A_i \cap A_j}$, $1 \leq i \leq n$, $1 \leq j \leq n$. In this case f is continuous if and only if each f_i is.

We shall often set about defining a continuous function $f: X \rightarrow Y$ by cutting up X into closed subsets A_i and defining f on each A_i separately in such a way that $f|_{A_i}$ is obviously continuous; we then have only to check that the different definitions agree on the "overlaps" $A_i \cap A_j$.

0.2. *The universal property of the cartesian product:* let $p_X: X \times Y \rightarrow X$, $p_Y: X \times Y \rightarrow Y$ be the projections onto the first and second factors respectively. Given any pair of functions $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ there is a unique function $h: Z \rightarrow X \times Y$ such that $p_X \circ h = f$, $p_Y \circ h = g$. h is continuous if and only if both f and g are. This property characterizes $X \times Y$ up to homeomorphism. The unique h will often be denoted by (f, g) .

In particular, to check that a given function $h: Z \rightarrow X \times Y$ is continuous it will suffice to check that $p_X \circ h$ and $p_Y \circ h$ are continuous.

0.3. *The universal property of the quotient:* let α be an equivalence relation on a topological space X , let X/α denote the space of equivalence classes and $p_\alpha: X \rightarrow X/\alpha$ the natural projection. Given a function $f: X \rightarrow Y$, there is a function $f': X/\alpha \rightarrow Y$ with $f' \circ p_\alpha = f$ if and only if $x \sim x'$ implies $f(x) = f(x')$. In this case f' is continuous if and only if f is. This property characterizes X/α up to homeomorphism.

An important example of a quotient which we frequently encounter is that of a space X with a closed subspace A collapsed to a point. Explicitly,

if $A \subset X$ is a closed non-empty subspace, we take the relation

$$\alpha = A \times A \cup \{(x, x) : x \in X\} \subset X \times X$$

and let $X/A = X/\alpha$. For $A = \emptyset$ we adopt the convention that $X/A = X/\emptyset = X^+ = X \cup \{a \text{ disjoint point}\}$. This convention will be seen to be justified by the fact that every theorem we state about X/A will hold equally well for X/\emptyset .

0.4. Product and quotient combined: if α is an equivalence relation on a topological space X and β is an equivalence relation on Y , then there is an obvious equivalence relation $\alpha \times \beta$ on $X \times Y$:

$$(x, y) \alpha \times \beta (x', y') \Leftrightarrow x \alpha x' \quad \text{and} \quad y \beta y'.$$

There is a unique continuous function

$$\phi: \frac{X \times Y}{\alpha \times \beta} \rightarrow (X/\alpha) \times (Y/\beta),$$

such that $\phi \circ p_{\alpha \times \beta} = p_\alpha \times p_\beta$ and which is even bijective—but not necessarily a homeomorphism. One important case in which ϕ is a homeomorphism is that where β is the identity relation 1 and Y is locally compact.

Proof: We must prove that if $T \subset \frac{X \times Y}{\alpha \times 1}$ is open then $\phi(T)$ is open in $(X/\alpha) \times Y$. T open means $T' = p_{\alpha \times 1}^{-1}(T)$ is open in $X \times Y$. Suppose $(x_0, y_0) \in \phi(T)$ and choose $x_0 \in X$ with $p_\alpha(x'_0) = x_0$. By the definitions of ϕ and T' , we have $(x'_0, y_0) \in T'$; and since T' is open, there exist neighborhoods U of x'_0 and K of y_0 in X, Y respectively such that $U \times K \subset T'$. We may assume K is compact.

Let $J' = \{x' \in X : x' \times K \subset T'\}$. We first show that J' is open. For every $x' \in J'$ we can, because K is compact, find an open neighborhood $U_{x'}$ of x' such that $U_{x'} \times K \subset T'$, which proves J' is open. If $J = \{x \in X/\alpha : x \times K \subset \phi(T)\}$, then clearly $p_\alpha^{-1}(J) = J'$, so J is open in X/α . Also $(x_0, y_0) \in J \times K \subset \phi(T)$, so $\phi(T)$ is open. \square

Example. The unit interval $I = [0, 1]$ is compact, so $\frac{X \times I}{\alpha \times 1} \cong (X/\alpha) \times I$.

0.5. A homotopy from X to Y is a continuous function $F: X \times I \rightarrow Y$. For each $t \in I$ one has $F_t: X \rightarrow Y$ defined by $F_t(x) = F(x, t)$ for all $x \in X$. The functions F_t are called the “stages” of the homotopy. If $f, g: X \rightarrow Y$ are two maps, we say f is *homotopic* to g (and write $f \simeq g$) if there is a homotopy $F: X \times I \rightarrow Y$ such that $F_0 = f$ and $F_1 = g$. In other words, f can be continuously deformed into g through the stages F_t . If $A \subset X$ is a subspace, then F is a *homotopy relative to A* if $F(a, t) = F(a, 0)$, all $a \in A, t \in I$.

0.6. The relation \simeq is an equivalence relation.

Proof: $f \simeq f$ is obvious; take $F(x, t) = f(x)$, all $x \in X, t \in I$. If $f \simeq g$ and F is a homotopy from f to g , then $G: X \times I \rightarrow Y$ defined by $G(x, t) = F(x, 1 - t)$, $x \in X, t \in I$, is a homotopy from g to f —i.e. $g \simeq f$. If $f \simeq g$ with homotopy F and $g \simeq h$ with homotopy G , then $f \simeq h$ with homotopy H defined by

$$H(x, t) = \begin{cases} F(x, 2t) & 0 \leq t \leq 1/2 \\ G(x, 2t - 1) & 1/2 \leq t \leq 1. \end{cases}$$

Note that here we use 0.1 to show H is continuous. \square

0.7. Thus the set of all continuous functions $f: X \rightarrow Y$ is partitioned into equivalence classes under the relation \simeq . The equivalence classes are called *homotopy classes*, and the set of all homotopy classes is denoted by $[X; Y]$. If $f: X \rightarrow Y$, then the homotopy class of f is denoted by $[f]$.

As an application of 0.4 we get the following proposition.

0.8. Proposition. If α is an equivalence relation on a topological space X and $F: X \times I \rightarrow Y$ is a homotopy such that each stage F_t factors through X/α —i.e. $x\alpha x' \Rightarrow F_t(x) = F_t(x')$ —then F induces a homotopy $F': (X/\alpha) \times I \rightarrow Y$ such that $F' \circ (p_\alpha \times 1) = F$.

Proof: The hypothesis on F is precisely that F factors through $\frac{X \times I}{\alpha \times 1}$ —i.e.

there exists a continuous function $F'': \frac{X \times I}{\alpha \times 1} \rightarrow Y$ such that $F'' \circ p_{\alpha \times 1} = F$.

By 0.4 $\phi: \frac{X \times I}{\alpha \times 1} \rightarrow (X/\alpha) \times I$ is a homeomorphism. Then $F' = F'' \circ \phi^{-1}$:

$(X/\alpha) \times I \rightarrow Y$ is the required homotopy. \square

Example. If A is a closed subspace of X and $F: X \times I \rightarrow Y$ is a homotopy such that $F(a, t) = F(a', t)$ for all $a, a' \in A, t \in I$, then F induces a homotopy $F': (X/A) \times I \rightarrow Y$.

0.9. Function spaces: if X and Y are topological spaces, we let Y^X denote the set of all continuous functions $f: X \rightarrow Y$. We give this set a topology, called the *compact-open topology*, by taking as a subbase for the topology all sets of the form $N_{K, U} = \{f: f(K) \subset U\}$, $K \subset X$ compact, $U \subset Y$ open.

0.10. The *evaluation function* $e: Y^X \times X \rightarrow Y$ defined by $e(f, x) = f(x)$ is continuous if X is locally compact. In all the cases we shall consider X will be $I = [0, 1]$.

0.11. The exponential law: if X, Z are Hausdorff spaces and Z is locally compact, then the natural function

$$E: Y^{Z \times X} \rightarrow (Y^Z)^X$$

defined by $(Ef(x))(z) = f(z, x)$ is a homeomorphism.

0.12. Base points: in what follows we shall often have to consider not just a topological space X but rather a space X together with a distinguished point $x_0 \in X$ called the *base point*. The pair (X, x_0) is called a *pointed space* (one also speaks of pointed sets). When we are concerned with pointed spaces (X, x_0) , (Y, y_0) , etc. we always require that all functions $f: X \rightarrow Y$ shall preserve base points—i.e. $f(x_0) = y_0$ —and that all homotopies $F: X \times I \rightarrow Y$ be relative to the base point—i.e. $F(x_0, t) = y_0$, all $t \in I$ —unless an explicit disclaimer to the contrary is made. We shall use the notation $[X, x_0; Y, y_0]$ to denote the homotopy classes of base point-preserving functions—where homotopies are rel x_0 , of course. $[X, x_0; Y, y_0]$ is a pointed set with base point f_0 , the constant function: $f_0(x) = y_0$, all $x \in X$.

Let us use the notation $(Y, B)^{(X, A)}$ to denote the subspace of Y^X consisting of those functions such that $f(A) \subset B$, where $A \subset X$, $B \subset Y$ are subspaces. There are obvious generalizations of this notation, such as $(Y, B, B')^{(X, A, A')}$, etc. In particular, if (X, x_0) , (Y, y_0) are pointed spaces, then we have the space $(Y, y_0)^{(X, x_0)}$ of base point-preserving functions. It has as base point f_0 , where $f_0(x) = y_0$, all x .

If (X, x_0) , (Y, y_0) and (Z, z_0) are three pointed spaces, we can form

$$((Y, y_0)^{(Z, z_0)}, f_0)^{(X, x_0)}, \quad f_0(z) = y_0, \quad \text{all } z \in Z,$$

which is a subspace of $(Y^Z)^X$. Thus we may ask what subspace of $Y^{Z \times X}$ corresponds to it under the exponential function of 0.11. One readily sees that the answer is

$$(Y, y_0)^{(Z \times X, Z \times \{x_0\} \cup \{z_0\} \times X)}.$$

We use the notation $Z \vee X$ for the subspace $Z \times \{x_0\} \cup \{z_0\} \times X$ of $Z \times X$. It can be thought of as the result of taking the disjoint union $Z \cup X$ and identifying z_0 with x_0 . $Z \vee X$ is again a pointed space with base point (z_0, x_0) . Then we have

0.13. Proposition. *The exponential function $E: Y^{Z \times X} \rightarrow (Y^Z)^X$ induces a continuous function*

$$E: (Y, y_0)^{(Z \times X, Z \vee X)} \rightarrow ((Y, y_0)^{(Z, z_0)}, f_0)^{(X, x_0)}$$

which is a homeomorphism if Z and X are Hausdorff and Z is locally compact.

Remark. The subspace $Z \times \{x_0\} \cup \{z_0\} \times X \subset Z \times X$ is called the *wedge sum* of Z and X and is characterized by the property that for any continuous functions $f: (Z, z_0) \rightarrow (W, w_0)$, $g: (X, x_0) \rightarrow (W, w_0)$ there is a unique continuous function $h: (Z \vee X, *) \rightarrow (W, w_0)$ such that $h|_Z = f$, $h|_X = g$. (Compare this with 0.2.) The unique function h will be denoted by