

third edition

---

# Diseases of the Ear

---

Stuart R. Mawson

# Diseases of the Ear

Third edition

**Stuart R. Mawson**

M.B.(Camb.), F.R.C.S.(Eng.), D.L.O.(Eng.)

Senior Consultant Surgeon, Department of Otolaryngology,  
King's College Hospital (including Belgrave Hospital for Children), London



Edward Arr

© Stuart R. Mawson 1974

First published 1963  
by Edward Arnold (Publishers) Ltd  
25 Hill Street, London W1X 8LL

Second edition 1967

Third edition 1974

ISBN: 0 7131 4221 9

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photo-copying, recording or otherwise, without the prior permission of Edward Arnold (Publishers) Ltd.

Printed in Great Britain by  
Alden & Mowbray Ltd  
at the Alden Press, Oxford

# Preface to the Third Edition

In a preface to the third edition of Politzer's *Textbook of the Diseases of the Ear*, published in 1894, Sir William Dalby wrote: 'as a treatise on diseases of the ear it is complete since it embraces all that is known of the subject'. How fortunate, in time spared from clinical practice, to be able to garner the totality of a subject, write it all down and, having done so, find oneself with a modest sized volume! Such has been the proliferation of knowledge since then, especially in the last few decades, that the problems of keeping fully informed, making appropriate selection and compressing within a reasonable length have assumed truly daunting proportions for any would-be author.

When the first edition of *Diseases of the Ear* appeared in 1963 surgical techniques were outstripping, certainly in interest, the basic sciences of otology. Now the reverse is true. Audiology, microhistology, physiology and chemical pathology are everywhere advancing and converging with new technical vocabularies to match. An attempt has been made to incorporate these growing points within the pre-existing textual framework without disturbing the balance of the whole, which remains poised on the assumption that there is still a place for an orthodox methodology from which to prepare for higher examinations in otology, and to obtain guidance for the safe and practical management of all aspects of diseases of the ear.

I would like, once more, to express my grateful thanks to my colleague and fellow otolaryngologist Harold Ludman, F.R.C.S., who has renewed some of the illustrations and introduced several others, and to Dr Stanley Mason, F.F.A.R.C.S., now senior anaesthetist at King's College Hospital, who has again revised the chapter on anaesthesia. I would above all like to dedicate this third edition to my dear wife who has ever sustained and supported me in the tasks thus far accomplished.

STUART MAWSON

# Preface to the First Edition 19

The last two decades have seen a remarkable revival of interest in the ear. Advances in treatment, of deafness in particular, have been as spectacular as those in any branch of surgery or medicine. Originally a separate speciality, otology has so enlarged in scope that it may be said again to occupy a wholly engrossing place within the general field of otorhinolaryngology.

At the same time, for those entering the speciality, and indeed for those already within it, methodical study and methodical teaching have been handicapped by the lack of a single, comprehensive text-book of recent origin. It was felt that a real need existed to assemble and present the subject of otology anew under uniform authorship. This is the task I have undertaken and attempted. Although written with the needs of the specializing postgraduate student primarily in view it is nevertheless also hoped that *Diseases of the Ear* will prove a useful book of reference for general practitioners, house surgeons, registrars, consultants, all, in fact, who may be called upon to treat patients with aural conditions.

Full understanding of the changes resulting from disease requires a very comprehensive knowledge of normality. A relatively large section has, therefore, been set aside for anatomy, physiology and other introductory subjects, corresponding with the requirements of the 1st parts of the examinations for the Fellowship (E.N.T.) of the Royal Colleges of Surgeons, and for the Diploma in Laryngology and Otology of London. The anatomical descriptions have been written with temporal bone dissections at hand, and reciprocal study of such specimens will facilitate assimilation of the text. Operative surgery has been relegated to a separate section at the end, where operating theatre details may be studied without distracting from the primary task of seeing a disease as a whole and of comprehending the principles of treatment upon which detail is based. Elsewhere the classical division of subjects into diseases of the external, middle and internal ear has been followed.

Every single one of the illustrations is original, and has been specially drawn in pen and ink by Harold Ludman, F.R.C.S. I have been incomparably fortunate in having as illustrator one already specializing in otolaryngology, and I am happy to express my deep gratitude to him for his great help and artistry.

I would also place on record my gratitude for the assistance and encouragement I have received at all times from my colleagues in the Ear, Nose and Throat Department at King's College Hospital, W. I. Daggett, F.R.C.S., Terence Cawthorne, F.R.C.S. and R. S. Lewis, F.R.C.S. For having read my manuscripts with the most critical care, kept alive my interest and answered many questions, despite heavy demands on their time, I am and always shall be greatly in their debt.

To Dr Stanley A. Mason, F.F.A.R.C.S., Anaesthetist to King's College Hospital, my grateful thanks are due, not only for the section on anaesthesia appearing in Chapter XIX, but for the many harmonious years of co-operation we have enjoyed together in the operating theatre. My preference for general anaesthesia for nearly all operations upon the ear derives from the excellence

of the methods advocated. Patients and surgeons who have had experience of a really good, safe general anaesthetic rarely remain in doubt of its advantages.

I would also like to acknowledge my indebtedness to Dr J. Blewett for the radiographs, and to Dr. D. I. Williams for some of the photographs of skin diseases appearing in Part III. Other photographs have been specially taken by Mr W. Smith, of the photographic department, whose skill and help have been invaluable.

To Miss Lorna Rutty, my medical secretary, I owe a very heavy debt of gratitude for her spare-time typing of the manuscript. But for her patience and forbearance the book might never have appeared. To her, to Miss Sargeant and the library staff at King's College Hospital, and to all others who have helped me in numerous ways, above all to my wife, I am most sincerely grateful.

STUART MAWSON

*Jucundi acti labores*

# Contents

## Part 1 Introductory

1	Prenatal and postnatal development of the ear	I
2	Anatomy	17
3	Physiology of hearing and balance	63
4	Methods of examination	79

## Part 2—Disturbance of Function

5	Auditory function	123
6	Vestibular function	176
7	Facial nerve palsy	183

## Part 3—Diseases of the External Ear

8	Congenital malformations	201
9	Diseases due to trauma or physical agents	218
10	Infections	233
11	Cysts and tumours	263

## Part 4—Diseases of the Middle Ear

12	Otitis media	283
13	Complications of suppurative otitis media	349
14	The Eustachian tube	414
15	Tumours of the middle ear and mastoid	422
16	Otosclerosis	435

## Part 5—Diseases of the Internal Ear

17	Cochleovestibular diseases	455
18	Neurocentral diseases	495

## Part 6—Operative Surgery

19	General principles	507
20	Operations on the external ear	514
21	Operations on the middle ear	524
22	Operations on the internal ear	565

## Index

**ADVANCED ORGANIC CHEMISTRY:  
REACTIONS, MECHANISMS, AND STRUCTURE**






Fig. 2. Dorsal view of germ disc to show primitive streak and formation of secondary mesoderm.

proliferates between ectoderm and endoderm as primordial secondary mesoderm (Fig. 2) From these three primitive layers the aural derivatives arise as follows:

Ectoderm	Mesoderm	Endoderm
External ear Auricle, external auditory meatus, skin, glands	Middle ear* Upper half, ossicles and mastoid	Middle ear Lower half
Tympanic membrane Outer layer	Tympanic membrane Middle layer	Tympanic membrane Inner layer
Membranous labyrinth Organ of Corti, utricle, saccule, semicircular canals	Bony labyrinth Petrous temporal bone	Pharyngo-tympanic (Eustachian) tube

\* Some authorities consider the mucosa of all air spaces is derived from endoderm.

Since the ear develops in close relation to the dorsal ends of the first and second branchial arches, these must next be considered (Fig. 3). Development begins at 3 weeks with the expansion of the cephalic part of the foregut, an entodermal derivative lying dorsal to the pericardium, to form the primitive pharynx, and is continued by the formation of 13 depressions in the walls of the pharynx, 5 bilaterally represented pouches, the rudimentary thyroid, the respiratory tract and the impermanent pouch of Siessel. The entodermal lining of the pouches is separated from surface ectoderm by the secondary mesoderm. On the external aspect of the embryo, opposite the pouches, 4 grooves or clefts appear in the ectoderm, dividing the intervening side wall of the pharynx into bars or arches. These bars are numbered from the cephalic end, and, as growth proceeds, bars 1 and 2 meet across the midline between the pericardium and the head. Bar 1 persists as the mandible, bar 2 as the hyoid, but bars 3, 4 and 5 remain small and unjoined and soon become overlain and recessed behind a caudal




Fig. 3. Front end of embryo to show pharyngeal pouches.

expansion of the second, hyoid, bar. This recess, known as the cervical sinus, is the nursery of branchial cysts and fistulae (Fig. 4).

Thus the first external groove or cleft is the only one not submerged beneath or overlain by the second bar or arch, and is the primordium of the external auditory meatus. At the same time, the internal aspect of the second bar forms part of the wall of a lateral extension of the upper part of the cavity of the primitive pharynx known as the tubotympanic recess. This recess also contains




Fig. 4. Development of the cervical sinus; (b) shows overgrowth of second arch caudally.

first and third pharyngeal pouches. The tympanic membrane, which separates the tympanic cavity from the external auditory meatus, is formed in the position of a separating membrane between the first pouch in the tubotympanic recess and the first external groove. But the membrane is a secondary formation since the deeper part of the meatus is first formed by a solid plug of ectodermal cells which later break down in the centre to form the meatus (Fig. 5).




Fig. 5. Development of tympanic membrane.

The third bar or arch, meanwhile, although overlain by the second, continues to grow forward under it, and succeeds in narrowing the inner part of the tubotympanic recess to form the Eustachian tube region, while the outer part of the recess develops into the tympanic cavity. The subsequent development of the ear may now be considered in detail.

### Internal ear

At 3 weeks a thickening of ectoderm, the auditory (otic) placode, appears opposite the hind brain immediately above the first external pharyngeal groove. By invagination the placode first becomes a pit, then a closed pit. Sinking beneath the surface ectoderm the auditory vesicle (otocyst), as it is now called, comes to rest in mesoderm. With the intervention of the tubotympanic recess between it and the surface ectoderm, the ultimate gross relations of the three parts of the ear are thus established (Fig. 6).




Fig. 6. Development of the otocyst.




Fig. 7. Development of membranous labyrinth (a) 5 weeks, (b) 6 weeks, (c) 8-9 weeks.

The vesicle elongates and develops an indenting groove on its medial side which demarcates a tubular diverticulum, soon further modified into the endolymphatic sac and duct. The major compartment of the vesicle, termed the utriculosaccular chamber, differentiates into semicircular canals and cochlea (Fig. 7). From the dorsal part of the chamber three disc-like evaginations appear in the right-angled disposition of the canals, acquire a corpuscular shape and, with further absorption of their centres, by 9 weeks become free semicircles. From the ventral part of the chamber a single evagination pushes medially as the cochlear duct and by 12 weeks it too, fully coiled, achieves definition. The remaining central portion of the utriculosaccular chamber represents the membranous vestibule, with the utricle developing dorsally and the saccule




Fig. 8. Diagrammatic representation of the development of the scala tympani and scala vestibuli.

ventrally. At 8 weeks the mesoderm investing the differentiating auditory vesicle shows signs of conversion into a cartilaginous capsule, later to become ossified. In this cartilage the scala tympani and scala vestibuli gradually extend along each side of the cochlear duct until, when the tip is reached, they approximate and fuse with the formation of an opening, the helicotrema (Fig. 8). Ossification of the cartilaginous capsule begins and is generally completed between the 16th and 23rd week.

### Fissula ante fenestram

Between the cartilaginous otic capsule and the membranous labyrinth, the mesoderm differentiates into a mesh-like connective tissue constituting the

periotic space or periotic labyrinth. Immediately in front of the oval window, where the vestibule unites with the scala vestibuli, a streak appears in the cartilage of the otic capsule which bears a close resemblance to the young mesh-like periotic tissue. This streak seems to develop in the direction of the middle ear as an extension of the tissue lining the vestibule, in other words as an evagination of periotic tissue into the capsule (Bast & Anson 1949). At the same time, when the streak has reached the middle ear it is in turn invaded by a vascular bud of middle ear epithelium which hastens the absorption of cartilage and growth of the streak. From the 16th to 20th week, the period during which cochlea and vestibule are attaining their full dimensions, the fissula is extending until at the end of this period it too has reached maximum size. In the adult the fissula appears as a fibrous slot in the bone. Its importance lies in that, in immediate proximity to the stapedial footplate, it forms the site of predilection for the formation of abnormal, otosclerotic bone.

### **Middle ear**

In the development of the middle ear there are three components to be considered: the air spaces, the lining membrane, and the ossicles. The lining membrane and air spaces of the Eustachian tube and lower tympanum, up to the level of the chorda tympani, are converted directly from the tubotympanic recess of entoderm, involving the first and second pharyngeal pouches. The ossicles are formed by ossification in cartilage out of mesoderm. Majority opinion holds that the malleus and incus are derived from the dorsal end of the mandibular (Meckel's) cartilage, and the stapes, with the possible exception of the footplate, from the dorsal end of the cartilage of the second arch.

### **Air spaces and lining membrane**

At 12 weeks the differentiating ossicles are at first embedded in the mesoderm overlying the tubotympanic recess (Fig. 9). This investing mesoderm loosens and decreases in cellularity until at about 20 weeks the ossicles are lying in a vacuolated connective tissue matrix. By this time the tympanic ring has already ossified, the tympanic membrane and outer attic wall (tympanic process of squamous portion of temporal bone) have formed, and a process of cartilage has grown out laterally from the otic capsule giving rise to the tegmen tympani and, more caudally, to the lateral wall of the Eustachian tube.

At 22 weeks the dorsal end of the tubotympanic recess or primitive tympanic cavity, hitherto slit-like and flattened, undergoes expansion. The lining epithelial membrane is believed to enter and invade the vacuolated mesoderm and to wrap itself round the ossicles, carrying in potential space of the tubotympanic recess in the same way as the peritoneal mesothelium covers the intestine (Bast & Anson 1949). The subsequent existence of mucosal folds is attributed to a four-pronged invasion of the tubotympanic process by pseudopodial buds or sacs, the folds being formed by the party walls between the buds. This vital process of tympanic expansion by active encroachment of tubotympanic (entodermal) cells into pre-loosened and vacuolated mesoderm is accepted as the basis for hypotheses attempting to explain lack of mastoid pneumatization on the grounds of either hereditary failure in this active process, or of interference with it by infective or other pathological changes. At 29 weeks the otic capsule is well ossified, the

middle ear is mostly pneumatized\* and the epithelium is extending towards the region of the antrum which is still occupied by the epitympanic connective tissue. The epithelium of the middle ear is then described as reaching the mouth of the antrum and by 34 weeks as having half pneumatized it. Thereafter the pneumatized antrum extends into the capsular bone to form antrum air cells and laterally into the growing tympanic process of the squamous bone to form air cells in the region of the primitive mastoid process. The concept is firmly one of continuous active expansion and invasion by epithelium cells originating from the primitive tympanum and continuing, in the case of the mastoid air cells, through infancy and childhood. This uncompromising support for the classical concept of pneumatization, as advanced by Frazer, Wittmaack and others, indicates the




Fig. 9. Development of middle ear.

position of present day orthodoxy. It cannot be denied, however, that the theory of entodermal invasion leaves important questions unanswered, chiefly the histological difference between the epithelium lining the Eustachian tube and lower part of the tympanic cavity, and the epithelium lining the epitympanum, antrum and mastoid air cells; a difference that is strikingly emphasized by markedly dissimilar response to infection by the two epithelial areas. Other observers (Schwarzbart 1959) have suggested that the process of pneumatization is accomplished primarily in mesoderm, that the vacuolated spaces appearing in the temporal bone simply represent bone marrow spaces similar to spaces in other bones, that these spaces acquire a lining of cells differentiated out of marrow cells,

\* Pneumatization here means formation of epithelial lined potential space.

that the tubotympanic recess takes up a position adjacent to the marrow spaces and that the party wall then ruptures, allowing air into the spaces immediately infantile respiration is established.

### The ossicles

As already indicated, the generally accepted view is that the malleus and incus are derived from the dorsal end of the mandibular (1st arch) cartilage, and the stapes from the cartilage of the dorsal end of the 2nd arch. Dissenting views vary between those which ascribe a non-branchial origin to the stapes and that which considers the second cartilage primordial to stapes, incus and most of malleus. On the available evidence it would appear that the majority view is correct, with the reservation that part of the footplate of the stapes may be derived from the otic capsule. At 12 weeks the ossicles are differentiating and at 16 weeks are fully formed in cartilage and beginning to ossify.




Fig. 10. Ossification of malleus and incus.

A centre of ossification first appears on the anterior surface of the long process of the incus and spreads in an encircling movement, the short process being the last to be invested with bone (Fig. 10). The malleus next exhibits a small plaque of bone on the medial surface of the body, near the area of articulation with the incus, from which centre of ossification bone spreads upwards over the body and downwards over the handle.

Last to ossify is the stapes. From a single centre appearing in the footplate, bone spreads along each crus towards the head. Ossification of the stapes is accompanied by complicated remodelling and, having regard to the importance of this bone in the surgery of otosclerosis, more detailed consideration must be given to its development than to that of the malleus and incus.

### Stapes (Fig. 11)

The stapes begins as a ring of cartilage through which passes the stapedia artery. At 16 weeks the ring, by flattening at one end and extrusion of a head



at the other, has become a recognizable cartilaginous precursor of the adult form. The changes next to be undergone are to result in the conversion of this solid cartilaginous miniature into the frail and excavated bone so readily fractured under manipulation.

The single ossification centre appears as a plaque on the tympanic surface of the footplate and spreads upwards upon the inner aspect of the crura to extend as an encirclement of periosteal bone over all the surfaces, except for the articular aspect of the head and the labyrinthine aspect of the footplate which remain




Fig. 11. Development of the stapes.

cartilaginous. At the same time as the bone is spreading superficially there is an invasion deeply, with dissolution of underlying cartilage, by buds of periosteal osteogenic tissue. As the cartilage disappears the crura become converted into tubes of periosteal bone, the core being occupied by marrow spaces. Then, unexpectedly, the early shell of periosteal bone on the inner aspect of each crus begins to undergo extensive erosion and to become widely cavitated, the marrow space thus exposed being replaced by an invasion of entodermal mucosa (see formation of air spaces and lining membrane, page 7). The crura, therefore, have neither the strength nor stability of a solid cylinder, nor even of a hollow cylinder, but eventually only of a hollow half-cylinder divided in its long axis.