

美国数学会经典影印系列



An Introduction to Measure Theory

测度论引论

Terence Tao



高等教育出版社

美国数学会经典影印系列



An Introduction to Measure Theory

测度论引论

Terence Tao



高等教育出版社·北京

图字：01-2016-2504 号

An Introduction to Measure Theory, by Terence Tao, first published by the American Mathematical Society.

Copyright © 2011 Terence Tao. All rights reserved.

This present reprint edition is published by Higher Education Press Limited Company under authority of the American Mathematical Society and is published under license.

Special Edition for People's Republic of China Distribution Only. This edition has been authorized by the American Mathematical Society for sale in People's Republic of China only, and is not for export therefrom.

本书最初由美国数学会于 2011 年出版，原书名为 *An Introduction to Measure Theory*，作者为 Terence Tao。Terence Tao 保留原书所有版权。

原书版权声明：Copyright © 2011 by Terence Tao。

本影印版由高等教育出版社有限公司经美国数学会独家授权出版。

本版只限于中华人民共和国境内发行。本版经由美国数学会授权仅在中华人民共和国境内销售，不得出口。

测度论引论

Cedulun Yinlun

图书在版编目 (CIP) 数据

测度论引论 = *An Introduction to Measure Theory* :

英文 / (澳) 陶哲轩 (Terence Tao) 著. —影印本.

—北京 : 高等教育出版社, 2017.1

ISBN 978-7-04-046905-9

I. ①测… II. ①陶… III. ①测度论—英文 IV.

①O174.12

中国版本图书馆 CIP 数据核字 (2016) 第 281914 号

策划编辑 赵天夫 责任编辑 赵天夫
封面设计 张申申 责任印制 毛斯璐

出版发行 高等教育出版社
社址 北京市西城区德外大街 4 号
邮政编码 100120

购书热线 010-58581118

咨询电话 400-810-0598

网址 <http://www.hep.edu.cn>

<http://www.hep.com.cn>

网上订购 <http://www.hepmall.com.cn>

<http://www.hepmall.com>

<http://www.hepmall.cn>

印刷:北京新华印刷有限公司

开本 787mm × 1092mm 1/16

印张 14.25

字数 290 千字

版次 2017 年 1 月第 1 版

印次 2017 年 1 月第 1 次印刷

定价 99.00 元

本书如有缺页、倒页、脱页等质量问题,

请到所购图书销售部门联系调换

版权所有 侵权必究

[物 料 号 46905-00]



美国数学会经典影印系列

出版者的话

近年来,我国的科学技术取得了长足进步,特别是在数学等自然科学基础领域不断涌现出一流的研究成果。与此同时,国内的科研队伍与国外的交流合作也越来越密切,越来越多的科研工作者可以熟练地阅读英文文献,并在国际顶级期刊发表英文学术文章,在国外出版社出版英文学术著作。

然而,在国内阅读海外原版英文图书仍不是非常便捷。一方面,这些原版图书主要集中在科技、教育比较发达的大中城市的大型综合图书馆以及科研院所的资料室中,普通读者借阅不甚容易;另一方面,原版书价格昂贵,动辄上百美元,购买也很不方便。这极大地限制了科技工作者对于国外先进科学技术知识的获取,间接阻碍了我国科技的发展。

高等教育出版社本着植根教育、弘扬学术的宗旨服务我国广大科技和教育工作者,同美国数学会(American Mathematical Society)合作,在征求海内外众多专家学者意见的基础上,精选该学会近年出版的数十种专业著作,组织出版了“美国数学会经典影印系列”丛书。美国数学会创建于1888年,是国际上极具影响力的专业学术组织,目前拥有近30000会员和580余个机构成员,出版图书3500多种,冯·诺依曼、莱夫谢茨、陶哲轩等世界级数学大家都是其作者。本影印系列涵盖了代数、几何、分析、方程、拓扑、概率、动力系统所有主要数学分支以及新近发展的数学主题。

我们希望这套书的出版,能够对国内的科研工作者、教育工作者以及青年学生起到重要的学术引领作用,也希望今后能有更多的海外优秀英文著作被介绍到中国。

高等教育出版社

2016年12月

To Garth Gaudry, who set me on the road;
To my family, for their constant support;
And to the readers of my blog, for their feedback and contributions.

Preface

In the fall of 2010, I taught an introductory one-quarter course on graduate real analysis, focusing in particular on the basics of measure and integration theory, both in Euclidean spaces and in abstract measure spaces. This text is based on my lecture notes of that course, which are also available online on my blog terrytao.wordpress.com, together with some supplementary material, such as a section on problem solving strategies in real analysis (Section 2.1) which evolved from discussions with my students.

This text is intended to form a prequel to my graduate text [Ta2010] (henceforth referred to as *An epsilon of room, Vol. I*), which is an introduction to the analysis of Hilbert and Banach spaces (such as L^p and Sobolev spaces), point-set topology, and related topics such as Fourier analysis and the theory of distributions; together, they serve as a text for a complete first-year graduate course in real analysis.

The approach to measure theory here is inspired by the text [StSk2005], which was used as a secondary text in my course. In particular, the first half of the course is devoted almost exclusively to measure theory on Euclidean spaces \mathbf{R}^d (starting with the more elementary Jordan-Riemann-Darboux theory, and only then moving on to the more sophisticated Lebesgue theory), deferring the abstract aspects of measure theory to the second half of the course. I found that this approach strengthened the student's intuition in the early stages of the course, and helped provide motivation for more abstract constructions, such as Carathéodory's general construction of a measure from an outer measure.

Most of the material here is self-contained, assuming only an undergraduate knowledge in real analysis (and in particular, on the Heine-Borel theorem, which we will use as the foundation for our construction of Lebesgue

measure); a secondary real analysis text can be used in conjunction with this one, but it is not strictly necessary. A small number of exercises, however, will require some knowledge of point-set topology or of set-theoretic concepts such as cardinals and ordinals.

A large number of exercises are interspersed throughout the text, and it is intended that the reader perform a significant fraction of these exercises while going through the text. Indeed, many of the key results and examples in the subject will in fact be presented through the exercises. In my own course, I used the exercises as the basis for the examination questions, and indicated this well in advance, to encourage the students to attempt as many of the exercises as they could as preparation for the exams.

The core material is contained in Chapter 1, and already comprises a full quarter's worth of material. Section 2.1 is a much more informal section than the rest of the book, focusing on describing problem solving strategies, either specific to real analysis exercises, or more generally, applicable to a wider set of mathematical problems; this section evolved from various discussions with students throughout the course. The remaining three sections in Chapter 2 are optional topics, which require understanding most of the material in Chapter 1 as a prerequisite (although Section 2.3 can be read after completing Section 1.4).

Notation

For reasons of space, we will not be able to define every single mathematical term that we use in this book. If a term is italicised for reasons other than emphasis or for definition, then it denotes a standard mathematical object, result, or concept, which can be easily looked up in any number of references. (In the blog version of the book, many of these terms were linked to their Wikipedia pages, or other on-line reference pages.)

Given a subset E of a space X , the *indicator function* $1_E : X \rightarrow \mathbf{R}$ is defined by setting $1_E(x)$ equal to 1 for $x \in E$ and equal to 0 for $x \notin E$.

For any natural number d , we refer to the vector space

$$\mathbf{R}^d := \{(x_1, \dots, x_d) : x_1, \dots, x_d \in \mathbf{R}\}$$

as *(d-dimensional) Euclidean space*. A vector (x_1, \dots, x_d) in \mathbf{R}^d has length

$$|(x_1, \dots, x_d)| := (x_1^2 + \dots + x_d^2)^{1/2}$$

and two vectors $(x_1, \dots, x_d), (y_1, \dots, y_d)$ have *dot product*

$$(x_1, \dots, x_d) \cdot (y_1, \dots, y_d) := x_1 y_1 + \dots + x_d y_d.$$

The *extended non-negative real axis* $[0, +\infty]$ is the non-negative real axis $[0, +\infty) := \{x \in \mathbf{R} : x \geq 0\}$ with an additional element adjoined to it, which we label $+\infty$; we will need to work with this system because many sets (e.g. \mathbf{R}^d) will have infinite measure. Of course, $+\infty$ is not a real number, but we think of it as an *extended* real number. We extend the addition, multiplication, and order structures on $[0, +\infty)$ to $[0, +\infty]$ by declaring

$$+\infty + x = x + +\infty = +\infty$$

for all $x \in [0, +\infty]$,

$$+\infty \cdot x = x \cdot +\infty = +\infty$$

for all non-zero $x \in (0, +\infty]$,

$$+\infty \cdot 0 = 0 \cdot +\infty = 0,$$

and

$$x < +\infty \text{ for all } x \in [0, +\infty).$$

Most of the laws of algebra for addition, multiplication, and order continue to hold in this extended number system; for instance, addition and multiplication are commutative and associative, with the latter distributing over the former, and an order relation $x \leq y$ is preserved under addition or multiplication of both sides of that relation by the same quantity. However, we caution that the laws of cancellation do *not* apply once some of the variables are allowed to be infinite; for instance, we cannot deduce $x = y$ from $+\infty + x = +\infty + y$ or from $+\infty \cdot x = +\infty \cdot y$. This is related to the fact that the forms $+\infty - +\infty$ and $+\infty / +\infty$ are indeterminate (one cannot assign a value to them without breaking many of the rules of algebra). A general rule of thumb is that if one wishes to use cancellation (or proxies for cancellation, such as subtraction or division), this is only safe if one can guarantee that all quantities involved are finite (and in the case of multiplicative cancellation, the quantity being cancelled also needs to be non-zero, of course). However, as long as one avoids using cancellation and works exclusively with non-negative quantities, there is little danger in working in the extended real number system.

We note also that once one adopts the convention $+\infty \cdot 0 = 0 \cdot +\infty = 0$, then multiplication becomes *upward continuous* (in the sense that whenever $x_n \in [0, +\infty]$ increases to $x \in [0, +\infty]$, and $y_n \in [0, +\infty]$ increases to $y \in [0, +\infty]$, then $x_n y_n$ increases to xy) but not *downward continuous* (e.g. $1/n \rightarrow 0$ but $1/n \cdot +\infty \not\rightarrow 0 \cdot +\infty$). This asymmetry will ultimately cause us to define integration from below rather than from above, which leads to other asymmetries (e.g. the monotone convergence theorem (Theorem 1.4.43) applies for monotone increasing functions, but not necessarily for monotone decreasing ones).

Remark 0.0.1. Note that there is a tradeoff here: if one wants to keep as many useful laws of algebra as one can, then one can add in infinity, or have negative numbers, but it is difficult to have both at the same time. Because of this tradeoff, we will see two overlapping types of measure and integration theory: the *non-negative* theory, which involves quantities taking values in $[0, +\infty]$, and the *absolutely integrable* theory, which involves quantities taking values in $(-\infty, +\infty)$ or \mathbf{C} . For instance, the fundamental convergence theorem for the former theory is the monotone convergence theorem (Theorem 1.4.43), while the fundamental convergence theorem for the latter is the dominated convergence theorem (Theorem 1.4.48). Both branches of the theory are important, and both will be covered in later notes.

One important feature of the extended non-negative real axis is that all sums are convergent: given any sequence $x_1, x_2, \dots \in [0, +\infty]$, we can always form the sum

$$\sum_{n=1}^{\infty} x_n \in [0, +\infty]$$

as the limit of the partial sums $\sum_{n=1}^N x_n$, which may be either finite or infinite. An equivalent definition of this infinite sum is as the supremum of all finite subsums:

$$\sum_{n=1}^{\infty} x_n = \sup_{F \subset \mathbf{N}, F \text{ finite}} \sum_{n \in F} x_n.$$

Motivated by this, given any collection $(x_\alpha)_{\alpha \in A}$ of numbers $x_\alpha \in [0, +\infty]$ indexed by an arbitrary set A (finite or infinite, countable or uncountable), we can define the sum $\sum_{\alpha \in A} x_\alpha$ by the formula

$$(0.1) \quad \sum_{\alpha \in A} x_\alpha = \sup_{F \subset A, F \text{ finite}} \sum_{\alpha \in F} x_\alpha.$$

Note from this definition that one can relabel the collection in an arbitrary fashion without affecting the sum; more precisely, given any bijection $\phi : B \rightarrow A$, one has the change of variables formula

$$(0.2) \quad \sum_{\alpha \in A} x_\alpha = \sum_{\beta \in B} x_{\phi(\beta)}.$$

Note that when dealing with signed sums, the above rearrangement identity can fail when the series is not absolutely convergent (cf. the *Riemann rearrangement theorem*).

Exercise 0.0.1. If $(x_\alpha)_{\alpha \in A}$ is a collection of numbers $x_\alpha \in [0, +\infty]$ such that $\sum_{\alpha \in A} x_\alpha < \infty$, show that $x_\alpha = 0$ for all but at most countably many $\alpha \in A$, even if A itself is uncountable.

We will rely frequently on the following basic fact (a special case of the *Fubini-Tonelli theorem*, Corollary 1.7.23):

Theorem 0.0.2 (Tonelli's theorem for series). *Let $(x_{n,m})_{n,m \in \mathbf{N}}$ be a doubly infinite sequence of extended non-negative reals $x_{n,m} \in [0, +\infty]$. Then*

$$\sum_{(n,m) \in \mathbf{N}^2} x_{n,m} = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x_{n,m} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} x_{n,m}.$$

Informally, Tonelli's theorem asserts that we may rearrange infinite series with impunity as long as all summands are non-negative.

Proof. We shall just show the equality of the first and second expressions; the equality of the first and third is proven similarly.

We first show that

$$\sum_{(n,m) \in \mathbf{N}^2} x_{n,m} \leq \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x_{n,m}.$$

Let F be any finite subset of \mathbf{N}^2 . Then $F \subset \{1, \dots, N\} \times \{1, \dots, N\}$ for some finite N , and thus (by the non-negativity of the $x_{n,m}$)

$$\sum_{(n,m) \in F} x_{n,m} \leq \sum_{(n,m) \in \{1, \dots, N\} \times \{1, \dots, N\}} x_{n,m}.$$

The right-hand side can be rearranged as

$$\sum_{n=1}^N \sum_{m=1}^N x_{n,m},$$

which is clearly at most $\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x_{n,m}$ (again by non-negativity of $x_{n,m}$). This gives

$$\sum_{(n,m) \in F} x_{n,m} \leq \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x_{n,m},$$

for any finite subset F of \mathbf{N}^2 , and the claim then follows from (0.1).

It remains to show the reverse inequality

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} x_{n,m} \leq \sum_{(n,m) \in \mathbf{N}^2} x_{n,m}.$$

It suffices to show that

$$\sum_{n=1}^N \sum_{m=1}^{\infty} x_{n,m} \leq \sum_{(n,m) \in \mathbf{N}^2} x_{n,m}$$

for each finite N .

Fix N . As each $\sum_{m=1}^{\infty} x_{n,m}$ is the limit of $\sum_{m=1}^M x_{n,m}$, the left-hand side is the limit of $\sum_{n=1}^N \sum_{m=1}^M x_{n,m}$ as $M \rightarrow \infty$. Thus it suffices to show that

$$\sum_{n=1}^N \sum_{m=1}^M x_{n,m} \leq \sum_{(n,m) \in \mathbf{N}^2} x_{n,m}$$

for each finite M . But the left-hand side is $\sum_{(n,m) \in \{1, \dots, N\} \times \{1, \dots, M\}} x_{n,m}$, and the claim follows. \square

Remark 0.0.3. Note how important it was that the $x_{n,m}$ were non-negative in the above argument. In the signed case, one needs an additional assumption of absolute summability of $x_{n,m}$ on \mathbf{N}^2 before one is permitted to interchange sums; this is *Fubini's theorem for series*, which we will encounter later in this text. Without absolute summability or non-negativity hypotheses, the theorem can fail (consider, for instance, the case when $x_{n,m}$ equals $+1$ when $n = m$, -1 when $n = m + 1$, and 0 otherwise).

Exercise 0.0.2 (Tonelli's theorem for series over arbitrary sets). Let A, B be sets (possibly infinite or uncountable), and $(x_{n,m})_{n \in A, m \in B}$ be a doubly infinite sequence of extended non-negative reals $x_{n,m} \in [0, +\infty]$ indexed by A and B . Show that

$$\sum_{(n,m) \in A \times B} x_{n,m} = \sum_{n \in A} \sum_{m \in B} x_{n,m} = \sum_{m \in B} \sum_{n \in A} x_{n,m}.$$

(*Hint:* Although not strictly necessary, you may find it convenient to first establish the fact that if $\sum_{n \in A} x_n$ is finite, then x_n is non-zero for at most countably many n .)

Next, we recall the *axiom of choice*, which we shall be assuming throughout the text:

Axiom 0.0.4 (Axiom of choice). Let $(E_\alpha)_{\alpha \in A}$ be a family of non-empty sets E_α , indexed by an index set A . Then we can find a family $(x_\alpha)_{\alpha \in A}$ of elements x_α of E_α , indexed by the same set A .

This axiom is trivial when A is a singleton set, and from mathematical induction one can also prove it without difficulty when A is finite. However, when A is infinite, one cannot deduce this axiom from the other axioms of set theory, but must explicitly add it to the list of axioms. We isolate the countable case as a particularly useful corollary (though one which is strictly weaker than the full axiom of choice):

Corollary 0.0.5 (Axiom of countable choice). Let E_1, E_2, E_3, \dots be a sequence of non-empty sets. Then one can find a sequence x_1, x_2, \dots such that $x_n \in E_n$ for all $n = 1, 2, 3, \dots$

Remark 0.0.6. The question of how much of real analysis still survives when one is not permitted to use the axiom of choice is a delicate one, involving a fair amount of logic and descriptive set theory to answer. We will not discuss these matters in this text. We will, however, note a theorem of Gödel [Go1938] that states that any statement that can be phrased in the first-order language of *Peano arithmetic*, and which is proven with the axiom of choice, can also be proven without the axiom of choice. So, roughly speaking, Gödel's theorem tells us that for any “finitary” application of real analysis (which includes most of the “practical” applications of the subject), it is safe to use the axiom of choice; it is only when asking questions about “infinitary” objects that are beyond the scope of Peano arithmetic that one can encounter statements that are provable using the axiom of choice, but are not provable without it.

Acknowledgments

This text was strongly influenced by the real analysis text of Stein and Shakarchi [StSk2005], which was used as a secondary text when teaching the course on which these notes were based. In particular, the strategy of focusing first on Lebesgue measure and Lebesgue integration, before moving onwards to abstract measure and integration theory, was directly inspired by the treatment in [StSk2005], and the material on differentiation theorems also closely follows that in [StSk2005]. On the other hand, our discussion here differs from that in [StSk2005] in other respects; for instance, a far greater emphasis is placed on Jordan measure and the Riemann integral as being an elementary precursor to Lebesgue measure and the Lebesgue integral.

I am greatly indebted to my students of the course on which this text was based, as well as many further commenters on my blog, including Maria Alfonseca, Marco Angulo, J. Balachandran, Farzin Barekat, Marek Bernát, Lewis Bowen, Chris Breeden, Danny Calegari, Yu Cao, Chandrasekhar, David Chang, Nick Cook, Damek Davis, Eric Davis, Marton Eekes, Wenying Gan, Nick Gill, Ulrich Groh, Tim Gowers, Laurens Gunnarsen, Tobias Hagge, Xueping Huang, Bo Jacoby, Apoorva Khare, Shiping Liu, Colin McQuillan, David Milovich, Hossein Naderi, Brent Nelson, Constantin Niculescu, Mircea Petrache, Walt Pohl, Jim Ralston, David Roberts, Mark Schwarzmann, Vladimir Slepnev, David Speyer, Blake Stacey, Tim Sullivan, Jonathan Weinstein, Duke Zhang, Lei Zhang, Pavel Zorin, and several anonymous commenters, for providing corrections and useful commentary on the material here. These comments can be viewed online at:

terrytao.wordpress.com/category/teaching/245a-real-analysis

The author is supported by a grant from the MacArthur Foundation, by NSF grant DMS-0649473, and by the NSF Waterman award.

郑重声明

高等教育出版社依法对本书享有专有出版权。任何未经许可的复制、销售行为均违反《中华人民共和国著作权法》，其行为人将承担相应的民事责任和行政责任；构成犯罪的，将被依法追究刑事责任。为了维护市场秩序，保护读者的合法权益，避免读者误用盗版书造成不良后果，我社将配合行政执法部门和司法机关对违法犯罪的单位和个人进行严厉打击。社会各界人士如发现上述侵权行为，希望及时举报，本社将奖励举报有功人员。

反盗版举报电话 (010) 58581999 58582371 58582488

反盗版举报传真 (010) 82086060

反盗版举报邮箱 dd@hep.com.cn

通信地址 北京市西城区德外大街4号

高等教育出版社法律事务与版权管理部

邮政编码 100120

美国数学会经典影印系列

1. **Lars V. Ahlfors**, Lectures on Quasiconformal Mappings, Second Edition, 978-7-04-047010-9
2. **Dmitri Burago, Yuri Burago, Sergei Ivanov**, A Course in Metric Geometry, 978-7-04-046908-0
3. **Tobias Holck Colding, William P. Minicozzi II**, A Course in Minimal Surfaces, 978-7-04-046911-0
4. **Javier Duoandikoetxea**, Fourier Analysis, 978-7-04-046901-1
5. **John P. D'Angelo**, An Introduction to Complex Analysis and Geometry, 978-7-04-046998-1
6. **Y. Eliashberg, N. Mishachev**, Introduction to the h -Principle, 978-7-04-046902-8
7. **Lawrence C. Evans**, Partial Differential Equations, Second Edition, 978-7-04-046935-6
8. **Robert E. Greene, Steven G. Krantz**,
Function Theory of One Complex Variable, Third Edition, 978-7-04-046907-3
9. **Thomas A. Ivey, J. M. Landsberg**,
Cartan for Beginners: Differential Geometry via Moving Frames and
Exterior Differential Systems, 978-7-04-046917-2
10. **Jens Carsten Jantzen**, Representations of Algebraic Groups, Second Edition, 978-7-04-047008-6
11. **A. A. Kirillov**, Lectures on the Orbit Method, 978-7-04-046910-3
12. **Jean-Marie De Koninck, Armel Mercier**, 1001 Problems in Classical Number Theory, 978-7-04-046999-8
13. **Peter D. Lax, Lawrence Zalcman**, Complex Proofs of Real Theorems, 978-7-04-047000-0
14. **David A. Levin, Yuval Peres, Elizabeth L. Wilmer**, Markov Chains and Mixing Times, 978-7-04-046994-3
15. **Dusa McDuff, Dietmar Salamon**, J -holomorphic Curves and Symplectic Topology, 978-7-04-046993-6
16. **John von Neumann**, Invariant Measures, 978-7-04-046997-4
17. **R. Clark Robinson**,
An Introduction to Dynamical Systems: Continuous and Discrete, Second Edition, 978-7-04-047009-3
18. **Terence Tao**,
An Epsilon of Room, I: Real Analysis: pages from year three of a mathematical blog, 978-7-04-046900-4
19. **Terence Tao**, An Epsilon of Room, II: pages from year three of a mathematical blog, 978-7-04-046899-1
20. **Terence Tao**, An Introduction to Measure Theory, 978-7-04-046905-9
21. **Terence Tao**, Higher Order Fourier Analysis, 978-7-04-046909-7
22. **Terence Tao**, Poincaré's Legacies, Part I: pages from year two of a mathematical blog, 978-7-04-046995-0
23. **Terence Tao**, Poincaré's Legacies, Part II: pages from year two of a mathematical blog, 978-7-04-046996-7
24. **Cédric Villani**, Topics in Optimal Transportation, 978-7-04-046921-9
25. **R. J. Williams**, Introduction to the Mathematics of Finance, 978-7-04-046912-7

Contents

Preface	ix
Notation	x
Acknowledgments	xv
Chapter 1. Measure theory	1
§1.1. Prologue: The problem of measure	2
§1.2. Lebesgue measure	15
§1.3. The Lebesgue integral	38
§1.4. Abstract measure spaces	66
§1.5. Modes of convergence	94
§1.6. Differentiation theorems	108
§1.7. Outer measures, pre-measures, and product measures	148
Chapter 2. Related articles	173
§2.1. Problem solving strategies	174
§2.2. The Rademacher differentiation theorem	187
§2.3. Probability spaces	192
§2.4. Infinite product spaces and the Kolmogorov extension theorem	194
Bibliography	201
Index	203