

美国数学会经典影印系列

 **AMS**
AMERICAN MATHEMATICAL SOCIETY

J-holomorphic Curves and Symplectic Topology

Second Edition

J-全纯曲线和辛拓扑

第二版

Dusa McDuff, Dietmar Salamon



高等教育出版社

美国数学会经典影印系列



J -holomorphic Curves and Symplectic Topology

Second Edition

J -全纯曲线和辛拓扑

第二版

Dusa McDuff, Dietmar Salamon



高等教育出版社·北京

图字: 01-2016-3517 号

J-holomorphic Curves and Symplectic Topology, Second Edition, by Dusa McDuff and Dietmar Salamon, first published by the American Mathematical Society.

Second Edition © 2012 by the American Mathematical Society. All rights reserved.

This present reprint edition is published by Higher Education Press Limited Company under authority of the American Mathematical Society and is published under license.

Special Edition for People's Republic of China Distribution Only. This edition has been authorized by the American Mathematical Society for sale in People's Republic of China only, and is not for export therefrom.

本书第二版由美国数学会于2012年出版,原书名为*J-holomorphic Curves and Symplectic Topology, Second Edition*,原书作者为Dusa McDuff and Dietmar Salamon。美国数学会保留原书所有版权。

原书版权声明: Second Edition © 2012 by the American Mathematical Society。

本影印版由高等教育出版社有限公司经美国数学会独家授权出版。

本版只限于中华人民共和国境内发行。本版经由美国数学会授权仅在中华人民共和国境内销售,不得出口。

J-全纯曲线和辛拓扑

第二版

J-Quanchun Quxian he

Xintuopu

图书在版编目(CIP)数据

J-全纯曲线和辛拓扑 = J-holomorphic Curves and Symplectic Topology : 第二版 : 英文 / (英) 杜莎·麦克达夫 (Dusa McDuff), (德) 迪特马尔·萨拉蒙 (Dietmar Salamon) 著. —影印本. —北京: 高等教育出版社, 2017.4

ISBN 978-7-04-046993-6

I. ①J… II. ①杜… ②迪… III. ①曲线—英文②拓扑—英文 IV. ①O123.3 ②O189

中国版本图书馆 CIP 数据核字 (2016) 第 326751 号

策划编辑 李 鹏 责任编辑 李 鹏
封面设计 张申申 责任印制 毛斯璐

出版发行 高等教育出版社
社址 北京市西城区德外大街4号
邮政编码 100120
购书热线 010-58581118
咨询电话 400-810-0598
网址 <http://www.hep.edu.cn>
<http://www.hep.com.cn>

网上订购 <http://www.hepmall.com.cn>
<http://www.hepmall.com>
<http://www.hepmall.cn>
印刷 北京新华印刷有限公司

开本 787mm×1092mm 1/16
印张 46.75
字数 1160千字
版次 2017年4月第1版
印次 2017年4月第1次印刷
定价 199.00元

本书如有缺页、倒页、脱页等质量问题,
请到所购图书销售部门联系调换
版权所有 侵权必究
[物料号 46993-00]



美国数学会经典影印系列

出版者的话

近年来,我国的科学技术取得了长足进步,特别是在数学等自然科学基础领域不断涌现出一流的研究成果。与此同时,国内的科研队伍与国外的交流合作也越来越密切,越来越多的科研工作者可以熟练地阅读英文文献,并在国际顶级期刊发表英文学术文章,在国外出版社出版英文学术著作。

然而,在国内阅读海外原版英文图书仍不是非常便捷。一方面,这些原版图书主要集中在科技、教育比较发达的大中城市的大型综合图书馆以及科研院所的资料室中,普通读者借阅不甚容易;另一方面,原版书价格昂贵,动辄上百美元,购买也很不方便。这极大地限制了科技工作者对于国外先进科学技术知识的获取,间接阻碍了我国科技的发展。

高等教育出版社本着植根教育、弘扬学术的宗旨服务我国广大科技和教育工作者,同美国数学会(American Mathematical Society)合作,在征求海内外众多专家学者意见的基础上,精选该学会近年出版的数十种专业著作,组织出版了“美国数学会经典影印系列”丛书。美国数学会创建于1888年,是国际上极具影响力的专业学术组织,目前拥有近30000会员和580余个机构成员,出版图书3500多种,冯·诺依曼、莱夫谢茨、陶哲轩等世界级数学大家都是其作者。本影印系列涵盖了代数、几何、分析、方程、拓扑、概率、动力系统所有主要数学分支以及新近发展的数学主题。

我们希望这套书的出版,能够对国内的科研工作者、教育工作者以及青年学生起到重要的学术引领作用,也希望今后能有更多的海外优秀英文著作被介绍到中国。

高等教育出版社

2016年12月

Preface to the second edition

This revision has two main purposes: first to correct various errors that crept into the first edition and second to update our discussions of current work in the field. Since the first edition of this book appeared in 2004, symplectic geometry has developed apace. It has found new applications in low dimensional topology, via Heegaard Floer theory [318] and the newly understood relations of embedded contact homology to gauge theory [74, 219]. Several important books have been published that develop powerful new ideas and techniques: Seidel [371] on the Fukaya category, Fukaya–Oh–Ohta–Ono [128] on Lagrangian Floer homology, and Cieliebak and Eliashberg [63] on the relations between complex and symplectic manifolds. Another exciting development is the introduction of sheaf-theoretic methods for proving fundamental rigidity results in symplectic geometry by Tamarkin [388] and Guillermou–Kashiwara–Shapira [166]. There has also been great progress on particular problems; for example Taubes [394] solved the Weinstein conjecture for 3-dimensional contact manifolds using Seiberg–Witten–Floer theory, Hingston [176] and Ginzburg [143] solved the Conley conjecture by new advances in Hamiltonian dynamics and Floer theory, and the nearby Lagrangian conjecture has been partially solved (by Fukaya–Seidel–Smith [131] and Abouzaid [1] among others) using Fukaya categories. A comprehensive exposition of Hamiltonian Floer theory is now available with the book by Audin–Damian [24], which presents all the basic analysis needed to set up Hamiltonian Floer theory for manifolds with $c_1 = 0$ as well as in the monotone case. Finally, the long series of papers and books by Hofer–Wysocki–Zehnder [184, 185, 186, 187, 188, 189] develops a new functional analytic approach to the theory of J -holomorphic curves. Their work will eventually give solid foundations to Lagrangian Floer theory and the various forms of Symplectic Field Theory.

We do not say much about the details of these developments. However, we have updated the introductions to the chapters where relevant, and also have extended the discussions of various applications of J -holomorphic curves in Chapters 9, 11 and 12, aiming to give a sense of the main new developments and the main new players rather than to be comprehensive.

Many of the corrections are rather minor. However, we have rewritten Section 4.4 on the isoperimetric inequality, the proof of Theorem 7.2.3, the proof of Proposition 7.4.8, and the proof of the sum formula for the Fredholm index in Theorem C.4.2. In Chapter 10 we added Section 10.9 with a new geometric formulation of the gluing theorem for z -independent almost complex structures, in Appendix C we expanded Section C.5 to include a proof of integrability of almost complex structures in dimension two, and in Appendix D we expanded Section D.4 to include the material previously in Sections D.4 and D.5 and added a new Section D.6 on the cohomology of the moduli space of stable curves of genus zero.

We warmly thank everyone who pointed out mistakes in the earlier edition, but particularly Aleksei Zinger who sent us an especially thorough and useful list of comments.

Dusa McDuff and Dietmar Salamon, April 2012

Preface

The theory of J -holomorphic curves has been of great importance to symplectic topologists ever since its inception in Gromov's paper of 1985. Its applications include many key results in symplectic topology, and it was one of the main inspirations for the creation of Floer homology. It has caught the attention of mathematical physicists since it provides a natural context in which to define Gromov–Witten invariants and quantum cohomology, which form the so-called A-side of the mirror symmetry conjecture. Insights from physics have in turn inspired many fascinating developments, for example highlighting as yet little understood connections between the theory of integrable systems and Gromov–Witten invariants.

Several years ago the two authors of this book wrote an expository account of the field that explained the main technical steps in the theory of J -holomorphic curves. The present book started life as a second edition of that book, but the project quickly grew. The field has been developing so rapidly that there has been little time to consolidate its foundations. Since these involve many analytic subtleties, this has proved quite a hindrance. Therefore the main aim of this book is to establish the fundamental theorems in the subject in full and rigorous detail. We also hope that the book will serve as an introduction to current work in symplectic topology. These two aims are, of course, somewhat in conflict, and in different parts of the book different aspects are predominant.

We have chosen to concentrate on setting up the foundations of the theory rather than attempting to cover the many recent developments in detail. Thus, we limit ourselves to genus zero curves (though we do treat discs as well as spheres). A more serious limitation is that we restrict ourselves to the semipositive case, where it is possible to define the Gromov–Witten invariants in terms of pseudocycles. Our main reason for doing this is that an optimal framework for the general case (which would involve constructing a virtual moduli cycle) has not yet been worked out. Rather than cobbling together a definition that would do for many applications but not suffice in broader contexts such as symplectic field theory, we decided to show what can be done with a simpler, more geometric approach. On the other hand, we give a very detailed proof of the basic gluing theorem. This is the analytic foundation for all subsequent work on the virtual moduli cycle and is the essential ingredient in the proof of the associativity of quantum multiplication. There are also five extensive appendices, on topics ranging from standard results such as the implicit function theorem, elliptic regularity and the Riemann–Roch theorem to lesser known subjects such as the structure of the moduli space of genus zero stable curves and positivity of intersections for J -holomorphic curves in dimension four. We have adopted the same approach to the applications, giving complete proofs of the foundational results and illustrating more recent developments by describing some key examples and giving a copious list of references.

The book is written so that the subject develops in logical order. Chapters 2 through 5 establish the foundational Fredholm theory and compactness results for J -holomorphic spheres and discs; Chapter 6 introduces the concepts need to define the Gromov–Witten pseudocycle for semipositive manifolds; Chapter 7 is the pivotal chapter in which the invariants are defined; and the later chapters discuss various applications. Since there is more detail in Chapters 2 through 6 than can possibly be absorbed at a first reading, we have written the introductory Chapter 1 to describe the outlines of the theory and to introduce the main definitions. It serves as a detailed guide to this book, pointing out where the key arguments occur and where to find the background details needed to understand various examples. Each chapter also has an introduction describing its main contents, which should help to orient the more knowledgeable readers. Wherever possible we have written the sections and chapters to be independent of each other. Hence the reader should feel free to skip parts that seem excessively technical.

We hope that Chapter 1 (supplemented by suitable parts of Chapters 2–6) will provide beginning students with enough of the essential background for understanding the main definitions in Chapter 7. Here is a brief outline of the contents of the remaining chapters. After the basic invariants are defined in Section 7.1 (with important supplemental ideas in Section 7.2 and Section 7.3), Section 7.4 discusses the fundamental example of rational curves in projective space. The chapter ends with a discussion of the Kontsevich–Manin axioms for the genus zero Gromov–Witten invariants, and deduces from them Kontsevich’s beautiful iterative formula for the number of degree d rational curves in the projective plane.

Chapter 8 sets up the theory of locally Hamiltonian fibrations over Riemann surfaces and shows how to count sections of such fibrations. This allows us to define Gromov–Witten invariants of arbitrary genus (but where the complex structure of the domain is fixed). It also provides the background for some important applications, for example Gromov’s result that every Hamiltonian system on a symplectically aspherical manifold has a 1-periodic orbit (see Theorem 9.1.1), and results about the group of Hamiltonian symplectomorphisms: a taste of Hofer geometry in Section 9.6 and a discussion of the Seidel representation in Sections 11.4 and 12.5.

Chapter 9 describes some of the main applications of J -holomorphic curve techniques in symplectic geometry. Besides the examples mentioned above and a discussion of the basic properties of Lagrangian submanifolds, it gives full proofs of McDuff’s results on the structure of rational and ruled symplectic 4-manifolds as well as Gromov’s results on the symplectomorphism group of the projective plane and the product of 2-spheres.

The other main application, quantum cohomology, requires a further deep analytic technique, that of gluing. The first rigorous gluing arguments are due to Floer (in the somewhat easier context of Floer homology) and Ruan–Tian (in the context relevant to quantum homology). In Chapter 10 we present a different, perhaps easier, method of gluing and derive from it a proof of the splitting axiom for the Gromov–Witten invariants in semipositive manifolds.

With this in hand, Chapter 11 defines quantum cohomology and explains some of the structures arising from it, such as the Gromov–Witten potential and Frobenius manifolds. As is clear from the examples in Section 11.3, this is the place where symplectic topology makes the deepest contact with other areas such as algebraic geometry, conformal field theory, mirror symmetry, and integrable systems. This

chapter should be accessible after Chapter 7. Finally, Chapter 12 is a survey that formulates the main outlines of Floer theory, omitting the analytic underpinnings. It explains the relations between Floer theory and quantum cohomology, using a geometric approach, and also indicates the directions of further developments, both analytic (the vortex equations) and geometric (Donaldson's quantum category).

There are five appendices. The first three set up the foundations of the classical theory of linear elliptic operators that is generalized in Chapters 3 and 4: Fredholm theory and the implicit function theorem for Banach manifolds in Appendix A, Sobolev spaces and elliptic regularity in Appendix B, and the Riemann–Roch theorem for Riemann surfaces with boundary in Appendix C. Appendix D provides background for Chapter 5. It explains the structure of the Grothendieck–Knudson moduli space of genus zero stable curves using cross ratios rather than the usual algebro-geometric approach. Appendix E was written jointly with Laurent Lazzarini. It contains a complete proof of positivity of intersections and the adjunction inequality for J -holomorphic curves in four-dimensional manifolds. Lazzarini provided the first draft of this appendix with complete proofs and we then worked together on the exposition. The results of Appendix E provide the basis for the structure theorems for rational and ruled symplectic 4-manifolds.

Those who wish to use this book as the basis for a graduate course must make some firm decisions about what kind of course they want to teach. As we know from experience, it is impossible in one semester to prove all the main analytic techniques as well as to describe interesting examples. One possibility, explained in more detail in Chapter 1, would be to concentrate on Chapter 1 (amplified by small parts of Chapter 2), Chapter 3 through Section 3.3 (together perhaps with some extra analysis from Appendices B and C), the basic compactness result for spheres with minimal energy in Section 4.2, very selected parts of Chapter 6 (the definition of pseudocycle), and then move to Section 7.1. Then either one could go directly to some of the geometric applications in Chapter 9 (for example, prove the nonsqueezing theorem or some of the results about symplectic 4-manifolds in Section 9.4) or one could discuss the Kontsevich–Manin axioms for Gromov–Witten invariants in Section 7.5 and then move to Chapter 11 to set up quantum cohomology. The idea here would be to develop a familiarity with the main analytic setup, prove some of the basic techniques, and then set them in context by discussing one set of applications

The above outline is perhaps still too ambitious, but there are ways to shorten the preliminaries. For example, it is possible to discuss many of the applications in Chapter 9 directly after the foundational material of Chapters 2–4 (and relevant parts of Chapter 8), without any reference to Chapters 5, 6 and 7. For if one considers only the simplest cases of these applications, rather than proving them in their most general form, the relevant moduli spaces are compact and so the results become accessible without any formal definition of the Gromov–Witten invariants. Alternatively, those aiming at quantum cohomology could state the results on Fredholm theory without proof and instead concentrate on explaining some of the compactness (bubbling) results in Chapters 4 and 5. These combine well with a study of the moduli space of stable maps and hence lead naturally to the Kontsevich–Manin axioms.

As indicated above, a first course, unless it moves incredibly fast or contains almost no applications, cannot both cover Fredholm theory and come to grips with

the analytic details of the compactness proof, even less go through all the details of gluing. Even though this proof in the main needs the same analytic background as Chapter 3, the proof of the surjectivity of the gluing map hinges on the deepest result from Chapter 4 (the behaviour of long cylinders with small energy) and relies on several technical estimates. We have written the gluing chapter to try to make accessible the outlines of the construction, together with the main analytic ideas. (These are summarized in Section 10.5.) Hence, for a more analytically sophisticated audience, one might base a course on Chapters 3, 4 and 10, with motivation taken from some of the examples in Chapter 9 or 11.

Despite the length of this book, its subject is so rich that it is impossible to treat all its aspects. We have given many references throughout. Here are some books on related areas that the reader might wish to consult both on their own account and for the further references that they contain: Cox–Katz [76] on mirror symmetry and algebraic geometry, Donaldson [87] on Floer homology and gauge theory, Manin [286] on Frobenius manifolds and quantum cohomology, Polterovich [330] on the geometry of the symplectomorphism group, and the paper by Eliashberg–Givental–Hofer [101] on symplectic field theory.

This book has been long in the making and would not have been possible without help from many colleagues who shared their insights and knowledge with us. In particular, Coates, Givental, Lalonde, Lazzarini, Polterovich, Popescu, Robbin, Ruan, and Seidel all gave crucial help with various parts of this book. We also wish to thank the many students and others who pointed out various typos and inaccuracies, and especially Eduardo Gonzalez, Sam Lisi, Jake Solomon, and Fabian Ziltener for their meticulous attention to detail.

Dusa McDuff and Dietmar Salamon, December 2003

Contents

Preface to the second edition	ix
Preface	xi
Chapter 1. Introduction	1
1.1. Symplectic manifolds	1
1.2. Moduli spaces: regularity and compactness	4
1.3. Evaluation maps and pseudocycles	7
1.4. The Gromov–Witten invariants	10
1.5. Applications and further developments	13
Chapter 2. J -holomorphic Curves	17
2.1. Almost complex structures	17
2.2. The nonlinear Cauchy-Riemann equations	19
2.3. Unique continuation	21
2.4. Critical points	26
2.5. Somewhere injective curves	30
2.6. The adjunction inequality	35
Chapter 3. Moduli Spaces and Transversality	39
3.1. Moduli spaces of simple curves	39
3.2. Transversality	49
3.3. A regularity criterion	56
3.4. Curves with pointwise constraints	61
3.5. Implicit function theorem	68
Chapter 4. Compactness	75
4.1. Energy	76
4.2. The bubbling phenomenon	80
4.3. The mean value inequality	85
4.4. The isoperimetric inequality	91
4.5. Removal of singularities	96
4.6. Convergence modulo bubbling	99
4.7. Bubbles connect	105
Chapter 5. Stable Maps	115
5.1. Stable maps	115
5.2. Gromov convergence	122
5.3. Gromov compactness	126
5.4. Uniqueness of the limit	134
5.5. Gromov compactness for stable maps	139

5.6. The Gromov topology	147
Chapter 6. Moduli Spaces of Stable Maps	153
6.1. Simple stable maps	155
6.2. Transversality for simple stable maps	158
6.3. Transversality for evaluation maps	165
6.4. Semipositivity	168
6.5. Pseudocycles	171
6.6. Gromov–Witten pseudocycles	177
6.7. The pseudocycle of graphs	182
Chapter 7. Gromov–Witten Invariants	201
7.1. Counting pseudoholomorphic spheres	203
7.2. Variations on the definition	210
7.3. Counting pseudoholomorphic graphs	220
7.4. Rational curves in projective spaces	225
7.5. Axioms for Gromov–Witten invariants	239
Chapter 8. Hamiltonian Perturbations	257
8.1. Trivial bundles	258
8.2. Locally Hamiltonian fibrations	264
8.3. Pseudoholomorphic sections	270
8.4. Pseudoholomorphic spheres in the fiber	277
8.5. The pseudocycle of sections	279
8.6. Counting pseudoholomorphic sections	285
Chapter 9. Applications in Symplectic Topology	295
9.1. Periodic orbits of Hamiltonian systems	296
9.2. Obstructions to Lagrangian embeddings	310
9.3. The nonsqueezing theorem	323
9.4. Symplectic 4-manifolds	329
9.5. The group of symplectomorphisms	344
9.6. Hofer geometry	353
9.7. Distinguishing symplectic structures	359
Chapter 10. Gluing	369
10.1. The gluing theorem	370
10.2. Connected sums of J -holomorphic curves	373
10.3. Weighted norms	376
10.4. Cutoff functions	380
10.5. Construction of the gluing map	382
10.6. The derivative of the gluing map	392
10.7. Surjectivity of the gluing map	400
10.8. Proof of the splitting axiom	406
10.9. The gluing theorem revisited	413
Chapter 11. Quantum Cohomology	417
11.1. The small quantum cohomology ring	418
11.2. The Gromov–Witten potential	436
11.3. Four examples	442

11.4.	The Seidel representation	467
11.5.	Frobenius manifolds	478
Chapter 12.	Floer Homology	487
12.1.	Floer's cochain complex	488
12.2.	Ring structure	499
12.3.	Poincaré duality	503
12.4.	Spectral invariants	505
12.5.	The Seidel representation	514
12.6.	Donaldson's quantum category	519
12.7.	The symplectic vortex equations	524
Appendix A.	Fredholm Theory	531
A.1.	Fredholm theory	531
A.2.	Determinant line bundles	533
A.3.	The implicit function theorem	538
A.4.	Finite dimensional reduction	545
A.5.	The Sard–Smale theorem	547
Appendix B.	Elliptic Regularity	549
B.1.	Sobolev spaces	549
B.2.	The Calderon–Zygmund inequality	562
B.3.	Regularity for the Laplace operator	568
B.4.	Elliptic bootstrapping	571
Appendix C.	The Riemann–Roch Theorem	579
C.1.	Cauchy–Riemann operators	579
C.2.	Elliptic estimates	586
C.3.	The boundary Maslov index (by Joel Robbin)	593
C.4.	Proof of the Riemann–Roch theorem	598
C.5.	The Riemann mapping theorem	604
C.6.	Nonsmooth bundles	613
C.7.	Almost complex structures	614
Appendix D.	Stable Curves of Genus Zero	619
D.1.	Möbius transformations and cross ratios	619
D.2.	Trees, labels, and splittings	622
D.3.	Stable curves	629
D.4.	The Grothendieck–Knudsen manifold	631
D.5.	The Gromov topology	640
D.6.	Cohomology	643
D.7.	Examples	648
Appendix E.	Singularities and Intersections (written with Laurent Lazzarini)	653
E.1.	The main results	654
E.2.	Positivity of intersections	658
E.3.	Integrability	664
E.4.	The Hartman–Wintner theorem	668
E.5.	Local behaviour	673

E.6. Contact between branches	678
E.7. Singularities of J -holomorphic curves	686
Bibliography	695
List of Symbols	711
Index	715

CHAPTER 1

Introduction

The theory of J -holomorphic curves, introduced by Gromov in 1985, has profoundly influenced the study of symplectic geometry, and now permeates almost all its aspects. The methods are also of interest in the study of Kähler manifolds, since often when one studies properties of holomorphic curves in such manifolds it is useful to perturb the complex structure to be generic. The effect of this is to ensure that one is looking at persistent rather than accidental features of these curves. However, the perturbed structure may no longer be integrable, and so again one is led to the study of curves that are holomorphic with respect to some nonintegrable almost complex structure J .

These curves satisfy a nonlinear analogue of the Cauchy–Riemann equations. Before one can do anything useful with them, one must understand the elements of the theory of these equations; for example, know what conditions ensure that the solution spaces are finite dimensional manifolds and know ways of dealing with the fact that these solution spaces are usually noncompact. As explained in the preface, this chapter introduces all the basic concepts and outlines the ingredients needed to establish these results. Readers, specially those unfamiliar with the theory, should start by reading this chapter and then branch out into more detailed study of whichever aspects of the theory interests them.

We assume that the reader is familiar with the elements of symplectic geometry. There are several introductory books. One possible reference is McDuff–Salamon [277], but there are now more elementary treatments such as Berndt [35] and Cannas da Silva [59] as well as classics such as Arnold [15].

1.1. Symplectic manifolds

A symplectic structure on a smooth $2n$ -dimensional manifold M is a closed 2-form ω which is nondegenerate in the sense that the top-dimensional form ω^n does not vanish anywhere. By Darboux’s theorem, all symplectic forms are locally diffeomorphic to the standard symplectic form

$$\omega_0 = dx_1 \wedge dy_1 + \cdots + dx_n \wedge dy_n$$

on Euclidean space \mathbb{R}^{2n} with coordinates $(x_1, y_1, \dots, x_n, y_n)$. This makes it hard to get a handle on the global structure of symplectic manifolds. Variational techniques have been developed which allow one to investigate some global questions in Euclidean space and in manifolds such as cotangent bundles which have some linear structure: see [190, 277] and the references contained therein. But the method which applies to the widest variety of symplectic manifolds is that of J -holomorphic curves.

Here J is an almost complex structure on M which is tamed by ω . An almost complex structure is an automorphism J of the tangent bundle TM of M which

satisfies the identity $J^2 = -\mathbb{1}$. Thus J can be thought of as multiplication by i , and it makes TM into a complex vector bundle of dimension n . The form ω is said to **tame** J if

$$\omega(v, Jv) > 0$$

for all nonzero $v \in TM$. Geometrically, this means that ω restricts to a positive form on each complex line $L = \text{span}\{v, Jv\}$ in the tangent space T_xM . Given ω the set $\mathcal{J}_\tau(M, \omega)$ of almost complex structures tamed by ω is always nonempty and contractible. Note that it is very easy to construct and perturb tame almost complex structures, because they are defined by pointwise conditions. Note also that, because $\mathcal{J}_\tau(M, \omega)$ is path connected, different choices of $J \in \mathcal{J}_\tau(M, \omega)$ give rise to isomorphic complex vector bundles (TM, J) . Thus the Chern classes of these bundles are independent of the choice of J and will be denoted by $c_i(TM)$.¹

In what follows we shall only need to use the first Chern class, and what will be relevant is the value $c_1(A) := \langle c_1(TM), A \rangle$ which it takes on a homology class $A \in H_2(M)$. If A is represented by a smooth map $u : \Sigma \rightarrow M$, defined on a closed oriented 2-manifold Σ then $c_1(A) = c_1(E)$ is the first Chern number of the pullback tangent bundle $E := u^*TM \rightarrow \Sigma$. But every complex bundle E over a 2-manifold Σ decomposes as a sum of complex line bundles $E = L_1 \oplus \cdots \oplus L_n$. Correspondingly

$$c_1(E) = \sum_i c_1(L_i).$$

Since the first Chern number of a complex line bundle is just the same as its Euler number, it is often easy to calculate the $c_1(L_i)$ directly. For example, if A is the class of the sphere $S = \text{pt} \times S^2$ in $M = V \times S^2$ then it is easy to see that

$$TM|_S = TS \oplus L_2 \oplus \cdots \oplus L_n,$$

where the line bundles L_k are trivial. It follows that

$$c_1(A) = c_1(TM|_S) = c_1(TS) = \chi(S) = 2$$

where $\chi(S)$ is the Euler characteristic of S .

A smooth map $\phi : (M, J) \rightarrow (M', J')$ from one almost complex manifold to another is said to be (J, J') -**holomorphic** if and only if its derivative $d\phi(x) : T_xM \rightarrow T_{\phi(x)}M'$ is complex linear, that is

$$d\phi(x) \circ J(x) = J'(\phi(x)) \circ d\phi(x).$$

These are the Cauchy–Riemann equations, and, when (M, J) and (M', J') are both subsets of complex n -space \mathbb{C}^n , they are satisfied precisely by the holomorphic maps. An almost complex structure J is said to be **integrable** if it arises from an underlying complex structure on M . This is equivalent to saying that one can choose an atlas for M whose coordinate charts are (J, i) -holomorphic where i is the standard complex structure on \mathbb{C}^n . In this case the coordinate changes are holomorphic maps (in the usual sense) between open sets in \mathbb{C}^n . When M has dimension 2 a fundamental theorem says that all almost complex structures J on M are integrable: for a proof see Section E.4. However this is far from true in higher dimensions.

¹There is another space of almost complex structures naturally associated to (M, ω) , namely the set $\mathcal{J}(M, \omega)$ of ω -**compatible structures** defined in Section 3.1. For the present purposes one can use either space. In fact, to make our results as general as possible, we will often work with $\mathcal{J}(M, \omega)$ because this is very slightly harder: $\mathcal{J}_\tau(M, \omega)$ is open in the space of all almost complex structures, while $\mathcal{J}(M, \omega)$ is not.