


MODERN
PRACTICE
IN
TUBERCULOSIS

VOLUME
I

C. HOLMES SELLORS
AND
J. L. LIVINGSTONE



MEDICAL
PUBLICATIONS

MODERN PRACTICE IN TUBERCULOSIS

Edited by

T. HOLMES SELLORS

M.A., D.M., M.Ch., F.R.C.S.

THORACIC SURGEON, MIDDLESEX HOSPITAL;
SURGEON, LONDON CHEST AND HAREFIELD HOSPITALS;
TEACHER OF SURGERY, MIDDLESEX HOSPITAL MEDICAL
SCHOOL AND THE INSTITUTE OF DISEASES OF THE CHEST,
UNIVERSITY OF LONDON

and

J. L. LIVINGSTONE

M.D., F.R.C.P.

PHYSICIAN, KING'S COLLEGE HOSPITAL AND BROMPTON
HOSPITAL, LONDON; FELLOW OF KING'S COLLEGE,
UNIVERSITY OF LONDON; TEACHER OF MEDICINE,
KING'S COLLEGE HOSPITAL MEDICAL SCHOOL AND
THE INSTITUTE OF DISEASES OF THE CHEST, UNIVERSITY
OF LONDON

[In two volumes]

VOL. 1

LONDON
BUTTERWORTH & CO. (PUBLISHERS) LTD.
BELL YARD, TEMPLE BAR
1952

BUTTERWORTHS MEDICAL PUBLICATIONS

MODERN PRACTICE SERIES

Under the General Editorship of

THE RT. HON. LORD HORDER, G.C.V.O., M.D., F.R.C.P.

The following titles have been chosen for the above-mentioned series:

ANAESTHESIA—Edited by FRANKIS T. EVANS, M.B., B.S., F.F.A.R.C.S., D.A. (*Published*)

DERMATOLOGY—Edited by G. B. MITCHELL-HEGGS, O.B.E., M.D., F.R.C.P. (*Published*)

OPHTHALMOLOGY—Edited by H. B. STALLARD, M.B.E., M.D., F.R.C.S. (*Published*)

PSYCHOLOGICAL MEDICINE—Edited by J. R. REES, C.B.E., M.D., F.R.C.P., D.P.H. (*Published*)

INFECTIOUS FEVERS (*in two volumes*)—Edited by H. STANLEY BANKS, M.A., M.D., F.R.C.P., D.P.H. (*Published*)

(Further titles are in course of preparation)

MODERN PRACTICE
IN
TUBERCULOSIS

VOL. 1

AFRICA: BUTTERWORTH & CO. (AFRICA), LTD.
DURBAN: 1 LINCOLN'S COURT, MASONIC GROVE

AUSTRALIA: BUTTERWORTH & CO. (AUSTRALIA), LTD.
SYDNEY: 8 O'CONNELL STREET
MELBOURNE: 430 BOURKE STREET
BRISBANE: 240 QUEEN STREET

CANADA: BUTTERWORTH & CO. (CANADA), LTD.
TORONTO: 1367 DANFORTH AVENUE

NEW ZEALAND: BUTTERWORTH & CO. (AUSTRALIA), LTD.
WELLINGTON: 49/51 BALLANCE STREET
AUCKLAND: 35 HIGH STREET

Distributors in U.S.A.
PAUL B. HOEBER INC.
49 EAST 33RD STREET
NEW YORK 16, N.Y.

LIST OF ILLUSTRATIONS

VOL. 1

	PAGE
FIG. 1.—Proportion of bodies at different ages showing tuberculous lesions at autopsy — — — — —	29
FIG. 2.—A tuberculin survey of the whole population (about 50,000) in the city of Orebro, Sweden, in 1944 — — — — —	29
FIG. 3.—Death rate per 100,000 from respiratory tuberculosis in various countries, 1920, 1925, 1930, 1935 — — — — —	41
FIG. 4.—Tuberculosis death rate per 100,000 living, at each age, in the United States of America, 1940 — — — — —	41
FIG. 5.—Tuberculosis attack rates in children of index cases by age, according to character of index cases — — — — —	42
FIG. 6.—Cumulative probability of developing tuberculosis for children of index cases by age, according to character of index cases — — — — —	42
FIG. 7.—Massachusetts death rates from tuberculosis per 100,000 in males, by age, 1880, 1910, 1930 — — — — —	43
FIG. 8.—Massachusetts death rates from tuberculosis per 100,000 in males, by age, in successive 10-year cohorts — — — — —	43
FIG. 9.—Death rates per 100,000 from tuberculosis, by age (United States of America, 1940) — — — — —	44
FIG. 10.—Tuberculosis mortality by sex and age per 100,000 living, England and Wales, 1939 — — — — —	45
FIG. 11.—Tuberculosis death rate and level of real wages, England and Wales, 1850–1940 (semi-logarithmic scale) — — — — —	47
FIG. 12.—Tuberculosis mortality per 100,000 adult males in three social classes England and Wales — — — — —	47
FIG. 13.—Standardized tuberculosis death rates per 100,000 occupied males aged 15–64 years in 10 states of the United States of America, 1930 — — — — —	47
FIG. 14.—Tuberculosis mortality, white population, by economic status, Buffalo, N.Y., 1939–41 — — — — —	47
FIG. 15.—Mean annual mortality rates from tuberculosis per 100,000 persons, classified according to family income, Hamburg, Germany, 1896–1900 — — — — —	48
FIG. 16.—Death rate from pulmonary tuberculosis by occupation, United States of America, 1930 (males) — — — — —	48
FIG. 17.—The degree of positiveness. The visual grading of sputa is not easy. When visual grading is compared with precise field counts, it becomes obvious that the visual gradings show a considerable degree of overlapping — — — — —	57
FIG. 18.—Sputum conversion curve. This diagram summarizes the results of a mathematical analysis of the conditions holding for direct smear examinations. It shows, as would be expected, that the process of transition from a positive to a negative film, or <i>vice-versa</i> , is a gradual one — — — — —	58
FIG. 19.—Distribution of colony counts in graded sputa. This diagram is constructed to show the way in which the various sputum classes grade one into the other — — — — —	59

LIST OF ILLUSTRATIONS

	PAGE
FIG. 20.—Discrepancies of six weeks. Where the colony count is low, it is common to find, for a pair of simultaneously sown tubes, that one is negative and the other positive — — — — —	61
FIG. 21.—Seasonal variations. A recent observation, included to show the care needed when quantitative bacteriology is utilized. Analysis of long-term records shows periodic variations in the proportion of positives found at different times of the year, these repeating themselves cyclically in successive years — —	61
FIG. 22.—Distribution of 1,001 colony counts. (This graph is equivalent to the lowest curve in Fig. 23.) — — — — —	63
FIG. 23.—Sputum tests—diagrammatic summary. The proportion of cases providing material suitable for examination by the various microscopic and culture techniques is indicated in the spectrum — — — — —	64
FIG. 24.—Examples of sputum conversion curves — — — — —	66
FIG. 25.—Estimating bacillary concentrations from direct smear results — —	70
FIG. 26.—Correlation of sputum bacteriology with blood sedimentation — —	74
FIG. 27.—Eosinophilia counts following thoracoplasty — — — — —	76
FIG. 28.—Healing primary complex in bronchiectatic lower lobe. Primary focus (F). Glandular component (G) — — — — —	80
FIG. 29.—Part of the wall of a small bronchus replaced by a tubercle — — —	80
FIG. 30.—Progressive primary complex — — — — —	81
FIG. 31.—(a) Progressive primary complex. Hilum of left lung showing caseating glands. (b) Lateral surface of left lung showing area of caseating pneumonia —	82
FIG. 32.—Lateral view of left lung showing confluent caseous pneumonia in the upper lobe and extensive broncho-pneumonia in lower lobe — — — —	85
FIG. 33.—Lower lobe showing a cavitating primary focus (F), caseating gland (G) and early bronchogenic spread — — — — —	85
FIG. 34.—Hilar surface of right lung dissected to show a calcified broncho-pulmonary gland (G) at a stenosis of the upper lobe bronchus — — —	86
FIG. 35.—Lateral surface of right lung cut to show an acute cavity (C) in the middle lobe and bronchiectasis in the collapsed upper lobe — — — —	86
FIG. 36.—Acute cavity with lung stones. Calcified focus (F) above cavity — —	87
FIG. 37.—(a) Tension cavity in the right lower lobe. (b) Section of bronchus draining tension cavity showing a sub-epithelial tubercle — — — — —	90
FIG. 38.—Right lung with a blocked cavity (C) in the lower lobe — — — —	91
FIG. 39.—Left lung showing a chronic cavity in the upper lobe — — — —	91
FIG. 40.—Lateral view of the left lung showing an empyema over the lingula —	94
FIG. 41.—Same case as Fig. 40. Close up of left lung showing cavity (C), fistula (F) and empyema (E) — — — — —	95
FIG. 42.—Right lung showing stenosis of the lower part of the main bronchus and bronchiectasis of the middle and lower lobes. Caseating gland (G) has been dissected away from the main bronchus and is now lying free, some distance away	97
FIG. 43.—Low-power magnification of the area of stenosis (S) in the lower part of the main bronchus — — — — —	97
FIG. 44.—Collapsed left lung showing tuberculous bronchiectasis. Bronchus (B) was mistaken in the film for a cavity — — — — —	99
FIG. 45.—A dissection of the right bronchial tree to show stenosis of the upper lobe bronchus (U), the lower lobe bronchus (L) and bronchiectasis in the middle lobe — — — — —	99

LIST OF ILLUSTRATIONS

	PAGE
FIG. 46.—Lateral view of the right lung showing bronchi (B) filled with caseous debris in the collapsed upper lobe (U) and the collapsed middle (M) and lower lobes (L). (a) Photograph of specimen; (b) Key to areas — — — — —	100
FIG. 47.—Lateral view of left lung showing total bronchiectasis following healed tuberculous stenosis of the left main bronchus — — — — —	101
FIG. 48.—On diagnosis; an acute lower-lobe lesion with distension cavity — —	148
FIG. 49.—Same case as in Fig. 48, after induction of pneumoperitoneum. Atelectasis of lower lobe with persisting cavity — — — — —	149
FIG. 50.—Same case as in Figs. 48 and 49 after partial re-expansion of lower lobe and administration of streptomycin. Cavity not seen; sputum cultures negative	149
FIG. 51.—Rapid lung destruction following bronchial obstruction and atelectasis	154
FIG. 52.—Premature pneumothorax (without bed rest) in an acute active lesion followed by atelectasis and empyema — — — — —	154
FIG. 53.—An acute lesion treated by 7 months' bed rest augmented by pneumoperitoneum and right phrenic crush. The subsequent artificial pneumothorax was uncomplicated by atelectasis — — — — —	154
FIG. 54.—Extensive active lesions treated by streptomycin. Uncomplicated artificial pneumothorax subsequently induced on right side. No collapse therapy necessary on left side — — — — —	154
FIG. 55.—A free pneumothorax but massive collapse of upper lobe. Subsequent contralateral spread. A common occurrence in such conditions— — —	154
FIG. 56.—Atelectasis allowed to persist with production of empyema — — —	154
FIG. 57.—Atelectatic lung re-aerated with subsequent establishment of a "clean" artificial pneumothorax — — — — —	154
FIG. 58.—A dangerous situation following adhesion section. Air at once removed with subsequent "clean" artificial pneumothorax — — — — —	154
FIG. 59.—A retrogressing lesion reactivated by pneumothorax which produced bronchial stenosis — — — — —	154
FIG. 60.—An inspissated cavity reopened and artificial pneumothorax was induced. The bronchus became kinked, but by trial and error a satisfactory cavity-bronchus relationship was found. The lung was alternately collapsed and re-expanded and also pneumoperitoneum was induced. The cavity was ultimately persuaded to close by these methods — — — — —	154
FIG. 61.—Phrenic nerve paralysis and pneumoperitoneum as a useful preliminary to pneumothorax — — — — —	155
FIG. 62.—Phrenic nerve paralysis and pneumoperitoneum as a useful adjunct to artificial pneumothorax — — — — —	155
FIG. 63.—Dropping the lung by artificial pneumothorax caused bronchial stenosis, but diaphragmatic rise was more effective treatment — — — — —	155
FIG. 64.—Increase in effectiveness of bronchial check valve by diaphragmatic rise with distension of cavity — — — — —	155
FIG. 65.—A case of tracheobronchitis in which diaphragmatic rise brought about complete atelectasis — — — — —	155
FIG. 66.—A shift of the mediastinum to the right when the left lung was over-collapsed was accompanied by distension of the cavities in the contralateral lung. Cavities disappeared on return of mediastinum to normal position and induction of pneumoperitoneum — — — — —	155

LIST OF ILLUSTRATIONS

	PAGE
FIG. 67.—Mediastinal shift again produced cavity distension in spite of pneumoperitoneum, but the cavities again disappeared with re-expansion of left lung —	155
FIG. 68.—Lipiodol introduced into a cavity through a drainage catheter shows the absence of bronchial drainage. Following first and second stage thoracoplasty bronchial drainage became established and the cavity closed — — — —	155
FIG. 69.—A cavity which became distended after thoracoplasty because the drainage bronchi failed to open. This cavity never closed although the thoracoplasty was completed — — — — — — — — — —	155
FIG. 70.—Lipiodol introduced into a cavity through Monaldi drainage tube. The bronchi are occluded and the Lipiodol is retained in the cavity — — —	157
FIG. 71.—Same case as in Fig. 70, immediately after thoracoplasty. The cavity is already reduced to a slit and the Lipiodol passes freely into the open bronchi —	157
FIG. 72.—A cavity persists after a 3-stage thoracoplasty. Lipiodol introduced into the cavity shows that the bronchi have not been freely opened — — — —	158
FIG. 73.—Same case as in Fig. 72, immediately after a revision thoracoplasty. Lipiodol introduced into the residual cavity shows that it has been reduced to a mere slit and that free bronchial drainage has been established— — — —	158
FIG. 74.—Male, aged 35 years. Minimal fibro-calcified lesion. History of cough lasting 3 months, 2 years prior to examination. Alive and well 28 years later	238
FIG. 75.—Female, aged 22 years. Minimal mid-zone lesions discovered on routine x-ray examination 2½ years after primary pleural effusion on opposite side, for which she had 7 months' in-patient treatment — — — — —	238
FIG. 76.—(a) Female, aged 26 years. Minimal lesion at right apex. Symptomless contact case. (b) Same case but taken 3 months later, showing spread and excavation — — — — — — — — — —	239
FIG. 77.—(a) Female, aged 13 years. Contact case showing minimal lesion at right apex. No symptoms. (b) Same patient. Condition remained unchanged until 6 years later, when extension of disease was noted for the first time — —	250
FIG. 78.—Processing unit with running water rinse — — — — —	254
FIG. 79.—Processing unit with spray rinse — — — — —	255
FIG. 80.—Diagram illustrating the broncho-pulmonary nomenclature — — —	256
FIG. 81.—Tomographs of the right upper zone, taken with 15 degree arc of swing. Cavity not shown in the 7 or 9 cms cut — — — — —	260
FIG. 82.—(a and b) Same patient as in Fig. 81. Tomographs with 30 degree arc of swing show cavity in the 8 cms cut (b) but not in the 7 cms cut (a) — — —	261
FIG. 83.—Tomograph of a cavity in the midst of an area of opacity. No bronchus nearby. Specimen showed cavity was completely walled off by fibrous tissue —	264
FIG. 84.—Atelectasis of the left lung from occlusion of the left main bronchus—the end-result of a localized tuberculous bronchitis. The clear area in the upper part is the right lung, herniated across — — — — —	267
FIG. 85.—(a) Atelectasis of the left upper lobe from tuberculous bronchitis. (b) Lateral view. Arrows point to the tongue-like shadow of the collapsed lobe	268
FIG. 86.—(a) Localized tuberculous bronchitis with distal bronchiectasis. Patchy clouding in the right posterior basal segment. (b) Lateral view. Arrow marks the site of the affected segment — — — — —	269

LIST OF ILLUSTRATIONS

	PAGE
FIG. 87.—(a) Same case as in Fig. 86: a bronchogram. Arrow marks filling defect in the opaque oil shadow. Bronchiectasis distal to this. (b) A lateral view bronchogram. Arrow marks the endobronchial mass projecting into the oil shadow —	270
FIG. 88.—Pneumothorax with circular translucency in the airless lung (marked by arrow). This is not a tuberculous cavity but a grossly dilated bronchus distal to an area of tuberculous bronchitis and stenosis — — — — —	272
FIG. 89.—A 3 cm circular shadow in the right lung ("tuberculoma") — — —	272
FIG. 90.—(a) A 3 cm circular shadow in the left upper zone. (b) The same case some months later. There is now an enormous cavity in the left upper zone. There is also a pleural effusion — — — — —	273
FIG. 91.—Neoplasm of the right upper lobe bronchus. The growth is invisible on the plain skiagram, but the distal inflammatory changes are clearly seen —	274
FIG. 92.—(a) Primary tuberculosis in a woman aged 41 years. Skiagram shows 3 cm area of clouding spreading out from the left hilum. (b) The lateral view shows consolidation of the anterior segment (lower limit marked by arrow) — — —	275
FIG. 93.—Tuberculous lesions in the right upper lobe. There is a large cavity, with fluid level in it, and surrounding consolidation below the clavicle. The apical translucency is actually a bulla — — — — —	277
FIG. 94.—(a) Constrictive pericarditis. Kymograph shows a small heart with diminished movement of both borders. (b) After surgical stripping of the thickened pericardium. The heart is larger, and vigorous movement is shown — — —	278
FIG. 95.—Twenty-four hours after first stage thoracoplasty, during which the pleura was accidentally opened. Skiagram shows collapsed airless lung. Arrow points to the fluid level in the pleura. There is a small fluid level with air above it in the extrapleural space — — — — —	280
FIG. 96.—Thoracoplasty with introduction of hollow Lucite balls (tomograph). The regular circle could not be mistaken for a cavity — — — — —	285
FIG. 97.—Six months after thoracoplasty on the left side with introduction of 18 solid Lucite balls. The appearances are the same as those of an extrapleural collection of fluid. On the right side there has been an artificial pneumothorax and rib resection for the decortication of a complicating empyema — — —	285
FIG. 98.—Tuberculosis of the small intestine. Skiagram taken 4 hours after a 10 oz. barium meal, showing areas of narrowing and irregularity in the small intestine, with some dilated loops high up on the left side — — — —	287
FIG. 99.—Tuberculosis of the colon. Skiagram taken 24 hours after a barium meal, showing a narrow area in the pelvic colon but no hypermotility — — —	287
FIG. 100.—Tuberculous strictures of the small intestine. Skiagram taken 1 hour after a 10 oz. barium meal, showing a grossly dilated coil of small intestine on the left side — — — — —	288
FIG. 101.—Tuberculosis of the ascending colon, with narrowing and indentation simulating a neoplasm. There is a fistula leading to the right kidney. The transverse colon is also affected — — — — —	289
FIG. 102.—Tuberculosis of the ileo-caecal area. Skiagram taken 6 hours after a barium meal, showing barium in the small intestine in the pelvis and in the transverse colon. There is poor filling of the irritable affected segment — —	289
FIG. 103.—Marshalling examinees for rapid fluorography — — — — —	294
FIG. 104.—Inserting the record card so that the appropriate number is photographed with the screen image of the examinee's chest — — — — —	294

LIST OF ILLUSTRATIONS

VOL. 2

PAGE

FIG. 1.—Primary infection with Mantoux conversion known to have taken place between December 1936 and April 1937. At time of Mantoux conversion x-ray was normal. (a) Showing enlarged hilar glands with partial collapse of middle lobe three months after Mantoux conversion. (b) Increased collapse of middle lobe eight to twelve months after primary infection. (c) Bronchogram showing blocking of middle lobe bronchus	5
FIG. 2.—Temperature chart of girl aged 3 years with gross hilar adenitis following primary tuberculous infection	6
FIG. 3.—(a) Primary tuberculous lesion with enlarged right paratracheal glands and collapse of middle lobe. (b) Healing lesion with calcified primary focus in right upper lobe and calcified lesion in glands at right hilum	8
FIG. 4.—Calcifying primary lesion of right lower lobe, with resolving localized pleural effusion	9
FIG. 5.—(a) Pleural effusion. (b) Showing effusion cleared. Enlarged glands left hilum and early primary lesion left third intercostal space	10
FIG. 6.—Antero-posterior x-ray of an atelectatic right upper lobe in a child	22
FIG. 7.—The same case as in Fig. 6: right upper lobe has re-aerated, not re-expanded. Coarse striations of bronchiectasis present in collapsed lobe	23
FIG. 8.—Bronchograph confirms a saccular bronchiectasis	23
FIG. 9.—Collapse of the right middle lobe	24
FIG. 10.—Film taken in inspiration. Obstructive emphysema due to granulations in right descending bronchus impinging on the orifice of the upper lobe bronchus	25
FIG. 11.—Film in expiration. Displacement of mediastinum more marked	25
FIG. 12.—X-ray of a boy with a resolved primary tuberculosis consolidation of the right upper and middle lobes	26
FIG. 13.—Same case as in Fig. 12. Tomograph showing collapse of right antero-lateral sector	26
FIG. 14.—A large tension cavity in the right posterior basic sector. A right artificial pneumothorax has just been induced	31
FIG. 15.—Phrenic paralysis is followed by atelectasis of the right lower lobe, the cavity persists	31
FIG. 16.—Lateral view of Fig. 15	32
FIG. 17.—Re-expansion of lower lobe and closure of cavity after recovery of phrenic nerve and freeing of lung	32
FIG. 18.—Tension cavity in lower lobe on the right. Extensive bilateral disease and bilateral artificial pneumothorax	33
FIG. 19.—The cavity enlarges in spite of manipulation of right artificial pneumothorax pressures and phrenic paralysis	33
FIG. 20.—Cavity closes after a course of streptomycin (confirmed by tomography)	34
FIG. 21.—Massive collapse of the left lung. Patient has a hectic pyrexia	40
FIG. 22.—Same case as Fig. 21. Bronchography revealed bronchiectasis of the middle and posterior basic sectors	40
FIG. 23.—Case 1. Male, aged 18 years. (a) Acute exudative disease throughout left upper lobe, with multiple small cavities. Left phrenic crush and pneumoperitoneum induced. (b) Pneumoperitoneum; but left diaphragm at level of 4th space, which is not high enough. Cavities still visible	52

LIST OF ILLUSTRATIONS

PAGE

FIG. 23 <i>cont.</i> —(c) Marked clearing of infiltration. Left diaphragm at proper level.	
(d) Heavy left hilar shadows with raised diaphragm: no calcification — — —	53
FIG. 24.—Case 2. Male, aged 18 years. (a) Routine radiograph in April 1946 shows scattered soft infiltration in left upper and mid zones with two cavities about 2 centimetres in diameter in postero-lateral segment of upper lobe. Minimal infiltration right apex. (b) Clearing of infiltration and apparent closure of cavities at left apex. Left diaphragm at level of 5th rib — — — — —	54
FIG. 24 <i>cont.</i> —(c) Tomograph at 5 centimetres level confirming closure of cavities at left apex. Note well-marked Assman's focus at right apex. (d) Radiograph showing cavitation at right apex. Calcified foci at left upper zone and left diaphragm now at level of 3rd space — — — — —	55
FIG. 24 <i>cont.</i> —(e) Radiograph shows right thoracoplasty. Left lung apparently healed: calcified foci in upper zone — — — — —	56
FIG. 25.—Case 3. Male, aged 23 years. (a) Infiltration in upper and middle zones, with cavity at right apex. Extensive infiltration in middle and lower lobes — —	56
FIG. 25 <i>cont.</i> —(b) Pneumoperitoneum with diaphragm to 3rd space. Tomograph shows cavity still persists at right apex. (c) Right thoracoplasty with large solid foci in middle and lower zones — — — — —	57
FIG. 26.—Case 4. Female, aged 28 years. Scattered nodular infiltration throughout left lung and in upper and middle zones of right lung. Large cavity (5.5 × 7 centimetres) in left upper lobe. (b) More penetrated film to show cavity — — —	58
FIG. 26 <i>cont.</i> —(c) Pneumoperitoneum with diaphragm to 3rd rib. Left upper lobe collapsed: cavity reduced to 2 centimetres. (d) Scattered calcified foci throughout both lungs. Left diaphragm to 5th rib — — — — —	59
FIG. 27.—Case 5. Female, aged 23 years. (a) Cavity 5 centimetres at right apex, with fluid level: infiltration in middle zone. Infiltration left lower zone with cavity, 2 centimetres; hard infiltration left apex. (b) Tomograph at 5 centimetres confirms cavitation at right upper zone but no cavity at left apex — — —	60
FIG. 27 <i>cont.</i> —(c) Tomograph at 5 centimetres shows no cavity at right apex. (d) Pneumoperitoneum with partial collapse of right upper lobe, which appears fairly clear — — — — —	61
FIG. 28.—Case 6. Female, aged 20 years. (a) Radiograph shows exudative disease with cavitation (2.2 centimetres in diameter) in apical segment of left lower lobe: minimal infiltration in right mid-zone. (b) Satisfactory pneumoperitoneum —	62
FIG. 28 <i>cont.</i> —(c) Tomograph at 3 centimetres shows round solid focus in position of original cavity. (d) Further infiltration in left lower lobe, although diaphragm maintained to 4th rib — — — — —	63
FIG. 28 <i>cont.</i> —(e) Tomograph at 3 centimetres shows blocked cavity has reopened. (f) 11 months after lobectomy. Left diaphragm still high with gastric air bubble	64
FIG. 29.—Case 7. Male, aged 29 years. (a) Cavity 3 centimetres in diameter at left apex with smaller cavity in first space and infiltration in left upper and mid zones	66
FIG. 29 <i>cont.</i> —(b) Pneumoperitoneum induced, left diaphragm to 3rd rib. Cavitation no longer visible. (c) Left costophrenic angle is obliterated, with linear opacity and emphysema at the base. Calcified foci in middle and upper zones —	67
FIG. 30.—Artificial pneumothorax. Diagrams of radiological appearances preceding onset of empyema — — — — —	70

LIST OF ILLUSTRATIONS

PAGE

FIG. 31.—Male, aged 23 years. (a) Extensive soft infiltration in upper and middle zones of left lung with two cavities 2 centimetres and 2.5 centimetres at left apex. Scattered soft infiltration in right upper zone. (b) Small extrapleural space: calcified foci in left middle zone and right upper zone — — — — —	73
FIG. 31 cont.—(c) Left oleothorax after the introduction of 500 millilitres of olive oil. (d) Left oleothorax: calcified foci in both lungs — — — — —	74
FIG. 32.—Plombage with Lucite spheres in a case of bronchiectasis where patient unfit for pneumonectomy. (a) Extensive bronchiectasis left lung (postero-anterior view). (b) Extensive bronchiectasis left lung (lateral view) — — — — —	91
FIG. 32 cont.—(c) After plombage (postero-anterior view). (d) After plombage (lateral view) — — — — —	92
FIG. 32 cont.—(e) Tomograph showing spheres in position (65 in number). (f) Condition 18 months after operation — — — — —	93
FIG. 33.—(a) Large pulmonary cavity at apex with contra-selective A.P. Almost complete collapse of lung. (b) After operation, E.P. and A.P. space very extensive, but cavity still persists — — — — —	94
FIG. 33 cont.—(c) A.P. and E.P. space smaller. Pulmonary cavity has disappeared. (d) A.P. and E.P. space smaller, lung coming out. Pulmonary cavity has disappeared — — — — —	95
FIG. 33 cont.—(e) Lung still creeping out. (f) Cap of thickened pleura has replaced the E.P. space. The pulmonary cavity remains healed — — — — —	96
FIG. 34.—Pulmonary cavities both apices. (a) Generalized bilateral disease. (b) Cavity left apex, spreading lesion right apex — — — — —	97
FIG. 34 cont.—(c) Tomograph shows bilateral apical cavities. (d) A.P. right side. Adhesions holding out apex — — — — —	98
FIG. 34 cont.—(e) Division of adhesions and pneumoperitoneum controlling right side. (f) After E.P. on left. (Note pulmonary cavity not yet obliterated) — — — — —	99
FIG. 34 cont.—(g) Control of pulmonary cavities on both sides. (h) Thoracoplasty left. Lung restored on right — — — — —	100
FIG. 35.—X-ray films showing active disease controlled by E.P. Extensive <i>Fruh infiltrat</i> left, unstable lung right. (b) Left E.P. — — — — —	101
FIG. 35 cont.—(c) E.P. replaced by firm "cap". Disease on left controlled — — — — —	102
FIG. 36.—(a) Apical cavity in lung with adhesive pleurae — — — — —	102
FIG. 36 cont.—(b) E.P. immediately after operation. Pleural detachment to 8th rib laterally. Retracted apex maintains general form. (c) E.P. a few weeks after operation. Lower part of detachment reattached. Apex of lung depressed. (d) Poor pneumothorax, with adherent apex. (e) Combined intrapleural and extrapleural pneumothorax showing "shelf". (f) Beginning of décollement. (g) Décollement continues, apex of lung peeled down. Creation of E.P. space — — — — —	103
FIG. 37.—Demonstrating usual progress of E.P. during 10 years. (a) Before operation—cavity in right apex. (b) Five years after E.P.—extensive "space" — — — — —	108
FIG. 37 cont.—(c) Ten years later—obliteration of E.P. space and replacement by thick "cap" of fibrous tissue. Cavity healed — — — — —	109
FIG. 38.—(a) Shows bilateral apical disease. Cavity right side and small apical cavity on left. (b) Shows right sided cavity — — — — —	120
FIG. 38 cont.—(c) Shows right thoracoplasty with cavity opposite anterior end of 2nd rib on left side. (d) Shows condition after bilateral thoracoplasty — — — — —	121

LIST OF ILLUSTRATIONS

	PAGE
FIG. 39.—(a) Shows apical cavitation after introduction of Monaldi tube with left artificial pneumothorax — — — — —	122
FIG. 39 <i>cont.</i> —(b) Shows condition after injection of Lipiodol into cavity before thoracoplasty. (c) Shows condition after thoracoplasty with left lung re-expanded. Some fibrosis and emphysema at left apex — — — — —	123
FIG. 40.—(a) Shows infiltration at right apex. Left lung normal — — — — —	124
FIG. 40 <i>cont.</i> —(b) Tomograph shows cavity at right apex. (c) Shows condition after 5-rib thoracoplasty. No evidence of cavitation. Left lung normal — — — — —	125
FIG. 41.—(a) Shows large cavity left apex with scattered infiltration throughout the whole of the left lung. Right side shows artificial pneumothorax — — — — —	126
FIG. 41 <i>cont.</i> —(b) Shows condition after 1st stage thoracoplasty. Cavity still obvious. (c) Shows condition after thoracoplasty. Right lung re-expanded. Fibrosis in middle zone with few scattered calcifications. Left side shows apex well collapsed. Scattered calcifications in lower zone — — — — —	127
FIG. 42.—(a) Shows cavity in left upper lobe, also cavity in apical segment of the lower lobe which is not clearly defined. Small patch of infiltration in right mid zone. Pneumoperitoneum with elevation of left leaf of diaphragm — — — — —	128
FIG. 42 <i>cont.</i> —(b) Tomograph confirms presence of left cavitation. (c) Shows condition after left thoracoplasty — — — — —	129
FIG. 43.—(a) Shows right apical cavity and left artificial pneumothorax with scattered calcified infiltration throughout both lungs. (b) Condition after right thoracoplasty. Left artificial pneumothorax being maintained — — — — —	130
FIG. 44.—Giant right apical cavity with fluid level. The film shows the appearance after a limited anterior thoracoplasty — — — — —	144
FIG. 45.—Excavation of almost all the right upper lobe following consolidation-collapse type of lesion. Right phrenicectomy and pneumoperitoneum were ineffective. (a) Endocavity drainage (Monaldi) established after a limited anterior thoracoplasty. (b) A posterior thoracoplasty with limited apicolysis followed and the tube was removed when the cavity had completely closed. Three years later the cavity has remained closed (c) and sputum is negative — — — — —	144
FIG. 46.—Giant dorsal (apex of lower) lobe cavity on the left (a). Treated by cavernostomy using a posterior approach. Antibiotics helped the relatively quick closure which has been permanent (b) — — — — —	146
FIG. 47.—Cavity in apex of the <i>right</i> lower lobe treated by cavernostomy. A left upper thoracoplasty had been performed some years previously (a). A posterior approach was used for a skin flap drainage of the cavity (b) — — — — —	147
FIG. 48.—Cavity at the apex of the right lower lobe which was not permanently controlled by AP or phrenicectomy (a) — — — — —	152
FIG. 48 <i>cont.</i> —The posterior situation of the cavity is seen in the lateral view (b). Segmental resection of the apex of the lower lobe gave a good result (c) — — — — —	153
FIG. 49.—Pneumonectomy for bronchostenosis and extensive disease in the left lung. A preliminary pneumoperitoneum was induced (a) and pneumonectomy performed with difficulty. Three weeks later a limited upper thoracoplasty was carried out for partial obliteration of the dead space. Note that the mediastinum has become central (b) — — — — —	155
FIG. 50.—Failure of thoracoplasty to close left apical cavities. In spite of an extensive apicolysis the lung "crept" out and there was considerable activation of disease in this area (a). After mobilization of the apex it was found possible to excise the diseased lobe (b) — — — — —	156

LIST OF ILLUSTRATIONS

PAGE

FIG. 51.—Pneumonectomy for “destroyed” lung. Active ulcerating disease in right lung uncontrolled by rest, antibiotics and pneumoperitoneum (<i>a</i>). AP was impracticable. Excision of the lung was followed by immediate improvement and a thoracoplasty was performed later (<i>b</i>)	158
FIG. 52.—Right upper lobe cavitation, also cavity in left mid-zone region (<i>a</i>). Marked endobronchitis treated by streptomycin and PAS resulting in a stricture in right upper lobe bronchus. Right phrenicectomy was added later and simultaneously the left cavity closed (<i>b</i>)	159
FIG. 53.—Extensive disease in left lung showing steady progression in addition to gross endobronchitis. Note elevation of left diaphragm (<i>a</i>). Following pneumonectomy there was remarkable improvement and the patient has remained well for 3 years without a thoracoplasty (<i>b</i>)	161
FIG. 54.—Female aged 16 years. (<i>a</i>) Film shows effusion left side and evidence of a haemic dissemination in the right lung. (<i>b</i>) Shows absorption of fluid and diffuse miliary spread of tuberculous foci throughout both lungs	166
FIG. 55.—Male aged 20 years. (<i>a</i>) Shows bilateral pleural effusion. (<i>b</i>) Further examination 17 years later, shows multiple calcified foci in the lung fields having the distribution of a haemic dissemination	168
FIG. 56.—Tension cavity: adhesion section: empyema: death	192
FIG. 57.—Tuberculous empyema: amyloid disease: pleurectomy: recovery	194
FIG. 58.—Tuberculous empyema: pleurectomy: recovery	197
FIG. 59.—Tuberculous empyema: drainage: thoracoplasty: recovery	199
FIG. 60.—Lung cavity: spontaneous tuberculous pyopneumothorax: pleuro-pneumonectomy: recovery	201
FIG. 61.—Bilateral pulmonary tuberculosis: bilateral artificial pneumothorax: right spontaneous pneumothorax: left tuberculous empyema: pleurectomy: recovery	203
FIG. 62.—Bilateral pulmonary tuberculosis: contra-selective artificial pneumothorax (left)—mixed infection empyema—penicillin sterilization: right A.P.—super-added spontaneous pneumothorax: death	206
FIG. 63.—Artificial pneumothorax: tuberculous empyema: aspirations and chemotherapy: thoracoplasty	207
FIG. 64.—Adhesion section: empyema with bronchopleural fistula and pleuro-cutaneous sinus: pleuropneumonectomy	209
FIG. 65.—Adhesion section: atelectasis and tuberculous empyema: aspiration and chemotherapy: bronchial fistula: thoracoplasty	211
FIG. 66.—Lobectomy to prevent pyothorax	213
FIG. 67.—Mixed infection pyothorax: actinomycosis: sinuses: sterilization: radical toilet: hypoproteinaemia and nephrosis: recovery	215
FIG. 68.—Artificial pneumothorax: endobronchial tuberculosis: pyothorax: phrenic crush: thoracoplasty	218
FIG. 69.—Inadequate artificial pneumothorax: tuberculous effusion: aspiration: azochloramide lavage: sterilization: re-expansion	220
FIG. 70.—Pleural effusion: tuberculous empyema: aspiration: residual encysted collection: late secondary infection and bronchial fistula: aspiration: re-expansion: recovery	222
FIG. 71.—Aspiration techniques	225

LIST OF ILLUSTRATIONS

	PAGE
FIG. 72.—Adhesion section: atelectasis: tuberculous empyema: aspiration: phrenic crush: re-expansion — — — — —	225
FIG. 73.—Post-lobectomy basal empyema: drainage: pneumoperitoneum: radical toilet (unroofing): healing — — — — —	227
FIG. 74.—Pleurectomy for tuberculous empyema—operative technique — —	229
FIG. 75.—André Maurer's articulated rib spreader — — — — —	232
FIG. 76.—Pyothorax and bronchial fistula after pneumothorax: pleurectomy and segmental resection: recovery — — — — —	233
FIG. 77.—Artificial pneumothorax: ruptured tension cavity: empyema: upper thoracoplasty: basal pleurectomy: recovery — — — — —	235
FIG. 78.—Contra-selective artificial pneumothorax: tension cavity: pyothorax: thoracoplasty without drainage: healing — — — — —	238
FIG. 79.—Lobar atelectasis: empyema: basal re-expansion after sterilization and aspiration: apical thoracoplasty — — — — —	240
FIG. 80.—Oleothorax: late bronchial perforation: pyopneumothorax: thoracoplasty: re-accumulation of empyema: drainage and radical toilet: recovery —	242
FIG. 81.—Tuberculous pyothorax: thoracoplasty without drainage: encysted residual collection: delayed bronchopleural fistula — — — — —	246
FIG. 82.—Perforation externally of undrained residual empyema under thoracoplasty	248
FIG. 83.—Adhesion section: pyopneumothorax: abdominal sinuses: thoracoplasty: radical toilet: streptomycin: recovery — — — — —	250
FIG. 84.—Upper lobectomy for tuberculous cavity: bronchopleural fistula: secondary lower lobectomy and total pleurectomy: recovery — — — —	254
FIG. 85.—Operative technique of radical toilet for residual empyema cavity — —	256
FIG. 86.—Healing after radical toilet for residual empyema cavity. Healing after the "saucerization" type of unroofing operation for residual empyema space depicted in Fig. 85 — — — — —	257
FIG. 87.—Tuberculous infection of chest wall after thoracoplasty — — —	258
FIG. 88.—Mixed infection pyothorax: cellulitis of chest wall: drainage and multiple incisions: thoracoplasty: healing — — — — —	259
FIG. 89.—Immobilization of the neck by a head or carapace splint — — —	278
FIG. 90.—Skiagrams of the neck showing calcified lymph nodes — — —	280
FIG. 91.—Skiagram of calcified lymph nodes removed at operation — — —	281
FIG. 92.—Premature fusion of the left knee epiphyses in tuberculosis of the left hip of a child after 4 years fixation — — — — —	305
FIG. 93.—A lesion of the lumbar spine considered by experienced surgeons to be pyogenic. The misleading feature was the stout buttress of osteophytes. <i>Myco. tuberculosis</i> was found in the pus from the psoas abscess. (This patient refused all treatment and yet obtained healing of his bony focus.) — — — — —	306
FIG. 94.—Six years after ischio-femoral arthrodesis for active disease in a man aged 52 years. Designed to shorten his treatment, the operation has failed in every way — — — — —	310
FIG. 95.—Disease of the lumbar spine still active in a man aged 20 years. At this stage an Albee fusion was done "to give him a hope of quicker cure" — —	312
FIG. 96.—The same lesion as in Fig. 95; fifteen months later the disease is still active and further destruction has occurred. Healing is being prevented because the graft is holding the diseased area open — — — — —	313

LIST OF ILLUSTRATIONS

	PAGE
FIG. 97.—Ilio-femoral fusion for tuberculosis of the hip in a woman aged 30 years	314
FIG. 98.—Diagram of Trumble's method of ischio-femoral fusion by a low extra-articular cortical graft	315
FIG. 99.—Diagram of Brittain's ischio-femoral arthrodesis	315
FIG. 100.—Double tibial onlay grafts (Girdlestone) for fusion of a lumbar focus in a man aged 26 years after 30 months of conservative treatment had produced quiescence of the disease	317
FIG. 101.—Ischio-femoral arthrodesis (Brittain) of para-articular type. X-ray appearances 11 months after operation in a woman aged 23 years	317
FIG. 102.—The Albee hip fusion as performed in 1913	318
FIG. 103.—The Hibbs hip fusion	318
FIG. 104.—A soundly united ilio-femoral fusion by tibial graft in a man aged 22 years, performed after 2½ years of conservative treatment	319
FIG. 105.—Arthrodesis of a tuberculous knee in a woman aged 47 years, using the Brittain cross-graft technique	319
FIG. 106.—(a) Arthrodesis of a tuberculous knee using Charnley's clamps. (b) Bony union five months later	320
FIG. 107.—X-ray 18 months after fusion of a quiescent tuberculous shoulder by the McKee-Brittain technique (posterior scapulo-humeral fusion with tibial graft) in a woman aged 22 years	321
FIG. 108.—While one focus in the dorsal spine of a man aged 48 years has healed and consolidated, a second has developed 2 segments below	322
FIG. 109.—Antero-lateral decompression for Pott's paraplegia	328
FIG. 110.—Illustrating why a kyphosis helps in bringing the compressing agent within easy access in antero-lateral decompression	328
FIG. 111.—(a) and (b) X-ray films of a dorsal spine lesion in a woman aged 25 years with profound paraplegia in extension	329
FIG. 112.—Same case as Fig. 111. Illustrated are films obtained two years later, showing the amount of bone removed, and also showing that the operation in no way impedes healing of the skeletal lesion	330
FIG. 113.—Endometrium with tuberculous endometritis without caseation	339
FIG. 114.—Tubercle follicle with giant cells, epithelioid proliferation and central caseation	339
FIG. 115.—A small tuberculoma in the cerebellum, reaching the surface in the depth of the sulcus. Typical tuberculous exudate is seen in the meninges on the upper surface of the cerebellum and is particularly marked in the interpeduncular fossa	354
FIG. 116.—Encapsulated tuberculoma in the lenticular nucleus destroying the internal capsule. Below, the optic chiasma is surrounded by exudate	354
FIG. 117.—Enlarged photograph of base of brain showing thick exudate filling the interpeduncular fossa, obscuring the optic chiasma and covering the upper part of the anterior aspect of the pons. The basilar and carotid arteries are obscured, as are the origins of the third nerves	355
FIG. 118.—Extensive dense exudate filling the interpeduncular fossa and burying the basilar artery. Extensions of this exudate are seen extending laterally and dorsally in the cisterna ambiens encircling the mid-brain. The pineal body is embedded in exudate	357