


经典英文化学教材系列

Modern Methods of Organic Synthesis

Fourth Edition

有机合成现代方法 第4版

William Carruthers and Iain Coldham



CAMBRIDGE

世界图书出版公司
www.wpcbj.com.cn

MODERN METHODS OF ORGANIC SYNTHESIS

W. CARRUTHERS

Formerly of the University of Exeter

IAIN COLDHAM

University of Sheffield



CAMBRIDGE
UNIVERSITY PRESS

图书在版编目 (C I P) 数据

有机合成现代方法: 第4版: 英文 / (英) 卡如萨
(Carruthers, W.) 著. —北京: 世界图书出版公司北
京公司, 2008.10

书名原文: Modern Methods of Organic Synthesis 4th
ed.

ISBN 978-7-5062-9270-2

I. 有… II. 卡… III. 有机合成-英文 IV. 0621.3

中国版本图书馆CIP数据核字 (2008) 第136101号

书 名: Modern Methods of Organic Synthesis 4th ed.

作 者: William Carruthers & Lain Coldham

中译名: 有机合成现代方法 第4版

责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河国英印务有限公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64015659

电子信箱: kjsk@vip.sina.com

开 本: 16开

印 张: 32

版 次: 2008 年 10 月第 1 次印刷

版权登记: 图字:01-2008-4043

书 号: 978-7-5062-9270-2 / O · 632

定 价: 99.00 元

Preface to the first edition

This book is addressed principally to advanced undergraduates and to graduates at the beginning of their research careers, and aims to bring to their notice some of the reactions used in modern organic syntheses. Clearly, the whole field of synthesis could not be covered in a book of this size, even in a cursory manner, and a selection has had to be made. This has been governed largely by consideration of the usefulness of the reactions, their versatility and, in some cases, their selectivity.

A large part of the book is concerned with reactions which lead to the formation of carbon-carbon single and double bonds. Some of the reactions discussed, such as the alkylation of ketones and the Diels-Alder reaction, are well established reactions whose scope and usefulness has increased with advancing knowledge. Others, such as those involving phosphorus ylids, organoboranes and new organometallic reagents derived from copper, nickel, and aluminium, have only recently been introduced and add powerfully to the resources available to the synthetic chemist. Other reactions discussed provide methods for the functionalisation of unactivated methyl and methylene groups through intramolecular attack by free radicals at unactivated carbon-hydrogen bonds. The final chapters of the book are concerned with the modification of functional groups by oxidation and reduction, and emphasise the scope and limitations of modern methods, particularly with regard to their selectivity.

Discussion of the various topics is not exhaustive. My object has been to bring out the salient features of each reaction rather than to provide a comprehensive account. In general, reaction mechanisms are not discussed except in so far as is necessary for an understanding of the course or stereochemistry of a reaction. In line with the general policy in the series references have been kept to a minimum. Relevant reviews are noted but, for the most part, references to the original literature are given only for points of outstanding interest and for very recent work. Particular reference is made here to the excellent book by H. O. House, *Modern Synthetic*

Reactions which has been my guide at several points and on which I have tried to build, I feel all too inadequately.

I am indebted to my friend and colleague, Dr K. Schofield, for much helpful comment and careful advice which has greatly assisted me in writing the book.

26 October 1970

Preface to the fourth edition

Some Modern Methods of Organic Synthesis was originally written by Dr W. (Bill) Carruthers, and three popular editions were published that have helped many students of advanced organic chemistry. Unfortunately, Dr Carruthers died in 1990, just prior to his retirement. As his successor at the University of Exeter, it was appropriate that I should take on the task of preparing the fourth edition of this text. In honour of Dr Carruthers, a similar format to previous editions has been taken, although of course the book has been completely re-written and brought up-to-date (through 2003) to take account of the many advances in the subject since the third edition was published. As in previous editions, the text begins with descriptions of some of the most important methods for the formation of carbon–carbon bonds, including the use of enolates and organometallic compounds for carbon–carbon single-bond formation (Chapter 1), methods for carbon–carbon double-bond formation (Chapter 2), pericyclic reactions (Chapter 3), radicals and carbenes (Chapter 4). There has been some re-organization of material and emphasis has been placed on reactions that are useful, high yielding or selective for organic synthesis. For example, Chapter 1 has been expanded to include some of the most popular and contemporary reactions using main-group and transition-metal chemistry (rather than placing reactions of organoboron and silicon compounds into a separate chapter). A new chapter describing the functionalization of alkenes has been devised, covering reactions such as hydroboration, epoxidation and dihydroxylation (Chapter 5). The book concludes with examples of pertinent oxidation (Chapter 6) and reduction (Chapter 7) reactions that are used widely in organic synthesis. The opportunity has been taken to add some problems at the end of each chapter, with answers at the end of the book. References have been compiled as footnotes on each relevant page for ease of use.

In common with the previous editions, the book is addressed principally to advanced undergraduates and to graduates at the beginning of their research careers. My aim has been to bring out the salient features of the reactions and reagents

rather than to provide a comprehensive account. Reaction mechanisms are not normally discussed, except where necessary for an understanding of the course or stereochemistry of a reaction. My hope is that the book will find widespread use as a helpful learning and reference aid for synthetic chemists, and that it will be a fitting legacy to Dr Carruthers.

The majority of the text was written at the University of Exeter before my move to the University of Sheffield and I would like to acknowledge the encouragement and help of the staff at Exeter.

Part of one chapter was written while I was a Visiting Professor at the University of Miami, and I am grateful to Professor Bob Gawley for hosting my visit. My thanks extend to various people who have proof-read parts of the text, including Chris Moody, Mike Shipman, Mark Wood, Alison Franklin, Joe Harrity, Steve Pih and Ben Dobson. Finally, I would like to thank my family for their patience during the writing of this book.

I. Coldham
January 2004

Contents

<i>Preface to the first edition</i>	<i>page</i> vii
<i>Preface to the fourth edition</i>	ix
1 Formation of carbon–carbon single bonds	1
1.1 Main-group chemistry	1
1.1.1 Alkylation of enolates and enamines	1
1.1.2 Conjugate addition reactions of enolates and enamines	19
1.1.3 The aldol reaction	27
1.1.4 Asymmetric methodology with enolates and enamines	36
1.1.5 Organolithium reagents	45
1.1.6 Organomagnesium reagents	63
1.1.7 Organozinc reagents	67
1.1.8 Allylic organometallics of boron, silicon and tin	71
1.2 Transition-metal chemistry	75
Problems	101
1.2.1 Organocopper reagents	75
1.2.2 Organochromium chemistry	81
1.2.3 Organocobalt chemistry	85
1.2.4 Organopalladium chemistry	89
2 Formation of carbon–carbon double bonds	105
2.1 β -Elimination reactions	105
2.2 Pyrolytic <i>syn</i> eliminations	111
2.3 Fragmentation reactions	118
2.4 Alkenes from hydrazones	120
2.5 Alkenes from 1,2-diols	123
2.6 Alkenes from alkynes	125
2.7 The Wittig and related reactions	132

2.8	Alkenes from sulfones	144
2.9	Alkenes using titanium or chromium reagents	148
2.10	Alkene metathesis reactions	151
	Problems	155
3	Pericyclic reactions	159
3.1	The Diels–Alder cycloaddition reaction	159
3.1.1	The dienophile	162
3.1.2	The diene	174
3.1.3	Regiochemistry of the Diels–Alder reaction	185
3.1.4	Stereochemistry of the Diels–Alder reaction	188
3.1.5	Intramolecular Diels–Alder reactions	193
3.1.6	The retro Diels–Alder reaction	199
3.1.7	Asymmetric Diels–Alder reactions	202
3.2	[2+2] Cycloaddition reactions	211
3.3	Cycloaddition reactions with allyl cations and allyl anions	219
3.4	1,3-Dipolar cycloaddition reactions	222
3.5	The ene reaction	231
3.6	[3,3]-Sigmatropic rearrangements	238
3.6.1	The Cope rearrangement	239
3.6.2	The Claisen rearrangement	244
3.7	[2,3]-Sigmatropic rearrangements	253
3.8	Electrocyclic reactions	259
	Problems	264
4	Radical and carbene chemistry	268
4.1	Radicals	268
4.1.1	Radical abstraction reactions	269
4.1.2	Radical addition reactions	280
4.2	Carbenes	299
	Problems	312
5	Functionalization of alkenes	315
5.1	Hydroboration	315
5.1.1	Reactions of organoboranes	322
5.2	Epoxidation and aziridination	331
5.2.1	Epoxidation	331
5.2.2	Asymmetric epoxidation	337
5.2.3	Aziridination	346

5.3	Dihydroxylation	349
5.3.1	Dihydroxylation with osmium tetroxide	349
5.3.2	Other methods of dihydroxylation	355
5.3.3	Amino-hydroxylation	358
5.4	Oxidative cleavage	360
5.5	Palladium-catalysed oxidation of alkenes	365
	Problems	367
6	Oxidation	370
6.1	Oxidation of hydrocarbons	370
6.1.1	Alkanes	370
6.1.2	Aromatic hydrocarbons	371
6.1.3	Alkenes	374
6.2	Oxidation of alcohols	378
6.2.1	Chromium reagents	378
6.2.2	Oxidation via alkoxysulfonium salts	381
6.2.3	Manganese reagents	384
6.2.4	Other metal-based oxidants	386
6.2.5	Other non-metal-based oxidants	389
6.2.6	Oxidation to carboxylic acids or esters	392
6.3	Oxidation of ketones	394
6.3.1	α,β -Unsaturated ketones	394
6.3.2	α -Hydroxy-ketones	396
6.3.3	Baeyer–Villiger oxidation of ketones	398
	Problems	402
7	Reduction	405
7.1	Catalytic hydrogenation	405
7.2	Reduction by dissolving metals	422
7.3	Reduction by hydride-transfer reagents	434
7.3.3	Derivatives of lithium aluminium hydride and sodium borohydride	443
7.3.4	Mixed lithium aluminium hydride–aluminium chloride reagents	444
7.3.5	Diisobutylaluminium hydride (DIBAL-H)	445
7.3.6	Sodium cyanoborohydride and sodium triacetoxymborohydride	446
7.3.7	Borane and derivatives	449

7.4 Other methods of reduction	454
7.4.1 Enzyme catalysed	454
7.4.2 Wolff–Kishner reduction	457
7.4.3 Reductions with diimide	459
7.4.4 Reductions with trialkylsilanes	460
Problems	462
<i>Answers to problems</i>	466
<i>Index</i>	487

1

Formation of carbon–carbon single bonds

The formation of carbon–carbon single bonds is of fundamental importance in organic synthesis. As a result, there is an ever-growing number of methods available for carbon–carbon bond formation. Many of the most useful procedures involve the addition of organometallic species or enolates to electrophiles, as in the Grignard reaction, the aldol reaction, the Michael reaction, alkylation reactions and coupling reactions. Significant advances in both main-group and transition-metal-mediated carbon–carbon bond-forming reactions have been made over the past decade. Such reactions, which have been finding useful application, are discussed in this chapter. The formation of carbon–carbon single bonds by pericyclic or radical reactions are discussed in chapters 3 and 4.

1.1 Main-group chemistry

1.1.1 Alkylation of enolates and enamines


It is well known that carbonyl groups increase the acidity of the proton(s) adjacent (α -) to the carbonyl group. Table 1.1 shows the pK_a values for some unsaturated compounds and for some common solvents and reagents.

The acidity of the C–H bonds in these compounds is caused by a combination of the inductive electron-withdrawing effect of the unsaturated groups and the resonance stabilization of the anion formed by removal of a proton (1.1). Not all groups are equally effective in ‘activating’ a neighbouring CH; nitro is the most powerful of the common groups, with the series following the approximate order $\text{NO}_2 > \text{COR} > \text{SO}_2\text{R} > \text{CO}_2\text{R} > \text{CN} > \text{C}_6\text{H}_5$. Two activating groups reinforce each other; for example, diethyl malonate has a lower pK_a (≈ 13) than ethyl acetate ($pK_a \approx 24$). Acidity is increased slightly by electronegative substituents


Table 1.1. Approximate acidities of some activated compounds and common reagents

Compound	pK _a	Compound	pK _a
CH ₃ CO ₂ H	5	C ₆ H ₅ COCH ₃	19
CH ₂ (CN)CO ₂ Et	9	CH ₃ COCH ₃	20
CH ₂ (COCH ₃) ₂	9	CH ₃ CO ₂ Et	24
CH ₃ NO ₂	10	CH ₃ CN	25
CH ₃ COCH ₂ CO ₂ Et	11	((CH ₃) ₃ Si) ₂ NH	26
CH ₂ (CO ₂ Et) ₂	13	CH ₃ SO ₂ CH ₃	31
CH ₃ OH	16	CH ₃ SOCH ₃	35
(CH ₃) ₃ COH	19	((CH ₃) ₂ CH) ₂ NH	36

(e.g. sulfide) and decreased by alkyl groups.



By far the most important activating group in synthesis is the carbonyl group. Removal of a proton from the α -carbon atom of a carbonyl compound with base gives the corresponding enolate anion. It is these enolate anions that are involved in many reactions of carbonyl compounds, such as the aldol condensation, and in bimolecular nucleophilic displacements (alkylations, as depicted in Scheme 1.2).



Enolate anions should be distinguished from enols, which are always present in equilibrium with the carbonyl compound (1.3). Most monoketones and esters contain only small amounts of enol (<1%) at equilibrium, but with 1,2- and 1,3-dicarbonyl compounds much higher amounts of enol (>50%) may be present. In the presence of a protic acid, ketones may be converted largely into the enol form,

implicated in many acid-catalysed reactions of carbonyl compounds.




Table 1.1 illustrates the relatively high acidity of compounds in which a C—H bond is activated by two or more carbonyl (or cyano) groups. It is therefore possible to use a comparatively weak base, such as a solution of sodium ethoxide in ethanol, in order to form the required enolate anion. An equilibrium is set up, as illustrated in Scheme 1.4, in which the conjugate acid of the base (BH) must be a weaker acid than the active methylene compound. Another procedure for preparing the enolate of an active methylene compound is to use sodium hydride (or finely divided sodium or potassium metal) in tetrahydrofuran (THF), diethyl ether (Et₂O) or benzene. The metal salt of the enolate is formed irreversibly with evolution of hydrogen gas. β-Diketones can often be converted into their enolates with alkali-metal hydroxides or carbonates in aqueous alcohol or acetone.



Much faster alkylation of enolate anions can often be achieved in dimethylformamide (DMF), dimethylsulfoxide (DMSO) or 1,2-dimethoxyethane (DME) than in the usual protic solvents. The presence of hexamethylphosphoramide (HMPA) or a triamine or tetramine can also enhance the rate of alkylation. This is thought to be because of the fact that these solvents or additives solvate the cation, but not the enolate, thereby separating the cation–enolate ion pair. This leaves a relatively free enolate ion, which would be expected to be a more reactive nucleophile than the ion pair.¹ Reactions with aqueous alkali as base are often improved in the presence of a phase-transfer catalyst such as a tetra-alkylammonium salt.²


Alkylation of enolate anions is achieved readily with alkyl halides or other alkylating agents.³ Both primary and secondary alkyl, allyl or benzyl halides may be used successfully, but with tertiary halides poor yields of alkylated product often result because of competing elimination. It is sometimes advantageous to proceed by way of the toluene-*p*-sulfonate, methanesulfonate or trifluoromethanesulfonate rather than a halide. The sulfonates are excellent alkylating agents and can usually be obtained from the alcohol in a pure condition more readily than

¹ H. E. Zaugg, D. A. Dunnigan, R. J. Michaels, L. R. Swett, T. S. Wang, A. H. Sommers and R. W. DeNet, *J. Org. Chem.*, **26** (1961), 644; A. J. Parker, *Quart. Rev. Chem. Soc. Lond.*, **16** (1962), 163; M. Goto, K. Akimoto, K. Aoki, M. Shindo and K. Koga, *Tetrahedron Lett.*, **40** (1999), 8129.


² M. Makosza and A. Jonczyk, *Org. Synth.*, **55** (1976), 91.

³ D. Caine, in *Comprehensive Organic Synthesis*, ed. B. M. Trost and I. Fleming, vol. 3 (Oxford: Pergamon Press, 1991), p. 1.

the corresponding halides. Primary and secondary alcohols can be used as alkylating agents under Mitsunobu conditions.⁴ Epoxides have also been used, generally reacting at the less substituted carbon atom. Attack of the enolate anion on the alkylating agent takes place by an S_N2 pathway and thus results in inversion of configuration at the carbon atom of the alkylating agent (1.5).⁵




With secondary and tertiary allylic halides or sulfonates, reaction of an enolate anion may give mixtures of products formed by competing attack at the α - and γ -positions (1.6). Addition of the enolate anion to a π -allylpalladium complex provides an alternative method for allylation (see Section 1.2.4).



A difficulty sometimes encountered in the alkylation of active methylene compounds is the formation of unwanted dialkylated products. During the alkylation of the sodium salt of diethylmalonate, the monoalkyl derivative formed initially is in equilibrium with its anion. In ethanol solution, dialkylation does not take place to any appreciable extent because ethanol is sufficiently acidic to reduce the concentration of the anion of the alkyl derivative, but not that of the more acidic diethylmalonate itself, to a very low value. However, replacement of ethanol by an inert solvent favours dialkylation. Dialkylation also becomes a more serious problem with the more acidic cyanoacetic esters and in alkylations with very reactive electrophiles such as allyl or benzyl halides or sulfonates.


Dialkylation may, of course, be effected deliberately if required by carrying out two successive operations, by using either the same or a different alkylating agent in the two steps. Alkylation of dihalides provides a useful route to three- to seven-membered ring compounds (1.7). Non-cyclic products are formed at the same time by competing intermolecular reactions and conditions have to be chosen carefully to suppress their formation (for example, by using high dilution).




⁴ O. Mitsunobu, *Synthesis* (1981), 1; J. Yu, J.-Y. Lai and J. R. Falck, *Synlett* (1995), 1127; T. Tsunoda, C. Nagino, M. Oguri and S. Itô, *Tetrahedron Lett.*, **37** (1996), 2459.

⁵ T. Sato and J. Otera, *J. Org. Chem.*, **60** (1995), 2627.

Under ordinary conditions, aryl or alkenyl halides do not react with enolate anions, although reaction can occur with aryl halides bearing strongly electro-negative substituents in the *ortho* and *para* positions. 2,4-Dinitrochlorobenzene, for example, with ethyl cyanoacetate gives ethyl (2,4-dinitrophenyl)cyanoacetate (90%) by an addition–elimination pathway. Unactivated aryl halides may react with enolates under more vigorous conditions, particularly sodium amide in liquid ammonia. Under these conditions, the reaction of bromobenzene with diethyl-malonate, for example, takes place by an elimination–addition sequence in which benzyne is an intermediate (1.8).




Enolate anions with extended conjugation can be formed by proton abstraction of α,β -unsaturated carbonyl compounds (1.9). Kinetically controlled alkylation of the delocalized anion takes place at the α -carbon atom to give the β,γ -unsaturated compound directly. A similar course is followed in the kinetically controlled protonation of such anions.



A wasteful side reaction which sometimes occurs in the alkylation of 1,3-dicarbonyl compounds is the formation of the *O*-alkylated product. For example, reaction of the sodium salt of cyclohexan-1,3-dione with butyl bromide gives the *O*-alkylated product (37%) and only 15% of the *C*-alkylated 2-butylcyclohexan-1,3-dione. In general, however, *O*-alkylation competes significantly with *C*-alkylation only with reactive methylene compounds in which the equilibrium concentration of enol is relatively high (as in 1,3-dicarbonyl compounds). The extent of *C*- versus *O*-alkylation for a particular 1,3-dicarbonyl compound depends on the choice of cation, solvent and electrophile. Cations (such as Li^+) that are more covalently bound to the enolate oxygen atom or soft electrophiles (such as alkyl halides) favour *C*-alkylation, whereas cations such as K^+ or hard electrophiles (such as alkyl sulfonates) favour *O*-alkylation.

Alkylation of malonic esters and other active methylene compounds is useful in synthesis because the alkylated products can be subjected to hydrolysis and decarboxylation (1.10). Direct decarboxylation under neutral conditions with an alkali metal salt (e.g. lithium chloride) in a dipolar aprotic solvent (e.g. DMF) is a popular alternative method.⁶



Proton abstraction from a monofunctional carbonyl compound (aldehyde, ketone, ester, etc.) is more difficult than that from a 1,3-dicarbonyl compound. Table 1.1 illustrates that a methyl or methylene group which is activated by only one carbonyl or cyano group requires a stronger base than ethoxide or methoxide ion to convert it to the enolate anion in high enough concentration to be useful for subsequent alkylation. Alkali-metal salts of tertiary alcohols, such as tert-butanol, in the corresponding alcohol or an inert solvent, have been used with success, but suffer from the disadvantage that they are not sufficiently basic to convert the ketone completely into the enolate anion. This therefore allows the possibility of an aldol reaction between the anion and unchanged carbonyl compound. An alternative procedure is to use a much stronger base that will convert the compound completely into the anion. Traditional bases of this type are sodium and potassium amide or sodium hydride, in solvents such as diethyl ether, benzene, DME or DMF. The alkali-metal amides are often used in solution in liquid ammonia. Although these bases can convert ketones essentially quantitatively into their enolate anions, aldol reaction may again be a difficulty with these bases because of the insolubility of the reagents. Formation of the anion takes place only slowly in the heterogeneous reaction medium and both the ketone and the enolate ion are present at some stage. This difficulty does not arise with the lithium dialkylamides, such as lithium diisopropylamide (LDA) or lithium 2,2,6,6-tetramethylpiperidide (LTMP) or the alkali-metal salts of bis(trimethylsilyl)amine (LHMDS, NaHMDS and KHMDS), which are soluble in non-polar solvents. These bases are now the most commonly used reagents for the generation of enolates.

An example illustrating the intermolecular alkylation of an ester is given in Scheme 1.11. Intramolecular alkylations also take place readily in appropriate cases and reactions of this kind have been used widely in the synthesis of cyclic compounds. In such cases, the electrophilic centre generally approaches the enolate

⁶ A. P. Krapcho, *Synthesis* (1982), 805; 893.