



Fudan Series in Graduate Textbooks



Advanced Quantum Mechanics

高等量子力学

苏汝铨 王 斌 编 著

Ru-Keng Su Bin Wang

复旦大学出版社



Fudan University Press



Fudan Series in Graduate Textbooks




Advanced Quantum Mechanics

高等量子力学

苏汝铨 王 斌 编 著
Ru-Keng Su Bin Wang

复旦大学出版社

 Fudan University Press

图书在版编目(CIP)数据

高等量子力学 = Advanced Quantum Mechanics/苏汝铿,王斌编著.
—上海:复旦大学出版社,2004.6
(博学·研究生系列)
ISBN 7-309-04013-9

I. 高… II. ①苏…②王… III. 量子力学-研究生-教材
IV. 0413.1

中国版本图书馆 CIP 数据核字(2004)第 040878 号

Advanced Quantum Mechanics

高等量子力学

苏汝铿 王 斌 编著

出版发行 复旦大学出版社

上海市国权路 579 号 邮编 200433

86-21-65118853(发行部) 86-21-65109143(邮购)

fupnet@fudanpress.com <http://www.fudanpress.com>

责任编辑 梁 玲

装帧设计 陈 萍

总 编 辑 高若海

出 品 人 贺圣遂

印	刷	上海浦东北联印刷厂
开	本	787×960 1/16
印	张	15.25 插页 2
字	数	315 千
版	次	2004 年 6 月第一版第一次印刷
印	数	1—2 500

书	号	ISBN 7-309-04013-9/O · 321
定	价	30.00 元

如有印装质量问题,请向复旦大学出版社发行部调换。

版权所有 侵权必究

**In celebration of
the 100th anniversary of Fudan University**

(1905-2005)

献给复旦大学一百周年校庆

作者简介

苏汝铿，1938年5月生于广东。1960年毕业于北京大学物理系。从1960年开始，一直在复旦大学物理系任教至今。著有《量子力学》、《统计物理学》、《物理学的挑战——物理学前沿和基础课题选》等书。主要从事温度场论、中高能核物理、量子场论、广义相对论、宇宙学及天体物理学等方面的研究工作，已发表学术论文200余篇，曾多次获国家教委、上海市及中国科学院科技进步奖及自然科学奖。现为教授、博士生导师。

王斌，1967年10月生于上海。1998年于复旦大学物理系获理学博士学位。2001年7月开始在复旦大学物理系任教授，2002年12月被聘为博士生导师。目前的研究领域为引力论和宇宙论，至今和合作者共发表论文50余篇。在复旦大学担任过《高等量子力学》、《广义相对论》、《电动力学》等研究生和本科生的教学工作。

Abstract

This book is intended for use as a textbook for a graduate course in Advanced Quantum Mechanics, and as a reference book for workers in the field.

Before the subject is presented, a brief but self-contained review on the foundation of Quantum Mechanics is given. A large part of this book is devoted to select applications of quantum mechanics, such as scattering theory, second quantization, superconductive theory, superfluidity, phase of wave function, path integral and relativistic quantum mechanics. The selection is guided by the interest of topic to physicists, its value as an illustration of calculating techniques, and our personal taste.

To read the book, the reader need basic knowledge of quantum mechanics, some intuitive feeling for electrodynamics and special relativity, and a good mathematical knowledge.

PREFACE

As an advanced study of quantum mechanics, this textbook deals with topics likely to be encountered in a graduate-level course. The authors believe there is no material difference between quantum mechanics and advanced quantum mechanics except that the latter targets an in-depth study. However, we can see basic differences between advanced quantum mechanics and quantum statistics and/or quantum field theory in that the former does not cope with effects of temperature and quantization of fields.

This book is written for advanced-level students or graduate students studying physics or the related disciplines. There are all together six chapters in this book, centering on such topics as relativistic quantum mechanics, path intergral, scattering theory, second quantization, superconductive theory, superfluidity as well as the phase of wave function. Though the textbook adopts a rigorous style, it is easy to read since no other background knowledge than the basics of quantum mechanics is required. In addition, the first chapter serves to equip the reader with the necessary basic knowledge. Throughout the book, the authors take painstaking efforts to demonstrate the mathematical steps and intermediate calculations in a thorough way, which will enable the reader to understand with ease.

We owe our thanks to Prof. Xun Wang for his encouragement, Prof. L. K. Wang for editing this book and Dr. Wei-Liang Qian for preparing the LATEX format for the current edition. We would like to dedicate this book to the 100th anniversary of Fudan University.

Ru-Keng Su
Bin Wang
Apr. 2004.
Fudan University

前 言

高等量子力学是一些高年级本科生及研究生学习的量子力学课程。作者认为,量子力学和高等量子力学并没有本质的不同,只不过后者涉及的面更深和更广。但是,高等量子力学与量子统计和量子场论有本质的不同,前者无需考虑温度效应和场的量子化。

本书是为高年级本科生或研究生学习物理学或相关课程所准备的。全书共分六章,包括相对论量子力学、路径积分、散射理论、二次量子化、超导理论、超流动性以及波函数的位相等。本书叙述严谨,但易于阅读和学习。它只要求读者具有在本书第一章中给出的量子力学的基础知识,而不要求其他。本书的数学推导力求详尽,以便读者更易于掌握。

感谢王迅教授的支持和鼓励,感谢汪礼康教授校订本书及钱卫良博士为本书准备的 LaTeX 文件。我们希望将本书献给母校——复旦大学一百周年校庆。

苏汝铨 王斌
2004 年 4 月于复旦大学

CONTENTS

PREFACE	1
Chapter 1 Foundation of Quantum Mechanics	1
1.1 State Vector, Wave Function and Superposition of States	1
1.2 Schrödinger Equation and Its Solutions	3
1.3 Operators	9
1.4 Approximation Methods	13
1.5 WKB Approximation	17
1.6 Density Matrix	27
Chapter 2 Scattering Theory	31
2.1 General Description of the Scattering Theory	31
2.2 The Partial Wave Method	37
2.3 Examples of the Partial Wave Method	44
2.4 Green Function Method and Born Approximation	49
2.5 Center-of-mass Coordinate and the Laboratory Coordinate	60
2.6 T Matrix	65
2.7 Lippman-Schwinger Equation	67
2.8 Dyson Equation	71
2.9 Scattering Matrix(S Matrix)	75
2.10 Scattering in the Complex Potential	81
2.11 General Theory of the Inelastic Scattering	84
2.12 Approximation of the Distortion Wave	90
Chapter 3 Many Body Problem	93
3.1 Second Quantization	93
3.2 Hartree-Fock Mean Field Approximation	103

3.3	Thomas-Fermi Method	107
3.4	Superconductive Theory	111
3.5	Superfluidity Theory	118
Chapter 4 Path Integral		124
4.1	Classical Action and the Amplitude in Quantum Mechanics	125
4.2	Path Integral	129
4.3	Gauss Integration	138
4.4	Path Integral and the Schrödinger Equation	142
4.5	The Canonical Form of the Path Integral	148
Chapter 5 Relativistic Quantum Mechanics		152
5.1	Klein-Gordon Equation	153
5.2	Dirac Equation	158
5.3	Solutions of the Free Field Dirac Equation	162
5.4	Dirac Equation in the Electromagnetic Field	169
5.5	Covariant Form of the Dirac Equation	171
5.6	Dirac Equation in the Central Force Field	178
5.7	Solution of the Dirac Equation in the Coulomb Field	187
5.8	Klein Paradox	192
5.9	MIT Bag Model	202
5.10	Chiral symmetry	207
Chapter 6 Phase of Wave Function		211
6.1	Aharonov-Bohm Effect	211
6.2	Aharonov-Casher Effect	216
6.3	Magnetic Flux of Superconductive Ring	217
6.4	Monopole	219
6.5	Berry Phase	222
6.6	Non-integrable Phase Factor	228
6.7	Vacuum Energy and Casimir Effect	230
Index		235

Foundation of Quantum Mechanics

1.1 State Vector, Wave Function and Superposition of States

Advanced quantum mechanics is a continuing course of *quantum mechanics* at undergraduate level. There is no material difference between the two courses except that the latter presents a detailed and insightful introduction to the fundamental concepts and their simple applications, but the former targets an in-depth study. This book introduces relativistic quantum mechanics, path integral, scattering theory, second quantization, superconductive theory, superfluidity as well as the further study of the phase of wave function. This chapter evolves from an attempt of a brief review over the basic ideas and formulae in undergraduate-level quantum mechanics. The details of this chapter can be found in the usual references of quantum mechanics^①.

In classical mechanics, the state of a particle is described in terms of its coordinate and momentum at some instant of time. The evolution of the state complies with the Newton equations or the Hamiltonian canonical equations. With the initial data of coordinate and momentum at a given instant, the Newton equation of motion can predict the behavior of the particle at all subsequent instants. In quantum mechanics the prediction is in principle impossible because the exact coordinate and the corresponding momentum cannot be given according to the uncertainty

① For example, Ru-Keng Su, *Quantum Mechanics*, Second Edition. High Education Press, 2002

principle.

In quantum mechanics, because of the wave-particle duality, the state of a quantum system is described in terms of a wave vector in a Hilbert space. The wave vector can be expressed as a wave function in a specific representation, i. e. a complete set of basics of the Hilbert space. For example, the eigenfunctions of coordinate operator x are δ -functions

$$x \delta(x - x') = x' \delta(x - x') . \quad (1.1.1)$$

If we choose the δ -functions to be the basics of the Hilbert space, the wave function at instant t is expressed as $\psi(x, t)$. According to the Born statistic explanation of wave function, the square of the modulus of this function determines the probability distribution of the values of the coordinates: $|\psi|^2 dx$ is the probability that a measurement made on the system will find the values of coordinates in the element dx of the configuration space. Since the sum of the probabilities of all possible values of the system coordinates must be unity, the wave function $\psi(x, t)$ satisfies the normalization condition

$$\int |\psi(x, t)|^2 dx = 1 . \quad (1.1.2)$$

If the integral of $|\psi|^2$ diverges and ψ cannot be normalized by using Eq. (1.1.2), $|\psi|^2$ will not determine the absolute values of the probability of the coordinates, but the ratio of the values of $|\psi|^2$ at two different points will still determine the relative probability of the corresponding values of coordinates.

Quantum states follow a general principle of superposition. Suppose a state with wave function $\psi_m(q)$, where measuring on it leads to a definite result m , and a state with $\psi_n(q)$ leads to result n . It is assumed that every linear combination of ψ_m and ψ_n ,

$$\psi = c_m \psi_m + c_n \psi_n \quad (1.1.3)$$

gives a state in which the measurement leads to either result m or n . The probability of belting result m or n is $\frac{|c_m|^2}{|c_m|^2 + |c_n|^2}$ or $\frac{|c_n|^2}{|c_m|^2 + |c_n|^2}$. If

ψ_i is an eigenfunction of Hermitian operator \hat{F} and f_i is the corresponding eigenvalue, then all eigenfunctions of \hat{F} form a complete set $\{\psi_i\}$ in Hilbert space. A wave function ψ can be expanded into

$$\psi = \sum_i c_i \psi_i. \quad (1.1.4)$$

If we make a measurement of dynamical quantity F in a state with wave function ψ , we reach the result that the square modulus $|c_i|^2$ of each coefficient in Eq. (1.1.4) determines the probability of the corresponding value f_n of the quantity F in a state with ψ provided that the wave functions satisfy

$$\int \psi^* \psi \, d\mathbf{r} = 1, \quad (1.1.5)$$

$$\int \psi_m^* \psi_n \, d\mathbf{r} = \delta_{mn}, \quad (1.1.6)$$

and

$$\sum_i |c_i|^2 = 1. \quad (1.1.7)$$

It is worth noting that the superposition of quantum states in quantum mechanics is with essentially different nature from that in the classical theory. The quantum coherence cannot be found in classical mechanics.

1.2 Schrödinger Equation and Its Solutions

In quantum mechanics, as in classical mechanics, a system is governed by the equation of motion which tells us that the state at one time determines the state at a later time. In classical mechanics, the equation of motion is Newton equation. In quantum mechanics, Schrödinger assumed that the equation of motion for wave function reads

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H}\psi = \left[-\frac{\hbar^2}{2m} \nabla^2 + U(\mathbf{r}, t) \right] \psi, \quad (1.2.1)$$

where

$$\hat{H} = -\frac{\hbar^2}{2m} \nabla^2 + U(\mathbf{r}, t) \quad (1.2.2)$$

is the Hamilton operator and $U(\mathbf{r}, t)$ is the potential energy. If the potential energy depends on \mathbf{r} alone, we can prove that the wave function becomes

$$\psi(\mathbf{r}, t) = \psi(\mathbf{r}) e^{-iEt/\hbar}. \quad (1.2.3)$$

Eq. (1.2.3) is called stationary state. In this state, the density of probability does not depend on time

$$|\psi(\mathbf{r}, t)|^2 = |\psi(\mathbf{r})|^2, \quad (1.2.4)$$

and the energy has a definite value. Schrödinger equation for stationary state reads

$$H\psi = E\psi, \quad (1.2.5)$$

$$\hat{H} = -\frac{\hbar^2}{2m} \nabla^2 + U(\mathbf{r}). \quad (1.2.6)$$

In principle, the solutions of Schrödinger equation can be obtained for definite boundary conditions. However, due to mathematical problems, only a few exact solutions have been found. Examples are given as follows:

1.2.1 One-dimensional infinite potential well

Considering one-dimension Schrödinger equation with potential

$$U(x) = \begin{cases} 0 & (|x| < a) \\ \infty & (|x| \geq a) \end{cases}, \quad (1.2.7)$$

we find that the energy levels and the normalized wave function are

$$E_n = \frac{n^2 \pi^2 \hbar^2}{8ma^2} \quad (n = 1, 2, 3, \dots), \quad (1.2.8)$$

$$\psi_n = \begin{cases} \frac{1}{\sqrt{a}} \sin \frac{n\pi}{2a} (x + a) & (|x| < a) \\ 0 & (|x| \geq a) \end{cases}, \quad (1.2.9)$$

respectively.

1.2.2 One-dimensional harmonic oscillator

$$\hat{H} = \frac{\hat{p}_x^2}{2m} + \frac{1}{2} m \omega^2 x^2, \quad (1.2.10)$$

where ω is the frequency of the oscillation. The corresponding

Schrödinger equation is

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \frac{1}{2} m \omega^2 x^2 \right) \psi(x) = E \psi(x). \quad (1.2.11)$$

One can prove that the energy levels and the normalized wave function are

$$E_n = \left(n + \frac{1}{2} \right) \hbar \omega \quad (n = 0, 1, 2, \dots), \quad (1.2.12)$$

$$\psi_n(x) = \left(\frac{\alpha}{\pi^{1/2} 2^n n!} \right)^{1/2} e^{-\frac{1}{2} \alpha^2 x^2} H_n(\alpha x), \quad (1.2.13)$$

where $\alpha = \sqrt{m\omega/\hbar}$ and $H_n(\alpha x)$ is the Hermit polynomials

$$\begin{aligned} H_n(\xi) &= (-1)^n e^{\xi^2} \frac{d^n}{d\xi^n} (e^{-\xi^2}) \\ &= (2\xi)^n - n(n-1)(2\xi)^{n-2} \\ &\quad + \frac{n(n-1)(n-2)(n-3)}{2!} (2\xi)^{n-4} \\ &\quad + \dots + (-1)^{\left[\frac{n}{2}\right]} \frac{n!}{\left[\frac{n}{2}\right]!} (2\xi)^{n-2\left[\frac{n}{2}\right]}, \end{aligned} \quad (1.2.14)$$

$$\left[\frac{n}{2} \right] = \begin{cases} n/2 & (n \text{ even}) \\ (n-1)/2 & (n \text{ odd}) \end{cases}. \quad (1.2.15)$$

Here is a list of the first few Hermit polynomials

$$\begin{aligned} H_0(\xi) &= 1, & H_1(\xi) &= 2\xi, \\ H_2(\xi) &= 4\xi^2 - 2, & H_3(\xi) &= 8\xi^3 - 12\xi. \end{aligned} \quad (1.2.16)$$

The energy spectra of linear oscillator can be obtained easily in the occupation representation. Introducing the creation operator \hat{a}^+ and the annihilation operator \hat{a}

$$\begin{aligned} \hat{a}^+ &= \left(\frac{m\omega}{2\hbar} \right)^{1/2} \left(\hat{x} - \frac{i}{m\omega} \hat{p} \right) \\ \hat{a} &= \left(\frac{m\omega}{2\hbar} \right)^{1/2} \left(\hat{x} + \frac{i}{m\omega} \hat{p} \right) \end{aligned}, \quad (1.2.17)$$

we can rewrite Eq. (1.2.10) into

$$\hat{H} = \hbar\omega \left(\hat{a}^+ \hat{a} + \frac{1}{2} \right), \quad (1.2.18)$$

and prove that the operators \hat{a} and \hat{a}^+ satisfy

$$[\hat{a}, \hat{a}^+] = 1. \quad (1.2.19)$$

Defining the operator of particle number \hat{N} as

$$\hat{N} = \hat{a}^+ \hat{a}, \quad (1.2.20)$$

the eigenvectors $|n\rangle$ of operator \hat{N} are

$$\hat{N} |n\rangle = n |n\rangle, \quad (1.2.21)$$

where $n = 0, 1, 2, \dots$ are the occupied numbers of the corresponding states $|0\rangle, |1\rangle, |2\rangle, \dots$ respectively. In occupation representation $\{|n\rangle\}$, we have

$$\begin{aligned} \hat{a}^+ |n\rangle &= \sqrt{n+1} |n+1\rangle \\ \hat{a} |n\rangle &= \sqrt{n} |n-1\rangle. \end{aligned} \quad (1.2.22)$$

Combining Eqs. (1.2.18) and (1.2.21), we get

$$E_n = \left(n + \frac{1}{2} \right) \hbar\omega. \quad (1.2.23)$$

1.2.3 Central potential

The central potential U only depends on the distance r of the particle from the center. Spherical coordinates are best adapted to the problem. The Schrödinger equation is

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + U(r) \right) \psi(r) = E \psi(r). \quad (1.2.24)$$

We express Eq. (1.2.24) in terms of spherical coordinates

$$-\frac{\hbar^2}{2m} \left[\frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} \right] \psi$$

$$+ U(r)\psi = E\psi. \quad (1.2.25)$$

The solution is

$$\psi(r, \theta, \varphi) = R_{nl}(r) Y_{lm}(\theta, \varphi), \quad (1.2.26)$$

where $Y_{lm}(\theta, \varphi)$ is the spherical harmonic functions

$$Y_{lm}(\theta, \varphi) = N_{lm} P_l^{|m|}(\cos \theta) e^{im\varphi}, \quad (1.2.27)$$

and

$$N_{lm} = \sqrt{\frac{(l - |m|)! (2l + 1)}{(l + |m|)! 4\pi}}, \quad (1.2.28)$$

is the normalized constant, and $P_l^{|m|}(\cos \theta)$ is the associated Legendre polynomials. The first few spherical harmonics are listed below:

$$\begin{aligned} Y_{00} &= \frac{1}{\sqrt{4\pi}}, \\ Y_{10} &= \sqrt{\frac{3}{4\pi}} \cos \theta, & Y_{1\pm 1} &= \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\varphi}, \\ Y_{20} &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1), & Y_{2\pm 2} &= \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{\pm i2\varphi}. \end{aligned} \quad (1.2.29)$$

The orthonormalized condition for Y_{lm} is

$$\int_0^{2\pi} \int_0^\pi Y_{lm}^*(\theta, \varphi) Y_{l'm'}(\theta, \varphi) \sin \theta d\theta d\varphi = \delta_{ll'} \delta_{mm'}. \quad (1.2.30)$$

If $U(r)$ is attractive Coulomb potential

$$U(r) = -\frac{Ze^2}{r}, \quad (1.2.31)$$

the radial part R_{nl} of solution Eq. (1.2.26) is

$$R_{nl}(r) = N_{nl} e^{-\xi/2} \xi^l F(-n + l + 1, 2l + 2, \xi), \quad (1.2.32)$$

where