


Microbiology and Biotechnology

CAMBRIDGE
MODULAR
SCIENCES



Microbiology and Biotechnology

Pauline Lowrie and Susan Wells



Published by the Press Syndicate of the University of
Cambridge
The Pitt Building, Trumpington Street, Cambridge CB2
1RP
40 West 20th Street, New York, NY 10011-4211, USA
10 Stamford Road, Oakleigh, Melbourne 3166,
Australia

© University of Cambridge Local Examinations
Syndicate 1994

First published 1994

Printed in Great Britain at the University Press,
Cambridge

A catalogue record for this book is available from the
British Library

ISBN 0 521 42204 3 paperback

Designed and produced by Gecko Ltd, Bicester, Oxon

Notice to teachers

It is illegal to reproduce any part of this work in
material form (including photocopying and electronic
storage) except under the following circumstances:

- (i) where you are abiding by a licence granted to your
school or institution by the Copyright Licensing Agency;
- (ii) where no such licence exists, or where you wish to
exceed the terms of a licence, and you have gained the
written permission of Cambridge University Press;
- (iii) where you are allowed to reproduce without
permission under the provisions of Chapter 3 of the
Copyright, Designs and Patents Act 1988.

This book is one of a series produced to support
individual modules within the Cambridge Modular
Sciences scheme. Teachers should note that written
examinations will be set on the content of each module
as defined in the syllabus. This book is the authors'
interpretation of the module.

Acknowledgements

Many thanks to Sue Kearsey for her meticulous
and thoughtful editing which the authors greatly
appreciate. However, the authors accept full
responsibility for the final content of the book.

The authors would also like to thank June
Withenshaw for her help with the simulated
newspaper on page 48.

The publishers would like to thank Michael
Reiss and Stephen Shaw for their invaluable help
with the text.

Photographs

1.1a, 1.21, NIBSC/Science Photo Library;
1.1b, Barry Dowsett/Science Photo Library;
1.1c, CNRI/Science Photo Library; 1.6, Dr L Caro/
Science Photo Library; 1.17a, Manfred Kage/
Science Photo Library; 1.17b, 1.24, Biophoto
Associates; 1.17c Dr Jeremy Burgess/Science Photo
Library; 1.17d David Scharff/Science Photo
Library; 2.1 Peter Gould; 3.1, Ronald Sheridan/
Ancient Art and Architecture Collection;
4.1, James Holmes/Celltech Ltd/Science Photo
Library; 4.2, Philippe Plailly/Science Photo Library;
4.4, Geoff Tompkinson/Science Photo Library;
5.1a, b, courtesy of Zeneca BioProducts;
5.5, courtesy of Wearne Associates; 5.7a, b,
courtesy of Gist-brocades; 6.1, courtesy of DISTA
Products Ltd; 7.4a, b, courtesy of Novo Nordisk ;
7.5, Sean Sprague/Panos Pictures.

Cover: A coloured TEM of *Salmonella enteritidis*.
The outer membrane, cell contents and hair-like
flagellae can be seen.
(Alfred Pasioka/Science Photo Library)

Contents

1 Microbiology 4

Kingdom Prokaryotae	5
Kingdom Protocista	12
Kingdom Fungi	14
Viruses	19
Retroviruses	23

2 Microbiological techniques 25

Laboratory safety and aseptic technique	25
Nutritional requirements	26
Culture media	27
Inoculating solid and liquid media	30
Growth in bacteria	31
Growth in fungi	34
Factors affecting microbial growth	35
Growth of bacteriophages	37

3 The scope of biotechnology 38

Principles of biotechnology	39
Genetic engineering	41
Gene cloning	42
Implications of biotechnology	43

4 Large-scale production 49

Methods of fermentation	51
Setting up an industrial process	52
Antibiotics	53
The industrial production of antibiotics	53

5 Biotechnology in food production 57

Improving crop plants by genetic engineering	58
Breeding disease-resistant plants	59
Breeding insect-resistant plants	60
Genetically engineered farm animals	61
Bovine somatotrophin (BST)	61
Single cell protein (SCP)	62
Yeast extract production	65
Making yogurt	65
Wine making	66
Beer manufacture	67
Bread manufacture	68
Enzymes in fruit juice production	68
Enzymes and meat tenderisation	69

6 Biotechnology in health care and medicine 71

Biosensors	73
------------	----

7 Biotechnology in manufacturing industry 76

Isolated enzymes	78
Industrial production of enzymes	78
Immobilised enzymes	80
Enzymes in washing powders	81
The production of methane	83
Energy from biomass	84
Microbial mining	86
Oil recovery and oil digestion	87

Answers to self-assessment questions	89
Index	96

Microbiology

By the end of this chapter you should be able to:

- 1 describe the major characteristics of the four groups of microorganisms, the Prokaryotae (Bacteria and Cyanobacteria), Protoctista, Fungi and Viruses;
- 2 describe, as a result of your observations, the structures and asexual reproduction of a named bacterium;
- 3 describe the general structure and life cycle of viruses, including a named bacteriophage and a named retrovirus.

The main purpose of this book is to give you an understanding of some of the many ways in which microbiology and biotechnology affect our lives. It is also important that you should understand why microorganisms are particularly suitable for use in industrial processes. However, before you can properly understand all these things, it is necessary that you learn a little about the structure and function of the different kinds of microorganisms.

Microorganisms (microbes) are so small that they can only be seen individually with a good



quality light microscope. **Microbiology** is the study of microorganisms and can be subdivided into several specialist branches such as **bacteriology** (the study of bacteria), **mycology** (the study of fungi) and **virology** (the study of viruses). There are four major groups of microorganisms: Prokaryotae, Protoctista, Fungi and Viruses.

All prokaryotic organisms are grouped into the kingdom Prokaryotae. Prokaryotic cells do not have a true, membrane-bound nucleus and organelles, nor flagella with the typical 9 + 2 arrangement of microtubules such as are found in the eukaryotes (see *Foundation Biology* in this series). The kingdom Fungi and the kingdom Protoctista are both eukaryotic. Viruses do not fit into a classification of living organisms because they are dependent on other cells for their reproduction.



● **Figure 1.1** Some examples of bacteria.

- a Scanning electron micrograph (SEM) of the rod-shaped Gram negative bacterium *Escherichia coli* ($\times 14\,500$).
- b TEM of *Campylobacter jejuni* which causes food poisoning in humans. The bacterium moves by means of two single terminal flagella ($\times 9\,000$).
- c Transmission EM of *Staphylococcus aureus*. The cell is in the process of cell division with the new cell wall half completed across the centre of the individual ($\times 62\,000$).



Structure	Features
*cell wall	10–100 nm thick, net-like, multilayered structure made of murein (a polymer of polysaccharide and peptide components); two distinct types are distinguished by the Gram's stain: Gram positive ; thicker, rigid murein network filled with peptides and polysaccharides, e.g. <i>Lactobacillus</i> Gram negative : thinner, more complex, made of murein coated with a smooth layer of lipids, e.g. <i>Escherichia coli</i>
cell surface membrane	phospholipid bilayer with proteins floating in membrane; proteins include enzymes involved in respiration, photosynthesis*, nucleic acid synthesis* and electron transport*
*ribosomes	small 70S ribosomes are the site of protein synthesis
*cytoplasm	uniform with few organelles; contains storage granules, ribosomes, plasmids; a nucleoid (nuclear region) is usually present
•flagellum	used in the locomotion of many motile bacteria; a rigid, hollow cylinder of protein, the base of which rotates propelling the cell along, e.g. <i>Rhizobium</i> and <i>Azotobacter</i>
•capsule (slime layer)	an outer protective layer visible with negative (background) staining; it protects against chemicals and desiccation, stores waste products and protects the bacterium from attack by phagocytic cells; it helps bacteria to form colonies and is important in soil bacteria where it helps bind soil particles into crumbs, e.g. <i>Azotobacter</i>
•photosynthetic membranes	sac-like or tubular infoldings of the cell surface membrane provide a large surface area for the inclusion of bacteriochlorophyll and other photosynthetic pigments in photoautotrophic bacteria, e.g. <i>Chromatium</i>
•mesosomes	tightly folded infoldings of the cell surface membrane, thought to be the site of respiration; also involved in cell division and the uptake of DNA
•pili (fimbriae)	numerous projections from the cell surface membrane through the cell wall found all over some bacterial cells; they are used in conjugation to bind cells together and exchange genetic material; they also have an antigenic effect, e.g. <i>Salmonella</i>
•endospores	a hard outer covering forms a resistant endospore which ensures survival in severe conditions of drought, toxic chemicals and extremes of temperature, e.g. <i>Bacillus anthracis</i> spores, which cause the disease anthrax, are known to be viable after 50 years in the soil

● **Table 1.1** Summary of the structure of bacteria (* always present, • sometimes present)

Kingdom Prokaryotae

This kingdom contains all the prokaryotic organisms, that is the Bacteria and Cyanobacteria (the blue-green bacteria, formerly called blue-green algae).

Bacteria


Examples: *Escherichia coli*, *Streptococcus lactis*, *Bacillus subtilis*, *Staphylococcus aureus*

Bacteria are found in a wide range of habitats: in soil, air, water, as well as in or on the surface of animals and plants. They range in size from 0.1–10 µm in length and are usually found in enormous numbers: one gram of soil may contain 100 million bacteria. Some bacteria have an

optimum temperature for growth which is greater than 45 °C and are called thermophilic bacteria; some can even thrive in hot volcanic springs at around 70 °C. Psychrophilic bacteria, on the other hand, grow best at temperatures below 20 °C and can withstand long periods of freezing. Bacteria are important because they help to decay and recycle organic waste. Some cause disease, but most are harmless and many are of increasing economic importance in biotechnology.

Structure of bacteria

Figure 1.1 shows a variety of shapes of bacteria and figure 1.2 shows the structure of a generalised bacterial cell. Table 1.1 gives more detail of their structure.




● **Figure 1.2** A generalised rod-shaped bacterial cell.

The bacterial cell wall


In 1884 Christian Gram developed a way of staining bacteria which divided them into two groups. These were called Gram positive and Gram negative. It is now known that bacteria have two different types of cell wall which the staining technique reveals.

The technique developed by Gram is still commonly used today. It involves heat-fixing a smear of bacteria to a clean microscope slide and then flooding it with crystal violet. All bacteria take up this stain. The smear is washed with Gram's iodine to fix the stain and then decolourised with alcohol or propanone. Gram positive bacteria retain the crystal violet/iodine complex to colour purple, but Gram negative bacteria do not. Finally the smear is counter-stained with a red stain such as safranin or carbol fuchsin. Gram negative bacteria take up this stain and become red. Gram positive bacteria stay purple.

The different reaction to the stain is due to the structure of the two basic types of cell wall (*figure 1.3*). Gram positive bacteria have a cell surface membrane surrounded by a rigid cell wall about 20–80 nm thick. This rigid layer is made of the peptidoglycan **murein**, which has a complex three-dimensional structure. Gram negative bacteria also have a rigid layer outside the surface membrane but it is much thinner, only 2–3 nm thick. On the outside of this is an outer membrane which contains lipopolysaccharides instead of phospholipids. This forms an extra physical barrier to substances, such as antibiotics and enzymes like lysozyme, which normally destroy or inhibit bacteria. The crystal violet/iodine complex is a large molecule and it is thought that, during Gram staining, it becomes trapped inside the cell wall, whereas it is more easily washed out of the thinner Gram negative cell wall.



● **Figure 1.3** The structure of bacterial cell walls based on electron micrographs of the cell walls of *Bacillus* (Gram positive) and *E. coli* (Gram negative).



● **Figure 1.4** The forms (shapes) of bacteria.

Shapes of bacteria


When viewed with a microscope, bacteria show several distinct shapes and these may be used to help in identification (*figure 1.4*).

Reproduction

Bacteria grow very quickly in favourable conditions. The generation time may be as little as 20 minutes, though for many species it is 15–20 hours. Division is usually by binary fission (*figure 1.5*).

Binary fission in *Escherichia coli*

- 1 The circular bacterial chromosome divides but there is no mitotic spindle. The chromosome attaches itself to the cell membrane or, in some cases, to the mesosome.
- 2 A septum starts to be synthesised to divide the cell. This often starts growing where there are mesosomes.
- 3 The septum grows right across the cell, dividing it into two daughter cells.




● **Figure 1.5** Binary fission in bacteria.

Conjugation, transformation and transduction

Some bacteria have 'mating' cells which come together and are joined by their pili in a process known as **conjugation** (figure 1.6). The donor passes a plasmid called the F factor, or fertility factor, to the recipient. The **plasmid** is a small, circular, non-chromosomal piece of DNA. Genetic information on the F factor provides the bacterial cell with everything needed to be a donor, including the capacity to synthesise the sex pilus. The F factor may exist as a free element within the cytoplasm, replicating independently of the bacterial chromosome, or it may become incorporated within the bacterial chromosome and be replicated whenever the chromosome replicates.

Transformation occurs when one bacterium releases DNA which is absorbed by a second bacterium. The second cell therefore acquires new characteristics.

Transduction is where new genes are inserted into the chromosome of a bacterium by a bacteriophage virus (see page 21).



● **Figure 1.6** Conjugation in *E. coli*.

Conjugation, transformation and transduction are not forms of sexual reproduction since fertilisation does not occur. In each of the above cases, DNA has been transferred from a donor to a recipient.

Plasmids

The cytoplasm of certain bacteria contains small circles of DNA called plasmids which are able to replicate independently of the main circular chromosome. Plasmids are known to carry genes which may help the bacterium to survive in adverse conditions. For example, plasmids known as R-factors cause resistance to antibiotics, virus infection and ultraviolet radiation. Plasmids can be transferred to another bacterium in conjugation, transformation or transduction.

The economic importance of bacteria

Some genera of bacteria contain species which are commercially useful and some which are harmful to humans. For example, most *Bacillus* spp. live in the soil, and are either aerobes or facultative anaerobes. *Bacillus subtilis* is a strict aerobe which is used in industry as a source of enzymes such as amylases. *Bacillus thuringiensis* has become important in genetic engineering because it causes a paralytic disease in many caterpillars and has been used to produce insect-resistant plants (page 61). However, some species of *Bacillus* are pathogenic, such as *Bacillus anthracis* which causes the disease anthrax.

Table 1.2 lists some bacteria which are commercially useful. Some of these are described in more detail in later chapters. Other bacteria are useful in different ways, for example in recycling nutrients and fixing nitrogen in ecological cycles. *Rhizobium* is a nitrogen-fixing bacterium which is present in the soil. It invades the root hairs of leguminous plants, causing the cells to divide and form nitrogen-fixing nodules. Many species of *Clostridium* can also fix nitrogen, such as *C. welchii*, *C. pastorianum* and *C. butyricum*, and are important in the nitrogen cycle.

Table 1.3 lists some bacteria which cause disease in humans. Some belong to genera which contain

SAQ 1.1

Match the words on the left to the definitions on the right.

plasmid	slimy layer surrounding some bacterial cell walls
mesosome	circular piece of DNA not joined to chromosome
cell wall	made of pilin, used in conjugation in some species
capsule	the main component of this is the peptidoglycan murein
pilus	infolding of membrane, probably used in cell division

Cyanobacteria (blue-green bacteria)

Examples: *Anabaena cylindris*, *Nostoc muscorum*, *Spirulina platensis*

Cyanobacteria are prokaryotic microorganisms similar to the true bacteria. They are photosynthetic but are not true algae because they do not have membrane-bound nuclei, and are considered to be very ancient life-forms. They have been found in fossil remains from over three billion years ago and may have been some of the first living organisms to evolve on Earth. They are found in the surface layer of fresh and sea water. On land they will grow wherever there is both light and moisture and are found as slime on the surface of mud, rocks, wood and on some living organisms. Their name comes from the photosynthetic pigments which give them a distinct dark greenish-blue colour.

Structure of cyanobacteria

Blue-greens have a typical prokaryotic cell structure since they have a naked coil of DNA and no true nucleus (figure 1.7). The cell wall is similar in structure and composition to that of

Name	Gram stain	Form (shape)	Use
<i>Lactobacillus bulgaricus</i>	+ve	rods	yogurt
<i>Streptomyces thermophilus</i>	+ve	filamentous	yogurt
<i>Streptococcus lactus</i>	+ve	cocci	cheese
<i>Streptococcus cremoris</i>	+ve	cocci	cheese
<i>Methylophilus methylotrophus</i>	variable	cocci	methane, methanol
<i>Clostridium acetobutylicum</i>	+ve	rods	propanone (acetone) and butanol
<i>Leuconostoc mesenteroides</i>	+ve	cocci	dextran
<i>Bacillus subtilis</i> *	+ve	rods	enzymes
<i>Streptomyces</i> spp.	+ve	filamentous	antibiotics
<i>Escherichia coli</i> *	-ve	rods	insulin, growth hormone, interferon
<i>Pseudomonas denitrificans</i>	-ve	rods	vitamin B ₁₂


* using genetic engineering

● **Table 1.2** Some useful bacteria

Name	Gram stain	Form (shape)	Disease
<i>Staphylococcus aureus</i>	+ve	cocci	boils
<i>Salmonella typhimurium</i>	-ve	rods	food poisoning
<i>Salmonella typhi</i>	-ve	rods	typhoid fever
<i>Mycobacterium tuberculosis</i>	variable	fine rods	tuberculosis
<i>Bordetella pertussis</i>	-ve	very short rods	whooping cough
<i>Neisseria gonorrhoea</i>	-ve	cocci	gonorrhoea
<i>Treponema pallidum</i>	variable	long spirals	syphilis
<i>Vibrio cholerae</i>	-ve	curved rods	cholera
<i>Clostridium tetani</i>	+ve	rods	tetanus
<i>Clostridium botulinum</i>	+ve	rods	botulism
<i>Corynebacterium diphtheriae</i>	+ve	short rods	diphtheria
<i>Listeria</i> spp.	+ve	round-ended rods	listeriosis
<i>Shigella sonnei</i>	-ve	rods	dysentery
<i>Yersinia pestis</i>	-ve	small rods	plague

● **Table 1.3** Some harmful bacteria

many non-pathogenic species, for example most species of *Staphylococcus* are facultative anaerobes which are found in the normal microflora of the skin. *S. aureus* normally causes boils. However, one strain of this species has developed resistance to most antibiotics, through overuse of antibiotics in treating patients, and has become a major problem in hospitals in many countries.



● **Figure 1.7** Diagram of a typical blue-green cell (based on electron micrographs).

Gram negative bacteria. Protein synthesis takes place on 70S ribosomes in the cytoplasm. Blue-greens are photosynthetic. They have chlorophyll and carotenoid pigments incorporated into infoldings of the cell surface membrane, called **lamellae**. They also have photosynthetic pigments, such as phycocyanin and phycoerythrin, which are present in phycobilisomes. These give the cells their distinctive colouration. The cells may occur singly or in colonies, but members of a colony remain independent.

Nitrogen fixation

Only a very few organisms are capable of fixing atmospheric nitrogen by reducing it to ammonia and combining it with organic acids to produce amino acids and proteins. Nitrogen-fixing bacteria can do this and so can some blue-greens. Cells able to fix nitrogen contain the enzyme nitrogenase. This enzyme is inactivated by oxygen and so conditions inside the nitrogen-fixing cell have to be


anaerobic. Some blue-greens, such as *Anabaena*, have special thick-walled cells called **heterocysts**. *Anabaena* has filaments made up of many normal photosynthetic cells that produce sugars and oxygen. Scattered along the filaments are a few distinct heterocyst cells that are able to fix atmospheric nitrogen in this way (figure 1.8).

Many filamentous blue-greens are also able to produce **akinetes**, or spores. These are able to survive adverse conditions, such as a period of overpopulation known as an algal 'bloom', and seem to develop from a vegetative cell near to a heterocyst. The cell increases in size and accumulates large food reserves. Photosynthesis within the akinete is reduced and gas vacuoles disappear. This means that the akinete slowly sinks to the bottom of the water. It may survive for several years, and will germinate as soon as conditions become favourable.

The economic importance of blue-greens

Spirulina platensis is a filamentous blue-green found naturally in shallow alkaline lakes in parts of Africa and South America. For thousands of years it has been collected and dried by the local people and used as a food. It is often fried or put in soups and sauces; it is also used as cattle food. *Nostoc* is another blue-green which is used as a food in Peru and in South-East Asia.

In agriculture, nitrogen-fixing blue-greens may be used as organic fertilisers. They are grown on a large scale in China, India and the Philippines, particularly where rice is cultivated in paddy fields. The water may be seeded with a starter culture of blue-greens at the beginning of the growing season. This method has been shown to increase the yield of rice by 15–20%.



● **Figure 1.8** A filament of *Anabaena*.

Research is taking place into the use of blue-greens in a solar energy conversion system. *Anabaena cylindrica* has heterocysts and is able to give off oxygen by photosynthesis in the vegetative cells. In the absence of nitrogen it gives off hydrogen by nitrogenase-catalysed electron

transfer to H^+ ions in the heterocysts. Both oxygen and hydrogen are in demand industrially.

SAQ 1.2

Why are blue-greens classified as prokaryotes?

Phylum	Example	Structure	Locomotion	Nutrition	Reproduction	Importance
Rhizopoda	<i>Amoeba proteus</i>	0.5 mm long, single cell, cytoplasm exists in two states: plasmasol and plasmagel	pseudopodia	phagocytosis and food vacuoles formed	asexual by binary fission	<i>Entamoeba histolytica</i> causes amoebic dysentery
Apicomplexa (sporozoans)	<i>Plasmodium vivax</i>	10 μ m long, different forms during life cycle: sporozoites long and thin, infective particles; merozoites smaller, pear-shaped	microtubules which slide over each other to penetrate host tissue	phagocytosis	asexual by multiple fission and sexual by gametocytes	<i>P. vivax</i> causes parasitic disease malaria, mosquito is the vector
Ciliophora (ciliates)	<i>Paramecium caudatum</i>	100 μ m long, enclosed by a pellicle with cilia, oral groove leads to 'mouth' with cytostome	cilia beat together in coordinated way	cilia beat and sweep food towards cytostome, where phagocytosis takes place	asexual by binary fission, sexual by conjugation	live in fresh water, important in sewage treatment and decomposition
Zoomastigina (flagellates)	<i>Trypanosoma gambiense</i>	unicellular, elongated spindle-shaped, contains trophonucleus (also concerned with nutrition) and kinetoplast near posterior end to control movement	single flagellum bound to the trypanosome along the length of the cell by the undulating membrane, free beyond that; wave-like movements propel the cell through the viscous blood plasma	heterotrophic, parasitic; lives in human blood plasma; soluble nutrients diffuse into cell	longitudinal binary fission in the blood of the host starting at anterior end	causes trypanosomiasis or 'sleeping sickness'; spread by the tsetse fly as it feeds on human blood; symptoms include headache, anaemia, inflammation of brain and spinal cord

● **Table 1.4** A summary of the Kingdom Protocista (continued on page 12)

Phylum	Example	Structure	Locomotion	Nutrition	Reproduction	Importance
Euglenophyta (euglenoid flagellates)	<i>Euglena viridis</i>	300 μm long slender cell, pellicle covers outside, blunt anterior end with reservoir	two flagella arise from base of reservoir, light-sensitive stigma used to orientate movement	chloroplasts present, therefore photoauto- trophic; stores food granules of paramylum, needs vitamins B_1 and B_{12} for growth	longitudinal binary fission (asexual)	found in fresh and salt water, and damp soil
Chlorophyta (green algae)	<i>Chlorella vulgaris</i>	50 μm diameter, unicellular green alga, thin cellulose cell wall, large cup-shaped chloroplast	floats in water	photoauto- trophic	asexual by fission (figure 1.9)	freshwater ponds
Oomycota (oomycetes)	<i>Phytophthora infestans</i> or <i>Pythium</i> spp.	long thread-like hyphae, aseptate (no cross walls), cytoplasm contains many nuclei, cell wall of chitin with mannan and glucan polymers	spore stages have two flagella	heterotrophic (plant parasites), hyphae secrete enzymes, food digested extra- cellularly, soluble nutrients absorbed	asexual by motile spores formed inside a sporangium on a special hypha (sporangiophore); sexual by fusion of male antheridium and female oogonium to produce sexual oospore which survives adverse conditions	<i>Phytophthora</i> causes late blight of potatoes (as in Irish potato famine) <i>Pythium</i> causes damping-off in seedlings

● Table 1.4 (cont.)

SAQ 1.3

It is thought that blue-greens may have been the first photosynthetic organisms on Earth and that they represent a very early stage in the evolution of life. Give as many reasons as you can why this might be so.


Kingdom Protocista

This kingdom has been created to contain all groups of eukaryotic organisms which are neither animals, plants, fungi nor prokaryotes. These


groups are not really related though they do have some similarities. They include all protozoa, all nucleated algae and the slime moulds (table 1.4 and figure 1.9).

The protozoa is a collective term for the phyla Rhizopoda, Zoomastigina, Apicomplexa and Ciliophora. They are found wherever moisture is present, in sea water, fresh water and soil. There are commensal, symbiotic and parasitic species in addition to many free-living types. Protozoa are eukaryotic. The nucleus has a nuclear envelope,


Amoeba




TS leaf infected by *Phytophthora*






Paramecium



Chlorella



Asexual reproduction in *Chlorella*



● **Figure 1.9** The structure of some representative protocists.

and movement is by means of a variety of locomotory structures such as flagella, cilia or pseudopodia. Since the cytoplasm of freshwater protozoans is usually hypertonic to (more concentrated than) the aqueous environment, they take in water by osmosis. To counteract this, they have contractile vacuoles that act as pumps to remove excess water from the cytoplasm. However, contractile vacuoles may also be found in some marine protozoans. All types of nutrition are found in protozoans: some are autotrophic, others are saprotrophic and many are heterotrophic. Digestion of food takes place in food vacuoles in the cytoplasm. Gas exchange is by diffusion across the cell membrane. Waste products from cell metabolism diffuse out of the cell. The main nitrogenous waste is ammonia.

The economic importance of protozoa

Many ciliates are saprobionts and are vital in the recycling of organic wastes, particularly in sewage treatment. Parasitic forms, such as *Entamoeba*, which causes amoebic dysentery, and *Plasmodium*, may cause loss of life. Malaria, for example, can be devastating to the economy of developing countries, incapacitating millions of workers every year.

SAQ 1.4

List **three** features shared by the four phyla classed as protozoa, and **two** features which distinguish them.

Kingdom Fungi


Fungi are obligate aerobes, which are killed in the absence of oxygen, or facultative anaerobes, which grow best when oxygen is available but are able to survive anaerobic conditions. They are found almost everywhere. They are eukaryotic organisms with a cell wall that is usually made of chitin. Fungi are not photosynthetic but heterotrophic, deriving their nutrients by absorption of organic materials.

Fungi are usually filamentous or thread-like. The individual threads are called **hyphae** which are usually multinucleate; these branch profusely, often fusing together to form a tangled mass of branched hyphae called the **mycelium**. Individual hyphae are surrounded by rigid cell walls and grow only at their tips. It is this form of **apical growth** that separates fungi from almost all other organisms, even filamentous ones.


Since fungi are eukaryotic, they have distinct nuclei each surrounded by a nuclear envelope with pores; they have chromosomes and a spindle that appears during nuclear division. Mitochondria are found in the cytoplasm and there is extensive endoplasmic reticulum (figure 1.10); ribosomes are found both free in the cytoplasm and attached to the ER. The cytoplasm also contains numerous vacuoles containing storage materials such as starch, lipid globules and volutin.

The cytoplasm and organelles are surrounded by a selectively permeable, phospholipid unit membrane. The cytoplasm is at its most dense at the tips of the hyphae. The older parts are often metabolically inactive with large vacuoles in the cytoplasm. In septate species there are pores in the cross walls to allow substances in solution, including nuclei, to move freely from one section to another.

The growth of hyphae is very rapid under favourable conditions. Each hypha grows at the tip and branches repeatedly along its




● **Figure 1.10** Diagram of the tip of a typical fungal hypha (based on electron micrographs).




● **Figure 1.11** Conditions required for fungal growth.


Growth of sporangiophore



Development of sporangium



Spore release and dispersal



length to reach new food supplies and grow away from its own waste products (*figure 1.11*).

Phylum Zygomycota (zygomycetes)

Example: *Mucor hiemalis*

In zygomycetes the hyphae are aseptate and make a large, well-developed branching mycelium. Asexual reproduction is by non-motile spores formed in a **sporangium** borne on a **sporangiophore** (*figure 1.12*). Sexual reproduction is shown in *figure 1.13*.


Phylum Ascomycota (ascomycetes)

Examples: *Penicillium notatum*, *Saccharomyces cerevisiae*, *Neurospora* spp.


All ascomycetes have aseptate hyphae. In asexual reproduction, non-motile spores, called **conidia**, are formed on special hyphal branches. Many zygotes form inside the perithecium and divide by mitosis and meiosis to form an **ascus** containing eight ascospores (four of each mating type). The ascospores germinate to form haploid mycelia when conditions are favourable.

Yeasts, such as *Saccharomyces cerevisiae*, are ascomycetes and are industrially important fungi. They have been used for centuries in the making of bread, the brewing of beer and in winemaking.

● **Figure 1.12** Asexual reproduction in *Mucor*.



● **Figure 1.13** Sexual reproduction in *Mucor*: conjugation.



● **Figure 1.14** Sexual reproduction in yeast *Saccharomyces cerevisiae*.