

JACQUES C. J. NIHOUL (editor)

---

# MARINE TURBULENCE

83  
J  

---

ELSEVIER OCEANOGRAPHY SERIES 28

*Elsevier Oceanography Series, 28*

# Marine Turbulence

PROCEEDINGS OF THE 11th INTERNATIONAL LIÈGE COLLOQUIUM  
ON OCEAN HYDRODYNAMICS

Edited by

JACQUES C.J. NIHOUL

*Professor of Ocean Hydrodynamics,  
University of Liège,  
Liège, Belgium*



ELSEVIER SCIENTIFIC PUBLISHING COMPANY

Amsterdam — Oxford — New York

1980

ELSEVIER SCIENTIFIC PUBLISHING COMPANY  
335 Jan van Galenstraat  
P.O. Box 211, 1000 AE Amsterdam, The Netherlands

*Distributors for the United States and Canada:*

ELSEVIER/NORTH-HOLLAND INC.  
52, Vanderbilt Avenue  
New York, N.Y. 10017

ISBN 0-444-41881-4 (Vol. 28)  
ISBN 0-444-41623-4 (series)

© Elsevier Scientific Publishing Company, 1980

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands

Printed in The Netherlands

# **Marine Turbulence**

## **FURTHER TITLES IN THIS SERIES**

- 1 J.L. MERO  
THE MINERAL RESOURCES OF THE SEA
- 2 L.M. FOMIN  
THE DYNAMIC METHOD IN OCEANOGRAPHY
- 3 E.J.F. WOOD  
MICROBIOLOGY OF OCEANS AND ESTUARIES
- 4 G. NEUMANN  
OCEAN CURRENTS
- 5 N.G. JERLOV  
OPTICAL OCEANOGRAPHY
- 6 V. VACQUIER  
GEOMAGNETISM IN MARINE GEOLOGY
- 7 W.J. WALLACE  
THE DEVELOPMENT OF THE CHLORINITY/SALINITY CONCEPT  
IN OCEANOGRAPHY
- 8 E. LISITZIN  
SEA-LEVEL CHANGES
- 9 R.H. PARKER  
THE STUDY OF BENTHIC COMMUNITIES
- 10 J.C.J. NIHOUL (Editor)  
MODELLING OF MARINE SYSTEMS
- 11 O.I. MAMAYEV  
TEMPERATURE-SALINITY ANALYSIS OF WORLD OCEAN WATERS
- 12 E.J. FERGUSON WOOD and R.E. JOHANNES  
TROPICAL MARINE POLLUTION
- 13 E. STEEMANN NIELSEN  
MARINE PHOTOSYNTHESIS
- 14 N.G. JERLOV  
MARINE OPTICS
- 15 G.P. GLASBY  
MARINE MANGANESE DEPOSITS
- 16 V.M. KAMENKOVICH  
FUNDAMENTALS OF OCEAN DYNAMICS
- 17 R.A. GEYER  
SUBMERSIBLES AND THEIR USE IN OCEANOGRAPHY AND OCEAN  
ENGINEERING
- 18 J.W. CARUTHERS  
FUNDAMENTALS OF MARINE ACOUSTICS
- 19 J.C.J. NIHOUL (Editor)  
BOTTOM TURBULENCE
- 20 P.H. LEBLOND and L.A. MYSAK  
WAVES IN THE OCEAN
- 21 C.C. VON DER BORCH (Editor)  
SYNTHESIS OF DEEP-SEA DRILLING RESULTS IN THE INDIAN OCEAN
- 22 P. DEHLINGER  
MARINE GRAVITY
- 23 J.C.J. NIHOUL (Editor)  
HYDRODYNAMICS OF ESTUARIES AND FJORDS
- 24 F.T. BANNER, M.B. COLLINS and K.S. MASSIE (Editors)  
THE NORTH-WEST EUROPEAN SHELF SEAS: THE SEA BED AND THE  
SEA IN MOTION
- 25 J.C.J. NIHOUL (Editor)  
MARINE HYDRODYNAMICS
- 26 H.-G. RAMMING and Z. KOWALIK  
NUMERICAL MODELLING OF MARINE HYDRODYNAMICS
- 27 R.A. GEYER (Editor)  
MARINE ENVIRONMENTAL POLLUTION

## FOREWORD

The concepts of mixing of water masses, dispersion of anomalies, transport of momentum and vorticity and the dissipation of kinetic energy by turbulent motions in the ocean, play a central role in almost every aspect of Oceanography. The action of turbulence appears not only in the circulation models of physical oceanographers but equally in the models of chemical distributions, of biological production and of sedimentation.

The interaction between atmosphere and ocean - with its important consequences for climate variation - can only be modelled by taking into account the transport of heat, salt and momentum by turbulent motions in the wind-mixed surface layer and the underlying thermocline.

The major advances that have been made in the past decade have revolutionized our thinking about turbulence in the ocean. The bulk effects of turbulence that were known fifty years ago can now for the first time be discussed in terms of models of the responsible flow patterns based on an understanding of the underlying dynamics and physics.

The IAPSO General Assembly in Grenoble (1976) accepted a proposal submitted by the British National Committee to sponsor a second Symposium on Turbulence in the Ocean, at which the emphasis would be on seeking a synthesis across the whole spectral band from millimetres to megametres. This proposal was subsequently discussed by SCOR which agreed to co-sponsor the meeting and by the IOC which offered support. A number of member states recommended that the Symposium should be recognised as a IDOE meeting. It was therefore designated the IAPSO-SCOR-IDOE Second Symposium on "Turbulence in the Ocean", with support from the IOC.

It was decided to make the first week an open meeting, during which both invited and contributed papers could be presented and discussed in open sessions to be attended by all comers. The plan was to follow this by a closed session during the second week. In order to accommodate the idea of an open first week, it was agreed that this would be combined with the annual Colloquium on Ocean Hydrodynamics held in Liège University.\*

\*J.D. Woods : Report to the IOC.

The international Liège Colloquia on Ocean Hydrodynamics are organized annually. Their topics differ from one year to another and try to address, as much as possible, recent problems and incentive new subjects in physical oceanography.

Assembling a group of active and eminent scientists from different countries and often different disciplines, they provide a forum for discussion and foster a mutually beneficial exchange of information opening on to a survey of major recent discoveries, essential mechanisms, impelling question-marks and valuable suggestions for future research.

The papers presented at the Eleventh International Liège Colloquium on Ocean Hydrodynamics report theoretical and experimental research and they address such different scales of motions as synoptic eddies, fronts, mesoscale eddies, and three-dimensional microscale fluctuations. Their unity resides in a common approach to the variability of the seas, based on the profound understanding of non-linear processes which the theory of turbulence provides.

Jacques C.J. NIHOUL.

The Scientific Organizing Committee of the Eleventh International Liège Colloquium on Ocean Hydrodynamics and all the participants wish to express their gratitude to the Belgian Minister of Education, the National Science Foundation of Belgium, The University of Liège, the Intergovernmental Oceanographic Commission and the Division of Marine Sciences (UNESCO) and the Office of Naval Research for their most valuable support.

## LIST OF PARTICIPANTS

ADAM, Y., Dr., Ministère de la Santé Publique et Environnement, Belgium.

BAH, A., Prof., Dr., Institut Polytechnique de Conakry, Rép. de Guinée.

BELYAEV, V.S., Prof., Dr., Institute of Oceanology, Moscow, U.S.S.R.

BERGER, A., Prof., Dr., Université Catholique de Louvain, Belgium.

BOWDEN, K.F., Prof., Dr., University of Liverpool, U.K.

BURT, W.V., Prof., Dr., Office of Naval Research, London, U.K.

BUTTI, C.H., Dr., Rijkswaterstaat, 's Gravenhage, The Netherlands.

CALDWELL, D.R., Prof., Dr., Oregon State University, U.S.A.

CHABERT d'HIERES, G., Ir., Institut de Mécanique de Grenoble, France.

COANTIC, M., Prof., Dr., Institut de Mécanique Statistique de la  
Turbulence, Marseille, France.

COLIN DE VERDIERE, A., Dr., C.O.B., Brest, France.

DELCOURT, B., Dr., Von Karman Institute for Fluid Dynamics,  
Rhode St. Genèse, Belgium.

DERENNE, M., Mr., Université de Liège, Belgium.

DESAUBIES, Y.J.F., Dr., University of Washington, Seattle, U.S.A.

DILLON, T.M., Ir., Oregon State University, U.S.A.

DISTECHE, A., Prof., Dr., Université de Liège, Belgium.

FEDOROV, K., Prof., Dr., Institute of Oceanology, Moscow, U.S.S.R.

FERGUSON, S.R., Mr., University of Liverpool, U.K.

GIBSON, C.H., Prof., Dr., University of California, U.S.A.

GOODMAN, L., Dr., O.N.R., Washington, U.S.A.

GORDON, C.M., Dr., Naval Research Laboratory, Washington, U.S.A.

GREGG, M.C., Prof., Dr., University of Washington, Seattle, U.S.A.

HAUGUEL, A., Ir., Electricité de France, Chatou, France.

HOPFINGER, E., Dr., Institut de Mécanique de Grenoble, France.

JOSSERAND, M., Ir., Institut de Mécanique de Grenoble, France.

KITAIGORODSKII, S.A., Prof., Dr., University of Copenhagen, Denmark.

KRUSEMAN, P., Dr., K.N.M.I., De Bilt, The Netherlands.

KULLENBERG, G.E.B., Prof., Dr., University of Copenhagen, Denmark.

LEBON, G., Prof., Dr., Université de Liège, Belgium.

LE PROVOST, C., Prof., Dr., Institut de Mécanique de Grenoble, France.

LEWALLE, A., Ir., Université de Liège, Belgium.

LOFFET, A., Ir., Université de Liège, Belgium.

LUMLEY, J.L., Prof., Dr., Cornell University, Ithaca, U.S.A.

MAHRT, K.H., Dr., Universität Kiel, W. Germany.

MICHAUX, T., Ir., Université de Liège, Belgium.  
MITCHELL, J.B., Dr., U.K. Meteorological Office, Bracknell, U.K.  
MULLER, P., Dr., Harvard University, Cambridge, U.S.A.  
NIHOUL, J.C.J., Prof., Dr., Université de Liège, Belgium.  
NIZET, J.L., Ir., Université de Liège, Belgium.  
OAKLEY, N.S., Dr., Bedford Institute of Oceanography, Dartmouth,  
Canada.  
OLBERS, D., Dr., Universität Kiel, W. Germany.  
ORLANSKI, I., Dr., Princeton University, U.S.A.  
OSBORN, T.R., Prof., Dr., University of British Columbia, Vancouver,  
Canada.  
OSTAPOFF, F., Dr., Sea-Air Interaction Laboratory, Miami, U.S.A.  
OZER, J., Ir., Université de Liège, Belgium.  
OZMIDOV, R.V., Prof., Dr., Institute of Oceanology, Moscow, U.S.S.R.  
PANCHEV, S., Prof., Dr., University of Sofia, Bulgaria.  
PASMANTER, R., Dr., Rijkswaterstaat, Den Haag, The Netherlands.  
PIACSEK, S.A., Dr., NORDA, Bay St. Louis, U.S.A.  
PICHOT, G., Ir., Ministère de la Santé Publique et de l'Environnement,  
Belgium.  
RENIER, I., Miss, Université de Liège, Belgium.  
REVAULT d'ALLONNES, M., Dr., Museum d'Histoire Naturelle, Paris, France.  
RHINES, P.B., Dr., Woods Hole Oceanographic Institution, Woods Hole,  
U.S.A.  
ROISIN, B., Ir., Université de Liège and Florida State University,  
U.S.A.  
RONDAY, F.C., Dr., Université de Liège, Belgium.  
RUNFOLA, Y., Ir., Université de Liège, Belgium.  
SMITZ, J., Ir., Université de Liège, Belgium.  
STROSCIO, M.A., Dr., Johns Hopkins University, U.S.A.  
SWIFT, M.R., Prof., Dr., University of New Hampshire, U.S.A.  
THOMASSET, F., Ir., IRIA Laboria, Le Chesnay, France.  
TURNER, J.S., Prof., Dr., Australian National University, Canberra,  
Australia.  
VETH, C., Dr., N.I.O.Z., Texel, The Netherlands.  
WOODS, J.D., Prof., Dr., Universität Kiel, W. Germany.  
WOLF, J., Mrs, I.O.S., Bidston, U.K.

CONTENTS

|   |     |
|---|-----|
| FOREWORD . . . . .  | V   |
| ACKNOWLEDGMENTS . . . . .   | VII |
| LIST OF PARTICIPANTS . . . . .  | IX  |
| J.C.J. NIHOUL : The turbulent ocean . . . . .   | 1   |
| A. COLIN DE VERDIERE : The influence of meridional boundaries<br>upon Rossby wave rectification processes . . . . .       | 21  |
| S. PANCHEV : Spectral structure of horizontal oceanic<br>turbulence - Semi-empirical models . . . . .                     | 43  |
| K.N. FEDOROV : Intrusive fine structure in frontal zones and<br>indication of double diffusion . . . . .                  | 57  |
| I. ORLANSKI and C.P. CERASOLI : Resonant and non-resonant wave-<br>wave interactions for internal gravity waves . . . . . | 65  |
| T.M. DILLON and D.R. CALDWELL : Kinetic energy dissipation<br>observed in the upper ocean . . . . .                       | 101 |
| M. REVAULT D'ALLONNES and G. CAULLIEZ : Observations of air-<br>sea interaction parameters in the open ocean . . . . .    | 103 |
| N.S. OAKEY and J.A. ELLIOTT : Dissipation in the mixed layer<br>near Emerald Basin . . . . .                              | 123 |
| M.C. GREGG : Zero crossings of temperature microstructure .   | 135 |
| T.R. OSBORN : Dissipation measurements of oceanic turbulence  | 143 |
| V.S. BELYAEV : Shear instability generation of the<br>turbulence in the ocean (field evidences) . . . . .                 | 157 |
| R.V. OZMIDOV : Small-scale oceanic turbulence . . . . .   | 167 |
| J.L. LUMLEY : Some fundamental aspects of turbulence with<br>implications in geophysical flows . . . . .                  | 175 |
| S.A. PIACSEK and J. TOOMRE : Nonlinear evolution and<br>structure of salt fingers . . . . .                               | 193 |
| C.H. GIBSON : Fossil temperature, salinity, and vorticity<br>turbulence in the ocean . . . . .                            | 221 |

## XII

|  |     |
|--|-----|
| K.F. BOWDEN and S.R. FERGUSON : Variations with height of<br>the turbulence in a tidally-induced bottom boundary layer | 259 |
| M.R. SWIFT : Spatially varying turbulence production<br>in tidal channels . . . . .                                    | 287 |
| C. VETH : A laser-Doppler velocimeter for small scale<br>turbulence studies in the sea . . . . .                       | 303 |
| J. WOLF : Estimation of shearing stresses in a tidal current<br>with application to the Irish Sea . . . . .            | 319 |
| J.C.J. NIHOUL, Y. RUNFOLA and B. ROISIN : Shear effect<br>dispersion in a shallow tidal sea . . . . .                  | 345 |
| A. WARLUZEL and J.P. BENQUE : Dispersion in a tidal sea . .  | 363 |
| Subject Index . . . . .  | 375 |

## THE TURBULENT OCEAN

Jacques C.J. NIHOUL<sup>1</sup>

Mécanique des Fluides Géophysiques, Université de Liège, Sart Tilman  
B6, B-4000 Liège, (Belgium).

<sup>1</sup>Also at the Institut d'Astronomie et de Géophysique, Université de  
Louvain (Belgium).

### ABSTRACT

The variability of the ocean over a wide range of scales, from the megameter to the millimeter, is examined in the light of turbulence theory.

The geophysical constraints which arise from the Earth's rotation and curvature and from the stratification are discussed with emphasis on the role they can play at different scales in inducing instabilities and a transfer of energy to other scales of motion.

### INTRODUCTION

Turbulence in the ocean is still a very controversial subject. An innocent physicist, emboldened by a solid background in the theory of turbulence, who would approach the subject relying on his good understanding of, say, turbulent channel flow, might soon find himself confronted with a maze of conflicting experimental data and a nightmarish farrago of theories where he recognizes very little of what he regards as "turbulence".

Assuming that he is able to muster enough of his high school latin and greek to find his way among iso-halines, iso-pycnals and other proliferating peculiar surfaces which seem to fill the ocean with complexity, he may still require some time to adjust to new cabalistic concepts such as enstrophy, red energy cascade, eddies "which do not overturn", double diffusion or fossil turbulence.

Used to regard a turbulent flow as the superposition of a mean motion and turbulent fluctuations, he must face the fact that, if such things exist in the ocean, they are several thousands orders of magnitude apart in scales and apparently separated by a jungle of

complicated movements which one refers to as the "variability of the ocean".

One of the most intriguing aspect of ocean variability is that, while the macroscale dynamics is claimed to be governed by a cascade of enstrophy to smaller scales, studies of mesoscale and microscale variability seem totally unconcerned with it as if it had disappeared somewhere on the way. (Planetary oceanographers talk about enstrophy "dissipation" but surely they mean something else like "annihilation"; even if one of them, answering a question at the Eleventh Liège Colloquium on Ocean Hydrodynamics, expressed the somewhat surprising view that "dissipated" enstrophy turns into heat).

For somebody trained in classical turbulence theory, it is not immediately obvious that the variability of the ocean is a form of turbulence.

If it is turbulence, it is clearly very different from the type of turbulence with which mechanical engineers, say, are familiar.

The reason why it should be so is found in the work of those oceanographers who, taking a quite opposite view, insist on describing ocean hydrodynamics in terms of non-turbulent theories such as linear wave propagation or molecular effects.

That they succeed in explaining some of the observations - after removing turbulence from the experimental signals in an operation which they gallantly call "decontamination" - is an indication of the mechanisms which are particular to geophysical flows and which are liable to modify, more or less drastically, geophysical turbulence.

The Coriolis force and the density stratification - which allow wave motions that do not exist in non-rotating non-stratified fluids -, the variations of density not only with temperature but also with salinity, - with different molecular diffusivities for momentum, heat and salt -, and their interference in other processes related to bottom or coastal topography and air-sea interactions, provide a great variety of mechanisms which eventually combine with mechanical effects to determine the stability or the instability of oceanic motions and the subsequent constraints on turbulence at different scales.

Including such geophysical constraints, it becomes possible to understand some essential characteristics of ocean variability and to build an image of what turbulence in the ocean may be.

## TIME SCALES, LENGTH SCALES AND LINEAR WAVE THEORY

The equations of Geophysical Fluid Dynamics admit linear wave solutions of different kinds (e.g. Monin et al, 1977). Whether these waves can be observed in the ocean depends on a series of factors. Very small amplitude waves will not be affected by non-linear interactions but, on the other hand, they may be masked by stronger motions and remain unnoticed. Interactions between larger amplitude waves may create an intricate field of waves and wave packets of all scales, wave breaking and turbulence, from which the ideal individual wave of linear theory cannot be sorted out.

The theory of linear waves is however always a useful mathematical exercise as it helps to identify the dominant length scales (wave numbers) and time scales (frequencies) of motions.

From this point of view, it is convenient to divide ocean waves into three categories (Nihoul, 1979) :

### A. Macroscale waves

These waves have frequencies ( $\omega$ ) in the range

$$10^{-8} \text{ s}^{-1} \lesssim \omega \lesssim 10^{-5} \text{ s}^{-1} \quad (1)$$

and operate over a range of horizontal wave-numbers ( $k$ )

$$10^{-6} \text{ m}^{-1} \lesssim k \lesssim 10^{-3} \text{ m}^{-1} \quad (2)$$

(Waves of larger periods could be considered but the limiting frequency  $10^{-8}$  corresponding to periods larger than 10 years, they are presumably rather irrelevant to the present discussion). Macroscale waves are directly related to the spatial variations of the Coriolis parameter  $f$  ( $f = 2 \Omega \sin \phi$  where  $\Omega$  is the angular velocity of the earth's rotation and  $\phi$  the latitude) i.e. to the parameter

$$\beta = \|\nabla f\| \sim 10^{-11} \text{ m}^{-1} \text{ s}^{-1} \quad (3)$$

Macroscale waves include (e.g. Rhines, 1977 ; Nihoul, 1979)

(i) very slow baroclinic waves ( $\omega k \ll \beta$ ) for which

$$\omega \sim \frac{N^2 \beta H^2}{f^2} k \quad (4)$$

where  $H$  is the depth of the ocean and  $N$  the Brunt-Väisälä frequency.

(The condition  $\omega k \ll \beta$  yields  $L \sim k^{-1} \gg R$  where  $R = NHf^{-1} \sim 10^5$  m is the so-called "Rossby internal scale". These waves are thus very slow large scale small amplitude waves which, if excited, are likely to break rapidly under the effect of bottom slope and general baroclinic instability).

(ii) barotropic Rossby waves ( $\omega k \sim \beta$ ) for which

$$\omega \sim \beta k^{-1} \sim f \gamma H^{-1} k^{-1} \quad (5)$$

where  $\gamma$  is the non-dimensional mean bottom slope chosen here of the order  $\gamma \sim 10^{-4}$  such that  $f \gamma H^{-1} \sim \beta \sim 10^{-11}$ .

(In a forced problem, with wind blowing across the ocean surface, these waves constitute the most important mode at "weatherlike" time scales over horizontal scales greater than  $R$ ).

(iii) fast baroclinic waves confined within a layer of thickness  $f L N^{-1}$  above the sloping bottom, for which

$$\omega \sim \gamma N \quad (6)$$

(If, in the dispersion relation of topographic Rossby waves  $\omega \sim f \gamma H^{-1} k^{-1}$ , one replaces the depth  $H$  by the "penetration height"  $f L N^{-1}$ , one obtains the dispersion relation (6). These waves may thus be regarded as topographic Rossby waves where density stratification provides a lid for vortex stretching).

#### B. Mesoscale waves

These waves have frequencies in the range

$$10^{-5} \text{ s}^{-1} \lesssim \omega \lesssim 10^{-2} \text{ s}^{-1} \quad (7)$$

and include gyroscopic waves and inertial oscillations, tides, and internal gravity waves (e.g. Tolstoy, 1963 ; Monin et al, 1977).

The effect of the Earth's curvature becomes here negligible and the essential factors in the dispersion relations are the Coriolis parameter  $f$  and the Brunt-Väisälä frequencies  $N_{\min}$  and  $N_{\max}$ .

#### C. Microscale waves

These waves have frequencies

$$10^{-2} \text{ s}^{-1} \lesssim \omega \quad (8)$$

They are essentially surface waves and acoustic waves. The former only affect the upper layer of the ocean and may be regarded as an

indispensable - unfortunately rather complicated - way of transferring momentum and energy directly from the wind to the sea ; the latter are marginally important in Ocean Hydrodynamics from which they are customarily excluded by the Boussinesq approximation.

#### MACROSCALE TURBULENCE

Macroscale motions in the ocean include large scale currents (gyres) and quasi-geostrophic or "synoptic" eddies which appear, from observational studies, to contain a large fraction of the ocean's kinetic energy.

The dynamics of the synoptic eddies is dominated by the earth's curvature - parameterized in terms of  $\beta$  - and their horizontal length scale is of the order of the Rossby internal scale  $R$ .

The spectral characteristics of the synoptic eddies, wave number  $\kappa_\beta$ , frequency  $\omega_\beta$  and energy level  $\kappa_\beta E(\kappa_\beta)$  where  $E(\kappa)$  is the horizontal kinetic energy spectral density, can be estimated by turbulence similarity arguments. One finds

$$\kappa_\beta \sim R^{-1} \sim 10^{-5} \text{ m}^{-1} \quad (9)$$

$$\omega_\beta \sim \beta \kappa_\beta^{-1} \sim 10^{-6} \text{ s}^{-1} \quad (10)$$

$$\kappa_\beta E(\kappa_\beta) \sim \beta^2 \kappa_\beta^{-4} \sim 10^{-2} \text{ m}^2 \text{ s}^{-2} \quad (11)$$

These estimates appear to be in good agreement with the observations (e.g. Koshlyakov and Monin, 1978).

The use of the term "synoptic" emphasizes the physical analogy between these eddies and the synoptic eddies of the atmosphere (cyclones and anticyclones, quasi-geostrophic motions at the Rossby scale).

The synoptic variability of the atmosphere, however, has time scales of the order of a week and is shaped by pressure lows and highs with characteristic horizontal scales of the order of the thousand of kilometers and one must exclude the hypothesis of the generation of synoptic ocean eddies by direct resonant interactions. Atmospheric disturbances - lows and highs - generate large-scale currents in the ocean and it is the barotropic and essentially baroclinic instability of these currents which provide the energy for the synoptic eddies (Koshlyakov and Monin, 1978).

(The kinetic - and approximately equal potential-energy of the synoptic eddies is essentially higher than the kinetic energy of the