

Nonlinear Waves, Solitons and Chaos

2nd Edition

Eryk Infeld & George Rowlands



Nonlinear Waves, Solitons and Chaos

Eryk Infeld

Institute for Nuclear Studies, Warsaw

George Rowlands

Department of Physics, University of Warwick

2nd edition



CAMBRIDGE
UNIVERSITY PRESS

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK <http://www.cup.cam.ac.uk>
40 West 20th Street, New York, NY 10011-4211, USA <http://www.cup.org>
10 Stamford Road, Oakleigh, Melbourne 3166, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain

First edition © Cambridge University Press 1990
Second edition © Infeld and Rowlands 2000

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 1990
Reprinted (with corrections) 1992
Second edition 2000

Printed in the United Kingdom at the University Press, Cambridge

Typeset in Times 9.5 13pt [vN]

A catalogue record for this book is available from the British Library

Library of Congress Cataloguing in Publication data

Infeld, E. (Eryk)
Nonlinear waves, solitons, and chaos / E. Infeld, G. Rowlands. 2nd ed.
p. cm.

Includes bibliographical references and index.

ISBN 0 521 63212 9 (hb)

1. Solitons. 2. Chaotic behavior in systems. 3. Nonlinear waves. I. Rowlands, G. (George) II. Title.
QC174.26.W28.I55 2000 530.14 dc21 99-045441

ISBN 0 521 63212 9 hardback
ISBN 0 521 63557 8 paperback

Nonlinear Waves, Solitons and Chaos

This revised and updated second edition of a highly successful book is the only text at this level to embrace a universal approach to three major developments in classical physics; namely nonlinear waves, solitons and chaos. The authors now include new material on biology and laser theory, and go on to discuss important recent developments such as soliton metamorphosis.

A comprehensive treatment of basic plasma and fluid configurations and instabilities is followed by a study of the relevant nonlinear structures. Examples of these are coherent entities like nonlinear waves and solitons, as well as the incoherent structures associated with chaos. The first part of the book is a self-contained introduction to general topics associated with nonlinear graduate physics, and would be accessible to final-year undergraduates and beginning graduate students. The remainder of the book, for example the treatment of cylindrical solitons, is more advanced and will have a wide appeal to specialists in a number of branches of physics. Each chapter concludes with a set of problems.

This text will be particularly valuable for students taking courses in nonlinear aspects of physics. In general, it will be of value to final-year undergraduates and beginning graduate students studying fluid dynamics, plasma physics or applied mathematics.

ERYK INFELD was born in Toronto in 1940. In 1950 he left Canada for Poland, as his father was Polish (the scientist Leopole Infeld, best known for his work with Einstein). He graduated from both Warsaw University (physics) and Cambridge University (mathematics). He is a professor of nonlinear dynamics in Warsaw and has visited Cambridge University, University College London, Culham Laboratories, Warwick University (Royal Society Visitor) and the University of Montreal (visiting professor). His publications include 120 papers and two CUP books. The first edition of the present book was designated Book of the Month by the Newbridge Book Club. He served as president of the Polish Electromagnetic Society and as a member of the Polish Academy Physics Committee. In 1990 he was awarded the prize of the Polish Academy. His main interests are nonlinear dynamics, plasma theory and hydrodynamics. He has worked with George Rowlands, J.M.T. Thompson, FRS, and about twenty other scientists.

GEORGE ROWLANDS graduated from Leeds with a PhD in physics in 1956. After post-doctoral work at the NRC in Ottawa Canada he was a Harwell Fellow for three years. He then worked for a six-year period as a Senior Scientific Officer studying magnetic fusion with the UKAEA at Culham, UK. In 1966 he joined the then new University of Warwick and has been there ever since, the last 10 years as Professor of Theoretical Physics. During his time at Warwick Prof. Rowlands has spent time abroad as Visiting Professor at (amongst others) the Lawrence Livermore National Laboratory the University of Wisconsin and NASA. For 30 years he has collaborated with fellow author Eryk Infeld from the University of Warsaw, and together they have published at least 40 research papers on nonlinear plasma physics. In recognition for their work on plasma physics they were jointly awarded a prize by the Polish Academy. Prof. Rowlands has co-authored three software packages on chaos and time series analysis with Prof. J.C. Sprott from the University of Wisconsin, which are published by the American Institute of Physics. He is also author of *Non-Linear Phenomena in Science and Engineering* (Ellis Horwood Ltd, 1990) which is a textbook for final-year undergraduates and beginning graduate students.

Foreword to the first edition

The last few decades have seen three important developments in nonlinear classical physics, all of which extend across the board of physical disciplines. They have, however, received uneven coverage in the literature.

Perhaps the best known outburst of activity is associated with the soliton, and the most famous development here is the inverse scattering method which has been with us now for over twenty years. There are, however, several other, less known methods for treating solitons. Indeed these compact, single hump wave entities have been known to scientists for over a century and a half (it might be interesting to look through some old ships' log books!). Nevertheless, books on the subject tend to concentrate on the inverse scattering method.

The second much publicized development is a new understanding of some deterministic aspects of chaos as well as the various roads a physical system can take to reach a chaotic state. Established views are being revised and new concepts and indeed even universal constants are being found. These important new developments derive from a realization that complex chaotic behaviour can be described by simple equations. The field has now reached the stage where a summary of basic theory can be given, though applications to specific physical problems are largely at the research stage.

The third development is somewhat less well publicized. Over the last three decades or so, scientists working on fluid dynamics and plasma and solid state theory have developed a multitude of new methods to deal with nonlinear waves. Some of these people were aware of the shortcomings of our linear physics education even before the above-mentioned two developments brought them to the attention of the scientific community.

We believe that, although there are now a number of books on all three topics, the time has come to try to bring them together in one volume. Thus the present book documents the three important developments in classical physics jointly, and, when possible, points out the similarities of approach.

The authors' research interest over the past twenty years has been in fluid dynamics and plasma theory and this is reflected in the book. However, the main aim is to cover a wider range of nonlinear wave phenomena than hitherto. A few examples of what is done are: treatment of both surface and volume wave phenomena, including recent results (e.g. instabilities and their pictorial representation, wavelength doubling, wave dynamics in three dimensions, splitting of signals observed experimentally, the universal theory of wave envelope dynamics); new developments in

soliton studies (e.g. many soliton experiments in rectangular, cylindrical and spherical devices and their theory); and a bringing together of theoretical and numerical results on various scenarios for reaching chaos. An example of what is not attempted is a coverage (or even mention) of the 100 or so instabilities found in plasmas and fluids. Instead we present the basic physics of a few of them, each representing a whole category in some sense. Thus, all in all, the ambition of the book could be summarized by the adage 'not many but much'. Some unsolved problems are indicated. References are extensive and exercises are given at the end of each chapter. Thus the more ambitious reader should be able to get into the field. On the other hand, little knowledge is assumed, thus also giving the general science graduate (or senior undergraduate), who would like to learn what these new developments are about, a chance to do so.

As one of us is based in Warsaw, an attempt has been made to do some justice to research performed in Poland and the Soviet Union.

The authors would like to thank Drs P. Frycz, P. Goldstein, T. Lenkowska, K. Murawski and L. Starkman for critical reading of the manuscript, and Professors P.N. Butcher*, J.P. Dougherty, A. Kuszell*, E. Kuznetsov, R. Raczka*, A.A. Skorupski, K.N. Stepanov and R. Zelazny for remarks on parts of the text. Additional thanks are due to Dr Frycz for preparation the material used in Sections 7.10 and A.1 and Figs. 9.2 and 9.3. We would also like to acknowledge a huge debt to Ms H. Gilder for typing the manuscript several times over. Finally, Cambridge University Press in the person of Dr S. Capelin has been both helpful and patient.

E. Infeld and G. Rowlands
Warwick

* Now deceased.

Foreword to the second edition

When this book was first published in 1990, it became more popular than we expected. A book club chose it as its Book of the Month. It was reprinted in 1992. Eight years have now gone by and we feel it is time for a proper revision. New results and references have been added. On the other hand, some chapters remain largely as they were, since we feel that the presentation of the basic ideas to be found there remains valid. Chapter 11 (Chapter 10 in the first edition) on chaotic phenomena is an example of this.

The only criticism anyone made to our faces was that we leaned too heavily on plasma physics and hydrodynamics for examples, whereas most phenomena and methods we consider have wider applications. These include optics, biology, solid state physics and other fields. This shortcoming has now been rectified to a certain extent. Also, a new chapter on soliton metamorphosis, including some colour plates, has been added (Chapter 10).

However, much of the text has been left as it was. Thus 'recent' should be read as recent in 1990. Some printing errors have been corrected. Once again, Dr Simon Capelin of the Cambridge University Press has been patient and helpful. Ms Lenkowska-Czerwińska spent a large portion of her time in Warwick helping us organize our material.

E. Infeld and G. Rowlands
Warwick

Contents

	<i>Foreword to the first edition</i>	page xi
	<i>Foreword to the second edition</i>	xiii
1	Introduction	1
1.1	Occurrence of nonlinear waves and instabilities in Nature	1
1.1.1	Nonlinear phenomena in our everyday experience	1
1.1.2	Nonlinear phenomena in the laboratory	3
1.2	Universal wave equations	5
1.2.1	The Korteweg–de Vries and Kadomtsev–Petviashvili equations and a first look at solitons	5
1.2.2	The nonlinear Schrödinger equation	8
1.2.3	Nonlinear optics	9
1.3	What is a plasma?	9
1.4	Wave modes on a water surface	12
1.4.1	Mathematical theory	12
1.4.2	Comments	14
1.5	Linear stability analysis and its limitations	18
1.6	Nonlinear structures	18
1.6.1	Coherent structures and pattern formation	22
1.7	Contents of Chapters 2–11	23
2	Linear waves and instabilities in infinite media	25
2.1	Introduction	25
2.2	Plasma waves	25
2.3	CMA diagrams	30
2.4	Instabilities	33
2.5	The Vlasov equation	37
2.6	Weak instabilities	43
	Exercises	47
3	Convective and non-convective instabilities; group velocity in unstable media	48
3.1	Introduction	48
3.2	Kinematics of unstable wavepackets	50

3.3	Moving coordinate systems	54
3.4	Higher dimensional systems	56
3.5	Summary	57
	Exercise	57
4	A first look at surface waves and instabilities	59
4.1	Introduction	59
4.2	Simple surface waves	61
4.3	The Rayleigh–Taylor instability	66
4.4	The Kelvin–Helmholtz instability	68
4.5	Solid–liquid interface instabilities	71
4.6	A first look at gravity wave instabilities	72
	4.6.1 The small amplitude onset of wave instability	73
	4.6.2 Further numerical results	75
4.7	Summary	81
	Exercises	81
5	Model equations for small amplitude waves and solitons; weakly nonlinear theory	82
5.1	Introduction	82
	5.1.1 Some physical equations ask for surgery	82
	5.1.2 Examples	83
5.2	A few model equations as derived by introducing a small parameter	85
	5.2.1 Shallow water, weak amplitude gravity waves	85
	5.2.2 Weak amplitude ion acoustic waves in an unmagnetized plasma	89
	5.2.3 Weak amplitude ion acoustic waves in a magnetized plasma	91
5.3	Weakly nonlinear waves	92
	5.3.1 Spreading, splitting and instabilities	92
	5.3.2 The story of deep water waves	98
	5.3.3 Mystery of the missing term	100
	5.3.4 Dynamics of a wavepacket	102
	5.3.5 Some generalizations	104
5.4	A general look at two families of model equations	107
5.5	A natural extension to finite amplitude waves due to Hayes	111
5.6	Temporal development of instabilities and wave-wave coupling	114
5.7	Concluding remarks	119
	Exercises	119
6	Exact methods for fully nonlinear waves and solitons	123
6.1	Introduction	123
6.2	Phase plane analysis and other methods	124

6.2.1	One stationary wave in a dissipationless medium	124
6.2.2	A two-fluid layer soliton pair	129
6.2.3	Weak ion acoustic shock waves in a collisional plasma	132
6.2.4	Solitons generated by laser fields	134
6.2.5	Solitons and domains in dipole chains	136
6.2.6	Discrete equations	139
6.3	Bernstein–Greene–Kruskal waves	143
6.3.1	Statistical description of a plasma and BGK waves	144
6.3.2	No trapped particles	145
6.3.3	Various limits	146
6.3.4	Trapped particle equilibria	147
6.3.5	Stability; subsequent developments	150
6.4	Lagrangian methods	152
6.5	Lagrangian interpolation	159
	Exercises	164
7	Cartesian solitons in one and two space dimensions	166
7.1	Introduction	166
7.2	The direct method	168
7.3	Constants of motion	171
7.4	Inverse scattering method	173
7.5	Bäcklund transformations	175
7.6	Entr'acte	177
7.7	Breathers and boundary effects	178
7.8	Experimental evidence	180
7.9	Plane soliton interaction in two space dimensions	181
7.9.1	Introducing the trace method	181
7.9.2	One and two soliton solutions	183
7.9.3	Some other developments and summary	188
7.10	Integrable equations in two space dimensions as treated by the Zakharov–Shabat method	192
7.10.1	Lax pairs and the PDEs they represent	193
7.10.2	Extension to x, y, t	194
7.10.3	How to proceed from the Lax pair to the general solution	195
7.10.4	An example: the Kadomtsev–Petviashvili equation	196
7.11	Summary	199
	Exercises	200
8	Evolution and stability of initially one-dimensional waves and solitons	202
8.1	A brief historical survey of large amplitude nonlinear wave studies	202
8.1.1	Solitons	204
8.1.2	Water waves are unstable	206
8.1.3	The geometrical optics limit	207

8.1.4	More recent results	211
8.1.5	What the remainder of Chapter 8 is about	212
8.2	Four methods as illustrated by the nonlinear Klein–Gordon equation	213
8.2.1	Whitham I	214
8.2.2	Whitham II	219
8.2.3	K expansion	220
8.2.4	Hayes	223
8.3	Higher dimensional dynamics	224
8.3.1	Kadomtsev–Petviashvili as analysed by Whitham II	224
8.3.2	Various limits	230
8.3.3	Common features of the weak amplitude and soliton limits for $\psi = 0$	231
8.3.4	Group velocity	233
8.3.5	Zakharov–Kuznetsov as analysed by K expansion	236
8.3.6	The variational method	244
8.4	A more physical approach leading to an assessment of models	245
8.4.1	Form of the waves considered	245
8.4.2	Unmagnetized plasmas, $\Omega_c = 0$	246
8.4.3	Magnetized plasmas, $\Omega_c > 0$	249
8.5	Dynamics of nonlinear wave, shock and soliton solutions to the cubic nonlinear Schrödinger equation	252
8.5.1	Results of a general stability calculation	253
8.5.2	One-dimensional dynamics: $\psi = 0$	254
8.5.3	Oblique and perpendicular propagation of perturbations	257
8.6	The direct K method	258
8.6.1	Transverse instability of Zakharov–Kuznetsov solitons	259
8.6.2	The Cahn–Hilliard equation	263
8.7	Some general conclusions and possible future lines of investigation	264
	Exercises	265
9	Cylindrical and spherical solitons in plasmas and other media	268
9.1	Interest in higher dimensional plasma solitons	268
9.2	Unidirectional cylindrical and spherical ion acoustic solitons	269
9.2.1	Model equations in non-Cartesian geometry	269
9.2.2	Cylindrical soliton equations CI and CII	269
9.2.3	Spherical solitons	271
9.2.4	Summary	272
9.3	Properties of unidirectional soliton equations	272
9.3.1	Integrability by inverse scattering	272
9.3.2	Conservation laws	273
9.4	Soliton solutions as compared with numerics and experiments	275
9.4.1	Exact solutions to CI	277
9.4.2	Initial value problem and experiments	277
9.4.3	Reflection from the axis (centre)	280
9.4.4	Models	284

9.4.5	Stability of cylindrical solitons	287
9.5	Langmuir solitons	287
9.5.1	Integrability	288
9.5.2	Stability of Langmuir solitons	288
9.6	Interacting solitons and some conclusions	291
9.7	Epilogue. Some other examples of spherical and cylindrical solitons	292
	Exercises	294
10	Soliton metamorphosis	296
10.1	The next step in investigating soliton behaviour	296
10.2	Decay of line KPI solitons in two dimensions	297
10.3	Decay of 2D solitons in three dimensions	301
10.3.1	2D solitons perturbed perpendicular to the motion	301
10.3.2	2D solitons perturbed parallel to the velocity	302
10.4	Conclusions	303
	Exercises	303
11	Non-coherent phenomena	304
11.1	Introduction	304
11.2	Bifurcation sequences and chaos	311
11.3	Flows and maps	326
11.4	Strange attractors	329
11.5	Effect of external noise	340
11.6	Experimental evidence for strange attractors	341
11.7	Other theories of turbulence	343
11.8	Conclusions	344
	Exercises	345
	Appendices	346
A1	Parameter stretching as suggested by the linear dispersion relations	346
A1.1	Ion acoustic waves in an unmagnetized plasma, $\Omega_c = 0$	346
A1.2	Magnetized plasmas, $\Omega_c > 0$	347
A2	Relation between the trace method and the inverse scattering method	349
A3	Some formulae for perturbed nonlinear ion acoustic waves and solitons	351
A3.1	No magnetic field	351
A3.2	$\Omega_c > 0$	352
A4	Colliding soliton theory	354
A5	A model equation for spherical solitons	356
A6	Stability calculation for 2D KPI soliton in 3D	358
	<i>References</i>	360
	<i>Author index</i>	379
	<i>Subject index</i>	387

Introduction

1.1 *Occurrence of nonlinear waves and instabilities in Nature*

This book is concerned with the propagation of waves and instabilities both linear and nonlinear, but concentrating on the latter.

The main advances in this subject have quite naturally come from studies involving fluids and more recently plasmas. The latter primarily because of the possibility of ‘cheap, unlimited’ (and hopefully safe) power from thermonuclear reactions. Everyone is of course familiar with waves on water if only being aware of the many instances where they provide examples of natural beauty. It is not so obvious that very similar waves can exist in a plasma which, to a good approximation, is usually a very dilute assembly of ions and electrons. We shall see later in this chapter that this is indeed so and fluids and plasmas have much in common. However, plasmas also show a much wider range of phenomena basically because they are composed of two or more components and also can be made strongly anisotropic by the introduction of magnetic fields, something that is not possible for simple fluids. This richer variety of phenomena has also been a reason why plasmas have had more than their share of attention.

The above remarks notwithstanding, there are numerous media other than plasmas and fluids which can support waves and/or propagate instabilities. As we will see, some of these are more intriguing than others.

1.1.1 *Nonlinear phenomena in our everyday experience*

As most of us are aware, waves generated by the wind can propagate across a field of corn. In this case the microscopic model is that of the ears moving, due to the stalk bending, in an harmonic manner, and ear interacting with adjacent ear only when in contact (a hard sphere potential). On a macroscopic level the corn can be considered as a dense fluid and now with the moving air flow over it one has the classic situation of a Kelvin–Helmholtz instability (Chapter 4).

The wind drives the instability and the stalks of corn bend in an analogous manner to how water waves are formed on lakes by the wind. The nonlinear requirement is different, in that if the bending of the stalk is too great it will break and produce a permanent record of the wave. Kelvin–Helmholtz type instabilities occur in plasma physics and they have been controlled to a



Figure 1.1 Two examples of herring-bone cloud formation. (After R. Scorer, *Clouds of the world*, Lothian, Melbourne (1972).)

certain extent by introducing perpendicular periodicities. The intriguing question naturally arises if similar conditions would stabilize the motion of corn heads and stop the breaking of the stalks. By analogy this could be done by planting trees periodically spaced in a line (or lines) perpendicular to the usual direction of the wind. Another example where Nature leaves a permanent record of a surface wave instability (at least until diffusion processes slowly remove it) is the so-called herring-bone cloud formation. Here the sky is broken up into alternate bands of cloud (high density of moisture regions) and apparently clear regions (low density regions). The moisture in this case is responsible for the permanent record, two layers of air moving relative to each other giving rise to the Kelvin–Helmholtz instability (Chapter 4) and subsequent nonlinear effects (Fig. 1.1).

Television coverage of soccer matches usually shows, incidentally, the swaying of the crowd. This is seen as a wave moving through the stadium. Unfortunately, in some circumstances this wave can build up and those near the barriers can get crushed.

Lighthill and Whitham (1955) and Richards (1956) have considered the flow of cars (discrete objects) in a fluid context and explained phenomena such as the effects of traffic lights in terms of the propagation of waves and, in particular, shocks.

The conclusion to be reached from these quite disparate examples is that they can be effectively studied in terms of the propagation of waves and instabilities, leading to nonlinear effects, in continuous media. If the natural wavelength is large compared to the underlying microscopic length, the above picture should be applicable.

1.1.2 *Nonlinear phenomena in the laboratory*

Waves in solids have received considerable attention both at the microscopic (atomic) and the macroscopic (continuous) level. The most interesting phenomenon concerning the propagation of disturbances in solids at a microscopic level is the effect of anisotropies and non-homogeneities in the media. Until recently, nonlinear effects in solids have received little attention, as the energy needed to produce them is very large. However, with the advent of intense power sources such as lasers it has been possible to show that the flow of heat in solids is closely related to the flow of solitons. The basic relationships between soliton amplitude, width and velocity have been verified in this context (Section 1.2 and, in some detail, Chapters 5 and 7).

At the microscopic level it is usually necessary to quantize the system and instead of talking about sound waves one talks about phonons. However, because of the lattice periodicity the phonons have a dispersive nature. Most interesting phenomena, such as thermal conductivity, depend on phonon–phonon interaction. Until recently, such interaction has been studied in what could be called weakly nonlinear theories. However, a major breakthrough was made when it was realized that a number of phenomena could best be explained by introducing solitons as elementary excitations. Then it was found that a statistical mechanics based on weakly interacting solitons and phonons gave better results than previous theories which attempted to consider phonon–phonon interaction outside the control of a weak interaction.

Many years ago, Fermi, Ulam and Pasta (1965), though the actual work was performed much earlier and published in a Los Alamos report in 1955, studied numerically the problem of the strong interaction of phonons. They found, somewhat perplexingly at that time, that the




Figure 1.2 Evolution of an initially periodic profile, $\cos \pi x$, as given by the Korteweg-de Vries equation (1.2.8). The breaking time for the wave profile (when the third term is neglected) is t_B . From Zabusky and Kruskal (1965). After a while, patterns roughly repeat themselves.

phonons did not come to thermal equilibrium, but rather they underwent nearly periodic variations. See Weissert (1998).

Much later Zabusky and Kruskal (1965) showed that this was the correct behaviour and *could* be explained in terms of solitons in a space periodic medium, Fig. 1.2. (A name given to reflect their quantal nature.) Nowadays people realize that solitons are not *necessary* to explain this phenomenon (Thyagaraja (1983)) Chapter 5. A theory that relies on the interaction of a small number of periodic modes also exists, see Infeld (1981c). Solitons are special wave pulses which interact with one another so as to keep their basic identity and so that they act as particles. Now the soliton has come of age in its mother subject, though in the meantime promising to be a useful concept in many other branches of physics, most of them being well out of the range of quantum effects. One of the first detailed experimental verifications of soliton type behaviour was in the study of nonlinear ion acoustic waves.

Davydov (1978), (1985) has applied some of the rules of solid state physics to the transport of energy down protein chains. He assumed that the idea of soliton propagation is relevant to a study of chemical changes taking place in long protein molecules. This leads to the transfer of ATP (adenosine triphosphate) and could be the basis for an understanding of muscle contraction. However, Davydov's theory is one-dimensional, whereas proteins are three-dimensional. Also, his mechanism for energy transport has been criticized. Zorski and Infeld (1997) approach protein dynamics by using a quadrilateral chip model. The dynamics were described by continuum equations in three dimensions. They obtained a helical structure and demonstrated stability. Solitons do appear in special cases, but have limited importance in their theory. Thus, the role of soliton dynamics in understanding protein chains still seems to be an open question.

The motion of electrons in solids is surprisingly well understood in terms of a simple Drude picture where the current J is linearly related to the electric field E by $J = (ne^2\tau/m)E$. Here n is the electron density, e its charge, and τ the mean free time. The effective mass ' m ' takes into account the presence of the periodic nature of atomic structure. Normally all the quantities are constants and we simply have Ohm's law. However, it can happen, particularly in semiconductors, that if E is large enough the electron can be excited to a higher band which can have a different value of m . Thus the conductivity $\sigma (= ne^2\tau/m)$ is dependent on E and Ohm's law is no longer linear. If the

mass is larger in the higher band, then an increase in E causes a decrease in J and thus we have a negative differential resistance. Obviously this is an instability mechanism and such a mechanism is observed in GaAs (gallium arsenide). The instability itself leads to the propagation of nonlinear stable pulses called Gunn domains, analogues to a soliton. These have been observed and are in fact the bases of many modern day oscillators. See Butcher (1967) for a review of Gunn domains; and Butcher and Rowlands (1968) for a study of the stability of the domain.

Another instability, leading to the propagation of nonlinear pulses that are also kinsmen of solitons and have been studied in the general area of solid state physics, is that associated with the acousto-electric effect. Here the instability mechanism arises because of a piezo-electric coupling between the propagation of sound waves and the flow of electrons. A nonlinear pulse can propagate down a crystal, for example of CdS or ZnO, but reach such an amplitude as to cause permanent distortion to the crystal. For a brief account of the general theory and for a discussion of the instability and nonlinear effects see Pawlik and Rowlands (1975).

It took a long time for it to be accepted that a homogeneous mixture of chemicals could lead to a periodic time variation in the concentration of a particular chemical or to an inhomogeneous spatial separation of the chemicals. Turing demonstrated that nonlinear chemical reactions together with diffusion could lead to a spatial separation of the chemicals and explored the idea in connection with the theory of morphogenesis, the formation of life. Zhabotinsky (1964) and later Zhabotinsky and Belousov found experimentally that a homogeneous mixture of certain chemicals could lead to a time periodic variation of the colour of the mixture. Later experiments showed that the same mechanism could lead to spatial colour patterns. All these phenomena can be explained in terms of nonlinear waves in time and space that have the inherent stability of a soliton.

In this revised version of the book we will look at nonlinear phenomena in new contexts, such as laser theory and biology. Waves following from discrete equations are also a novelty, as are illustrations of soliton metamorphosis (solitons are a kind of compact wave, about which more in Subsection 1.2).

1.2 *Universal wave equations*

It has now been realized that the study of many different types of waves in numerous media can often be based on a few universal nonlinear equations. These replace the usual linear wave equations, such as

$$\left(\frac{\partial^2}{\partial t^2} - \nabla^2\right)\Phi = 0.$$

1.2.1 *The Korteweg–de Vries and Kadomtsev–Petviashvili equations and a first look at solitons*

Most classical media propagate longitudinal plane waves at or near a characteristic velocity c (acoustic-type waves). These waves are, for very small amplitude, given by