



南京航空航天大学
研究生系列精品教材

Introduction to Matrix Theory

(矩阵论引论)

Zhengsheng Wang




科学出版社

南京航空航天大学研究生系列精品教材

Introduction to Matrix Theory

(矩阵论引论)

Zhengsheng Wang



Responsible Editors: Zhongxing Zhang

Copyright© 2015 by Science Press
Published by Science Press
16 Donghuangchenggen North Street
Beijing 100717, China

Printed in Beijing

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the copyright owner.

ISBN 978-7-03-045394-5 (Beijing)

Preface

About matrix theory

The study of matrix theory occupies a singular place within mathematics. No applied mathematician can be properly trained without some basic understanding of matrix theory and linear algebra. It is still an area of active research, and it is used by every mathematician and by many scientists working in various specialities. Here we list several examples to illustrate its versatility.

(1) Scientific computing libraries begin growing around matrix computation. As a matter of fact, the discretization of partial differential operators is an endless source of linear finite-dimensional problems.

(2) At a discrete level, the maximum principle is related to nonnegative matrices.

(3) Control theory and stabilization of systems with finitely many degrees of freedom involve special analysis of matrices.

(4) The discrete Fourier transform (FT), including the fast Fourier transform (FFT), makes use of Toeplitz matrices.

(5) Statistics is widely based on correlation matrices.

(6) The generalized inverse is involved in least-squares approximation.

(7) Symmetric matrices are inertia, deformation, or viscous tensors in continuum mechanics.

(8) Markov process involves stochastic or bistochastic matrices.

(9) Graphs can be described in a useful way by square matrices.

(10) Quantum chemistry is intimately related to matrix groups and their representations.

About this book

Our goal is to provide a textbook for a course in matrix theory and methods accessible to advanced undergraduate and beginning graduate students. Through the course, students learn, practice, and master basic matrix results and techniques that are very useful for applications in various fields such as mathematics, statistics, physics, computer science, and engineering, etc. This book is an attempt to provide some of the required knowledge and understanding. It is written in a spirit that considers matrix theory not merely as a tool for solving applied problems but also as a challenging and rewarding part of mathematics.

This book has evolved over six years from the author's lecture notes of the graduate level course "matrix theory" for international students at Nanjing University of Aeronautics and Astronautics (NUAA). The book can serve as textbook or reference for graduate students in Master or PhD degree. The only prerequisites are a decent background in elementary

linear algebra and calculus. This book can also be used as a reference for researchers and engineers. In order to make the book accessible not only to mathematicians but also to scientists and engineers, I have planned it to be as self-contained as possible.

This book is organized into nine chapters and consists of the following topics.

- (1) Review and Miscellanea: Basic Concepts in Linear Algebra.
- (2) Linear Space and Inner-product Space.
- (3) Linear Transformation.
- (4) Jordan Canonical Form.
- (5) Matrix Factorization.
- (6) Hermitian Matrix and Positive Definite Matrix.
- (7) Matrix Norm and Matrix Analysis.
- (8) Generalized Inverse Matrix.
- (9) An Introduction to MATLAB.

The author has benefited from numerous books and journals. This book would not exist without the earlier works of a great number of authors. The references at the end of the book are intended as a possible guide to some of the literature covering the topics of the individual chapters more exhaustively.

The author would appreciate any comments, suggestions, questions, criticisms, or corrections that readers may take the trouble of communicating to me. Readers are welcome to communicate me via e-mail: wangzhengsheng@nuaa.edu.cn.

Zhengsheng Wang
wangzhengsheng@nuaa.edu.cn
NUAA, Nanjing
March 2015

Acknowledgments

I would like to reiterate my thanks to many persons who have assisted me in the writing of this book.

I am grateful to Prof. Hua Dai. I was taught matrix theory and computation in the 1990s by my MS and PhD thesis adviser Professor Hua Dai at Nanjing University of Aeronautics and Astronautics (NUAA). Prof. Dai's perspective on teaching mathematics in general and numerical algebra in particular had a great and long-lasting impact on my own teaching.

I gratefully acknowledge a number of professors and friends for many valuable suggestions and helpful remarks that both greatly improved the quality of the book and have been improving my research work in matrix theory and computation. Among them are Hua Dai (NUAA), Zhong-zhi Bai (Institute of Computational Mathematics and Scientific/Engineering Computing of Chinese Academy of Sciences) and Lothar Reichel (Kent State University, USA).

I am thankful to my colleagues and my students who have had to put up with my talking to them so often about matrices and their constant interest and useful comments. I am also grateful to Science Press staff members who handled this book, especially editor Zhongxing Zhang, the NUAA Graduate School staff members for help with publishing this book. I also thank College of Science of NUAA for providing released time for me to work on the project.

This work is partly supported by the Natural Science Foundation of Jiangsu Province of China under grant No.BK20141408, the Graduate Education Excellent Project Program of NUAA, the Jiangsu Oversea Research & Training Program for Prominent Young & Middle-Aged University Teachers and Presidents, the National Natural Science Foundation of China under grant No.61473148 and the Fundamental Research Funds for the Central Universities under grant No.NZ2014101.

Finally, I would also like to express my appreciation to my family.

Zhengsheng Wang
NUAA, Nanjing
March 2015



Frequently Used Notations and Terminology

\mathcal{C} -All complex numbers

\mathcal{R} -All real numbers

\mathcal{C}^n -All complex vectors of order n

\mathcal{R}^n -All real vectors of order n

$\mathcal{C}^{m \times n}$ -All $m \times n$ complex matrices

$\mathcal{R}^{m \times n}$ -All $m \times n$ real matrices

$\dim(V)$ -Dimension of vector space V

$A = (a_{ij})$ -Matrix A with (i, j) -entry a_{ij}

I -Identity matrix

A^T (A^H) -Transpose (Conjugate transpose) of matrix A

A^{-1} -Inverse of matrix A

$\text{Rank}(A)$ -Rank of matrix A

$|A|$ or $\det(A)$ -Determinant of matrix A

$\text{tr}(A)$ -Trace of matrix A

$\|\cdot\|$ -Norm of a vector or matrix

(u, v) -Inner product of vectors u and v

$\rho(A)$ -Spectral radius of matrix A

$\sigma_{\max}(A)$ -Largest singular value (spectral norm) of matrix A

$\lambda_{\max}(A)$ -Largest eigenvalue of matrix A

An $n \times n$ matrix A is said to be

(1) Hermitian if $A^H = A$

(2) normal if $A^H A = A A^H$

(3) unitary if $A^H A = A A^H = I$, i.e., $A^{-1} = A^H$

(4) positive semidefinite if $x^H A x \geq 0$ for all vectors $x \in \mathcal{C}^n$

(5) similar to B if $P^{-1} A P = B$ for some invertible matrix P

- (6) unitary similar to B if $U^H A U = B$ for some unitary matrix U
- (7) diagonalizable if $P^{-1} A P$ is diagonal for some invertible matrix P
- (8) upper-triangular if all entries below the main diagonal are zero
- (9) λ is an eigenvalue of $A \in \mathcal{C}^{n \times n}$ if $Ax = \lambda x$ for some nonzero vector $x \in \mathcal{C}^n$

Contents

Chapter 1	Review and Miscellanea: Basic Concepts in Linear Algebra	1
1.1	Matrix Concept and Special Matrices	1
1.2	Matrix Algebra	3
1.3	Eigenvalue and Eigenvector	7
1.4	Reading: Cayley and Sylvester	10
1.5	Self Assessment Exercises	11
Chapter 2	Linear Space and Inner-product Space	13
2.1	Linear Space (Vector Space)	13
2.2	Basis and Dimension of Vector Space	17
2.3	Subspaces	20
2.4	Inner Product Space	21
2.5	Reading: Gram and Schmidt	24
2.6	Self Assessment Exercises	25
Chapter 3	Linear Transformation	27
3.1	Introduction	27
3.2	Linear Transformation with Properties	27
3.3	The Range and Kernel	30
3.4	Matrix Representation of Linear Transformation	32
3.5	Similarity	36
3.6	Reading: Cramer and Gershgorin	38
3.7	Self Assessment Exercises	39
Chapter 4	Jordan Canonical Form	41
4.1	Diagonalizability	41
4.2	Jordan Block Matrix and Jordan Form Matrix	44
4.3	λ -matrix and Smith Standard Form	45
4.4	Jordan Canonical Form	49
4.5	Cayley-Hamilton Theorem and Minimal Polynomial	53
4.6	MATLAB Commands for Jordan Canonical Form	55
4.7	Reading: Camille Jordan	57
4.8	Self Assessment Exercises	58
Chapter 5	Matrix Factorization	60
5.1	Introduction	60
5.2	Full Rank Decomposition	60
5.3	LU Factorization	63

5.4	QR Factorization	67
5.5	Schur Decomposition	69
5.6	Singular Value Decomposition(SVD)	71
5.7	An Example of Application: Image Compression	74
5.8	MATLAB Commands for Matrix Factorization	76
5.9	Reading: Householder and Householder Award	78
5.10	Self Assessment Exercises	80
Chapter 6 Hermitian Matrix and Positive Definite Matrix		81
6.1	Hermitian Matrix	81
6.2	Positive Definite Matrix	84
6.3	Reading: Charles Hermite	87
6.4	Self Assessment Exercises	88
Chapter 7 Matrix Norm and Matrix Analysis		89
7.1	Introduction	89
7.2	Vector Norm	89
7.3	Matrix Norm	91
7.4	Matrix Sequence, Series and Function	94
7.5	MATLAB Commands for Norms	98
7.6	Reading: Frobenius	99
7.7	Self Assessment Exercises	100
Chapter 8 The Moore-Penrose Generalized Inverse		102
8.1	Introduction	102
8.2	The Moore-Penrose Generalized Inverse	102
8.3	The Solvability of Linear Systems	105
8.4	MATLAB Commands for the Moore-Penrose Generalized Inverse Matrix	107
8.5	Reading: Moore and Penrose	108
8.6	Self Assessment Exercises	109
Chapter 9 An Introduction to MATLAB		110
9.1	A Glance of MATLAB	110
9.2	Starting Up	111
9.3	MATLAB as A Calculator	112
9.4	Plotting	119
9.5	Programming	122
9.6	Commonly Used Commands Summary	127
9.7	Reading: Golub and Moler	130
Answers to Selected Exercises		132
Bibliography		149

Chapter 1

Review and Miscellanea: Basic Concepts in Linear Algebra

We briefly review, mostly without proof, the basic concepts and results taught in an elementary linear algebra course.

1.1 Matrix Concept and Special Matrices

The term of *matrix* was first introduced by the British mathematician James Joseph Sylvester in 1890. The word “matrix” is derived from the Indo-European root mater, meaning “mother”. Matrices are indeed the core of linear algebra.

Firstly, we see the following two numbers tables (Table 1.1 and Table 1.2) in real life.

Table 1.1 The price table of four kinds of cans in three supermarkets

	P1	P2	P3	P4
S1	17	7	11	21
S2	15	9	13	19
S3	18	8	15	19

We can write it in brief in the form

$$\begin{bmatrix} 17 & 7 & 11 & 21 \\ 15 & 9 & 13 & 19 \\ 18 & 8 & 15 & 19 \end{bmatrix}_{3 \times 4}$$

Table 1.2 The distances of three cities in China (km)

	Beijing	Nanjing	Shanghai	Hongkong
Beijing	0	900	1050	1980
Nanjing	900	0	300	1140
Shanghai	1050	300	0	1120
Hongkong	1980	1140	1120	0

We can write it in the brief form

$$\begin{bmatrix} 0 & 900 & 1050 & 1980 \\ 900 & 0 & 300 & 1140 \\ 1050 & 300 & 0 & 1120 \\ 1980 & 1140 & 1120 & 0 \end{bmatrix}_{4 \times 4}$$

It is symmetric.

Definition 1.1.1 (Matrix) An array of numbers (or symbols) in m rows and n columns is called an $m \times n$ matrix.

The notation

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

or

$$A = (a_{ij})_{m \times n}$$

Usually, we denote

$$A \in \mathcal{R}^{m \times n}, \quad A \in \mathcal{C}^{m \times n}$$

where $\mathcal{R}^{m \times n}$ and $\mathcal{C}^{m \times n}$ denote the set of $m \times n$ real and complex matrices.

The entry that appears at the intersection of the i th row and j th column is called the (i, j) entry.

An $n \times n$ matrix is called a square matrix of order n . If the number of rows is not equal to the number of columns, the the matrix is called rectangular.

A matrix obtained from a given matrix A by deleting some (but of course not all) of its rows and columns is called the submatrix of A .

Example 1.1.1

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Here, A is a 4×3 rectangular matrix, B is a square matrix of order 3, and B is a submatrix of A .

Definition 1.1.2 (Equality of matrices) Let $A = (a_{ij})_{m \times n}$, $B = (b_{ij})_{p \times q}$. Then A is said to be equal to B if

a) A and B have the same number of rows and the same number of columns, that is, $m = p$, $n = q$, and

b) the corresponding entries of A and B are equal, that is, $a_{ij} = b_{ij}$ for each pair of subscripts (i, j) .

Here some special matrices which have special structure or/and special properties are introduced as follows.

Definition 1.1.3 (Zero matrix) The $m \times n$ matrix in which every entry is 0 is called the $m \times n$ zero matrix and is denoted by $0_{m \times n}$ or simply 0.

Note that

$$A + (-A) = A - A = 0, \quad A + 0 = A$$

Definition 1.1.4 (Diagonal matrix) An $n \times n$ matrix is called a diagonal matrix if each entry that is not on the diagonal is 0.

Example 1.1.2

$$D = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3}$$

Definition 1.1.5 (Identity matrix) An $n \times n$ matrix is called an identity matrix if it is a diagonal matrix and all the entries on the diagonal are equal to 1.

Example 1.1.3

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3}$$

Definition 1.1.6 (Upper (Lower) triangular matrix) An $n \times n$ matrix is called an upper (lower) triangular matrix if all the entries below (above) the diagonal are zero.

Example 1.1.4

$$L = \begin{bmatrix} 3 & 0 & 0 \\ 4 & 5 & 0 \\ 7 & 9 & 1 \end{bmatrix}_{3 \times 3}, \quad U = \begin{bmatrix} 3 & 5 & 7 \\ 0 & 5 & 10 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3}$$

1.2 Matrix Algebra

Definition 1.2.1(Scalar multiplication) Let α be a number, and let $A = (a_{ij})_{m \times n}$ be a matrix. Then

$$\alpha A = A\alpha = (\alpha a_{ij})_{m \times n}$$

This means that αA is the matrix obtained by multiplying each entry of A by the same number α .

Example 1.2.1

$$3 \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} 3 & 6 & 9 \\ 12 & 15 & 18 \\ 21 & 24 & 27 \end{bmatrix}$$

Definition 1.2.2 (Addition of matrices) Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{m \times n}$ be $m \times n$ matrices. Then

$$A + B = (c_{ij})_{m \times n}$$

where $c_{ij} = a_{ij} + b_{ij}$ for each pair (i, j) .

Example 1.2.2

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} + \begin{bmatrix} 0 & -1 & 1 \\ 2 & -5 & 8 \\ 1 & 0 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 4 \\ 6 & 0 & 14 \\ 8 & 8 & 7 \end{bmatrix}$$

The negative of A is written as $-A$.

$$A = \begin{bmatrix} 1 & 1 & 4 \\ 6 & 0 & 14 \\ 8 & 8 & 7 \end{bmatrix}, \quad -A = \begin{bmatrix} -1 & -1 & -4 \\ -6 & 0 & -14 \\ -8 & -8 & -7 \end{bmatrix}$$

Note that matrix addition is both commutative and associative, that is the theorem as follows.

Theorem 1.2.1 Let A, B, C be $m \times n$ matrices. Then

- a) $A + B = B + A$ (commutative law of addition),
- b) $(A + B) + C = A + (B + C)$ (associative law of addition).

Definition 1.2.3 (Matrix multiplication) If $A = (a_{ij})_{m \times n}$, $B = (b_{ij})_{n \times r}$, then the product $AB = C = (c_{ij})_{m \times r}$ matrix whose entries are defined by

$$c_{ij} = a(i, :)b(:, j) = \sum_{k=1}^n a_{ik}b_{kj}$$

Example 1.2.3

$$A = \begin{bmatrix} 3 & -2 \\ 2 & 4 \\ 1 & -3 \end{bmatrix}, \quad B = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 1 & 6 \end{bmatrix}$$

then

$$C = AB = \begin{bmatrix} -14 & 1 & -3 \\ 12 & 6 & 30 \\ -14 & -2 & -15 \end{bmatrix}$$

Matrix multiplication obeys many of the familiar laws of arithmetic apart from the commutative law.

Theorem 1.2.2 Suppose $A_{m \times n}$, $B_{n \times p}$, $C_{p \times q}$.

- 1) $(AB)C = A(BC)$;
- 2) $t(AB) = (tA)B = A(tB)$, $A(-B) = (-A)B = -(AB)$;
- 3) $(A + B)C = AC + BC$;
- 4) $D(A + B) = DA + DB$.

In general, $AB \neq BA$. If $AB = BA$, we call that A commutes with B .

Example 1.2.4 (Application: A simple model for marital status computations) In a certain town, 30 percent of the married women get divorced each year and 20 percent of the single women get married each year. There are 8000 married women and 2000 single women. Assuming that the total population of women remains constant, how many married women and how many single women will there be after 1 year? After 2 years?

Solution Form a matrix A as follows. The entries in the first row of A will be the percent of married and single women, respectively, that are married after 1 year. The entries in the second row will be the percent of women who are single after 1 year. Thus

$$A = \begin{bmatrix} 0.70 & 0.20 \\ 0.30 & 0.80 \end{bmatrix}$$

If we let $x = \begin{bmatrix} 8000 \\ 2000 \end{bmatrix}$, the number of married and single women after 1 year can be computed by multiplying A times x .

$$Ax = \begin{bmatrix} 0.70 & 0.20 \\ 0.30 & 0.80 \end{bmatrix} \begin{bmatrix} 8000 \\ 2000 \end{bmatrix} = \begin{bmatrix} 6000 \\ 4000 \end{bmatrix}$$

After 1 year there will be 6000 married women and 4000 single women. To find the number of married women and single women after 2 years, compute

$$A^2x = A(Ax) = \begin{bmatrix} 0.70 & 0.20 \\ 0.30 & 0.80 \end{bmatrix} \begin{bmatrix} 6000 \\ 4000 \end{bmatrix} = \begin{bmatrix} 5000 \\ 5000 \end{bmatrix}$$

After 2 years, half of the women will be married and half will be single. In general, the number of married and single women after n years can be determined by computing $A^n x$.

Definition 1.2.4 (Transpose of a matrix) Let $A = (a_{ij}) \in \mathcal{C}^{m \times n}$ be an $m \times n$ matrix. Then the transpose of A , written A^T , is the $n \times m$ matrix whose (i, j) entry is a_{ji} for all i, j . In other words, the i th row of A^T is the i th columns of A for all i .

Definition 1.2.5 (Symmetric matrix) An $n \times n$ matrix A is called symmetric if $A^T = A$. In other words, $A = (a_{ij})$ is symmetric if $a_{ij} = a_{ji}$ for all i, j .

For example, the following matrix is symmetric.

Example 1.2.5

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 6 \\ 3 & 6 & 9 \end{bmatrix}$$

Definition 1.2.6 (Conjugate transpose of a matrix) Let $A = (a_{ij}) \in \mathcal{C}^{m \times n}$ be an $m \times n$ matrix. Then the conjugate transpose of A , written $A^H (A^*)$, is the $n \times m$ matrix whose (i, j) entry is $\overline{a_{ji}}$ for all i, j .

Example 1.2.6

$$A = \begin{bmatrix} 1 & 2-i & 3 \\ 4 & 5i & 6 \\ 7+8i & 8 & 9 \end{bmatrix}, \quad A^T = \begin{bmatrix} 1 & 4 & 7+8i \\ 2-i & 5i & 8 \\ 3 & 6 & 9 \end{bmatrix}$$

$$A^H = \begin{bmatrix} 1 & 4 & 7-8i \\ 2+i & -5i & 8 \\ 3 & 6 & 9 \end{bmatrix}$$

Definition 1.2.7 (Trace of matrix) The trace of an $n \times n$ matrix $A = (a_{ij})$ is defined to be

$$\text{tr}(A) = a_{11} + a_{22} + \cdots + a_{nn} = \sum_{i=1}^n a_{ii}$$

Note that

$$\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B), \quad \text{tr}(AB) = \text{tr}(BA)$$

Example 1.2.7

$$\text{tr}(A) = \text{tr} \left(\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \right) = 15$$

Definition 1.2.8 (Determinant of 1×1 and 2×2 matrices)

$$\det(A) = |A| = \left| \begin{bmatrix} a_{11} \end{bmatrix} \right| = a_{11}$$

$$\det(A) = |A| = \left| \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \right| = a_{11}a_{22} - a_{12}a_{21}$$

Example 1.2.8

$$\det(A) = |A| = \left| \begin{bmatrix} 3 \end{bmatrix} \right| = 3$$

$$\left| \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \right| = 1 \times 4 - 2 \times 3 = -2$$

Definition 1.2.9 (Laplace expansion) Assume that the determinant function has been defined for matrices of size $(n-1) \times (n-1)$. Then $\det(A)$ is defined by the so called first-row Laplace expansion:

$$\begin{aligned} \det(A) &= a_{11}M_{11}(A) - a_{12}M_{12}(A) + \cdots + (-1)^{1+n}M_{1n}(A) \\ &= \sum_{j=1}^n (-1)^{1+j}a_{1j}M_{1j}(A) \\ &= \sum_{j=1}^n a_{1j}C_{1j} \end{aligned}$$

Example 1.2.9 $A = [a_{ij}]$ is a 3×3 matrix, the Laplace expansion gives

$$\begin{aligned} \det(A) &= a_{11}M_{11}(A) - a_{12}M_{12}(A) + a_{13}M_{13}(A) \\ &= a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31}) \\ &= a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31} \end{aligned}$$

Definition 1.2.10 (Minor) Let $M_{ij}(A)$ (or simply M_{ij} if there is no ambiguity) denote the determinant of the $(N-1) \times (N-1)$ submatrix of A formed by deleting the i th row and j th column of A . ($M_{ij}(A)$ is called the (i, j) minor of A or the minor of entry a_{12})

Definition 1.2.11 (Cofactors) The cofactors of entry a_{ij} is definition as

$$C_{ij} = (-1)^{i+j}M_{ij}(A)$$

Definition 1.2.12 (The adjoint of matrix) The adjoint of matrix A is defined as

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ \vdots & \vdots & & \vdots \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{bmatrix}$$

Definition 1.2.13 (Inverse of matrix (Nonsingular matrix)) A square matrix $A \in \mathcal{C}^{n \times n}$ is called nonsingular or invertible if there exists a matrix $B \in \mathcal{C}^{n \times n}$ such that

$$AB = I_n = BA$$

Any matrix B with the above property is called an inverse of A , denote $A^{-1} = B$. If A does not have an inverse, A is called singular.

Theorem 1.2.3 (Inverses are unique) If A has inverses B and C , then $B = C$.

Theorem 1.2.4 (sufficient and necessary condition of nonsingular) If A is nonsingular (or invertible) $\Leftrightarrow \det(A) \neq 0$. In another word, a square matrix is invertible if and only if its determinant is nonzero. and

$$A^{-1} = \frac{1}{\det(A)} \text{adj}(A)$$

Definition 1.2.14 (Elementary row operations on a matrix: elementary row matrices) There are three types, E_{ij} ; $E_i(t)$; $E_{ij}(t)$, corresponding to the three kinds of elementary row operation:

- 1) $E_{ij}(i \neq j)$ is obtained from the identity matrix I_n by interchanging rows i and j ;
- 2) $E_i(t)(t \neq 0)$ is obtained by multiplying the i th row of I_n by t ;
- 3) $E_{ij}(t)(i \neq j)$ is obtained from I_n by adding t times the j th row of I_n to the i th row.