

CLASSICS IN MATHEMATICS

Kôsaku Yosida

Functional Analysis

Sixth Edition

泛函分析

第6版

Springer-Verlag

世界图书出版公司

Kôzaku Yosida

Functional Analysis

Sixth Edition

Springer-Verlag

世界图书出版公司

北京 · 广州 · 上海 · 西安

书 名: Functional Analysis 6th ed.

作 者: K.Yosida

中译名: 泛函分析 第6版

出版者: 世界图书出版公司北京公司

印刷者: 北京世图印刷厂

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64015659, 64038347

电子信箱: kjsk@vip.sina.com

开 本: 24 开 印 张: 21.5

出版年代: 1999 年 6 月第 1 版 2004 年 7 月第 2 次印刷

书 号: 7-5062-2611-1/O · 242

定 价: 75.00 元

世界图书出版公司北京公司已获得 Springer-Verlag 授权在中国大陆
独家重印发行。

Kôzaku Yosida

1157-28 Kajiware, Kamakura, 247/Japan

AMS Subject Classification (1970): 46-XX

ISBN 3-540-10210-8 Springer-Verlag Berlin Heidelberg New York

ISBN 0-387-10210-8 Springer-Verlag New York Heidelberg Berlin

ISBN 3-540-08627-7 5. Auflage Springer-Verlag Berlin Heidelberg New York

ISBN 0-387-08627-7 5th edition Springer-Verlag New York Heidelberg Berlin

Library of Congress Cataloging in Publication Data. Yosida, Kôzaku, 1909 -
Functional analysis (Grundlehren der mathematischen Wissenschaften; 123).

Bibliography: p. Includes index. 1. Functional analysis, I. Title. II. Series.

QA320.16. 1980. 515.7. 80-18567.

ISBN 0-387-10210-8 (U.S.)

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks. Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher.

© by Springer-Verlag Berlin Heidelberg 1965, 1971, 1974, 1978, 1980

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the People's Republic of China only and not for export therefrom.

Reprinted in China by Beijing World Publishing Corporation, 1999

Preface to the First Edition

The present book is based on lectures given by the author at the University of Tokyo during the past ten years. It is intended as a textbook to be studied by students on their own or to be used in a course on Functional Analysis, i.e., the general theory of linear operators in function spaces together with salient features of its application to diverse fields of modern and classical analysis.

Necessary prerequisites for the reading of this book are summarized, with or without proof, in Chapter 0 under titles: Set Theory, Topological Spaces, Measure Spaces and Linear Spaces. Then, starting with the chapter on Semi-norms, a general theory of Banach and Hilbert spaces is presented in connection with the theory of generalized functions of S. L. SOBOLEV and L. SCHWARTZ. While the book is primarily addressed to graduate students, it is hoped it might prove useful to research mathematicians, both pure and applied. The reader may pass, e.g., from Chapter IX (Analytical Theory of Semi-groups) directly to Chapter XIII (Ergodic Theory and Diffusion Theory) and to Chapter XIV (Integration of the Equation of Evolution). Such materials as "Weak Topologies and Duality in Locally Convex Spaces" and "Nuclear Spaces" are presented in the form of the appendices to Chapter V and Chapter X, respectively. These might be skipped for the first reading by those who are interested rather in the application of linear operators.

In the preparation of the present book, the author has received valuable advice and criticism from many friends. Especially, Mrs. K. HILLE has kindly read through the manuscript as well as the galley and page proofs. Without her painstaking help, this book could not have been printed in the present style in the language which was not spoken to the author in the cradle. The author owes very much to his old friends, Professor E. HILLE and Professor S. KAKUTANI of Yale University and Professor R. S. PHILLIPS of Stanford University for the chance to stay in their universities in 1962, which enabled him to polish the greater part of the manuscript of this book, availing himself of their valuable advice. Professor S. ITO and Dr. H. KOMATSU of the University of Tokyo kindly assisted the author in reading various parts

Preface

of the galley proof, correcting errors and improving the presentation. To all of them, the author expresses his warmest gratitude.

Thanks are also due to Professor F. K. SCHMIDT of Heidelberg University and to Professor T. KATO of the University of California at Berkeley who constantly encouraged the author to write up the present book. Finally, the author wishes to express his appreciation to Springer-Verlag for their most efficient handling of the publication of this book.

Tokyo, September 1964

KÔSAKU YOSIDA

Preface to the Second Edition

In the preparation of this edition, the author is indebted to Mr. FLORET of Heidelberg who kindly did the task of enlarging the Index to make the book more useful. The errors in the second printing are corrected thanks to the remarks of many friends. In order to make the book more up-to-date, Section 4 of Chapter XIV has been rewritten entirely for this new edition.

Tokyo, September 1967

KÔSAKU YOSIDA

Preface to the Third Edition

A new Section (9. Abstract Potential Operators and Semi-groups) pertaining to G. HUNT's theory of potentials is inserted in Chapter XIII of this edition. The errors in the second edition are corrected thanks to kind remarks of many friends, especially of Mr. KLAUS-DIETER BIERSTEDT.

Kyoto, April 1971

KÔSAKU YOSIDA

Preface to the Fourth Edition

Two new Sections "6. Non-linear Evolution Equations 1 (The Kōmura-Kato Approach)" and "7. Non-linear Evolution Equations 2 (The Approach Through The Crandall-Liggett Convergence Theorem)" are added to the last Chapter XIV of this edition. The author is grateful to Professor Y. Kōmura for his careful reading of the manuscript.

Tokyo, April 1974

KÔSAKU YOSIDA

Preface to the Fifth Edition

Taking advantage of this opportunity, supplementary notes are added at the end of this new edition and additional references to books have been entered in the bibliography. The notes are divided into two categories. The first category comprises two topics: the one is concerned with the time reversibility of Markov processes with invariant measures, and the other is concerned with the uniqueness of the solution of time dependent linear evolution equations. The second category comprises those lists of recently published books dealing respectively with Sobolev Spaces, Trace Operators or Generalized Boundary Values, Distributions and Hyperfunctions, Contraction Operators in Hilbert Spaces, Choquet's Refinement of the Krein-Milman Theorem and Linear as well as Non-linear Evolution Equations.

A number of minor errors and a serious one on page 459 in the fourth edition have been corrected. The author wishes to thank many friends who kindly brought these errors to his attention.

Kamakura, August 1977

KÔSAKU YOSIDA

Preface to the Sixth Edition

Two major changes are made to this edition. The first is the rewriting of the Chapter VI,6 to give a simplified presentation of Mikusiński's Operational Calculus in such a way that this presentation does not appeal to Titchmarsh's theorem. The second is the rewriting of the Lemma together with its Proof in the Chapter XII,5 concerning the Representation of Vector Lattices. This rewriting is motivated by a letter of Professor E. Coimbra of Universidad Nova de Lisboa kindly suggesting the author's careless phrasing in the above Lemma of the preceding edition.

A number of misprints in the fifth edition have been corrected thanks to kind remarks of many friends.

Kamakura, June 1980

KÔSAKU YOSIDA

Contents

0. Preliminaries	1
1. Set Theory	1
2. Topological Spaces	3
3. Measure Spaces	15
4. Linear Spaces	20
I. Semi-norms	23
1. Semi-norms and Locally Convex Linear Topological Spaces .	23
2. Norms and Quasi-norms	30
3. Examples of Normed Linear Spaces	32
4. Examples of Quasi-normed Linear Spaces	38
5. Pre-Hilbert Spaces	39
6. Continuity of Linear Operators	42
7. Bounded Sets and Bornologic Spaces	44
8. Generalized Functions and Generalized Derivatives	46
9. B -spaces and F -spaces	52
10. The Completion	56
11. Factor Spaces of a B -space	59
12. The Partition of Unity	60
13. Generalized Functions with Compact Support	62
14. The Direct Product of Generalized Functions	65
II. Applications of the Baire-Hausdorff Theorem	68
1. The Uniform Boundedness Theorem and the Resonance Theorem	68
2. The Vitali-Hahn-Saks Theorem	70
3. The Termwise Differentiability of a Sequence of Generalized Functions	72
4. The Principle of the Condensation of Singularities	72
5. The Open Mapping Theorem	75
6. The Closed Graph Theorem	77
7. An Application of the Closed Graph Theorem (Hörmander's Theorem)	80
III. The Orthogonal Projection and F. Riesz' Representation Theo- rem	81
1. The Orthogonal Projection	81
2. "Nearly Orthogonal" Elements	84

Contents

3. The Ascoli-Arzelà Theorem	85
4. The Orthogonal Base, Bessel's Inequality and Parseval's Relation	86
5. E. Schmidt's Orthogonalization	88
6. F. Riesz' Representation Theorem	90
7. The Lax-Milgram Theorem	92
8. A Proof of the Lebesgue-Nikodym Theorem	93
9. The Aronszajn-Bergman Reproducing Kernel	95
10. The Negative Norm of P. LAX	98
11. Local Structures of Generalized Functions	100
 IV. The Hahn-Banach Theorems	 102
1. The Hahn-Banach Extension Theorem in Real Linear Spaces	102
2. The Generalized Limit	103
3. Locally Convex, Complete Linear Topological Spaces . . .	104
4. The Hahn-Banach Extension Theorem in Complex Linear Spaces	105
5. The Hahn-Banach Extension Theorem in Normed Linear Spaces	106
6. The Existence of Non-trivial Continuous Linear Functionals	107
7. Topologies of Linear Maps	110
8. The Embedding of X in its Bidual Space X''	112
9. Examples of Dual Spaces	114
 V. Strong Convergence and Weak Convergence	 119
1. The Weak Convergence and The Weak* Convergence . . .	120
2. The Local Sequential Weak Compactness of Reflexive B -spaces. The Uniform Convexity	126
3. Dunford's Theorem and The Gelfand-Mazur Theorem . . .	128
4. The Weak and Strong Measurability. Pettis' Theorem . . .	130
5. Bochner's Integral	132
 Appendix to Chapter V. Weak Topologies and Duality in Locally Convex Linear Topological Spaces	 136
1. Polar Sets	136
2. Barrel Spaces	138
3. Semi-reflexivity and Reflexivity	139
4. The Eberlein-Smul'yan Theorem	141
 VI. Fourier Transform and Differential Equations	 145
1. The Fourier Transform of Rapidly Decreasing Functions .	146
2. The Fourier Transform of Tempered Distributions.	149
3. Convolutions	156
4. The Paley-Wiener Theorems. The One-sided Laplace Transform	161
5. Titchmarsh's Theorem	166
6. Mikusiński's Operational Calculus	169
7. Sobolev's Lemma	173

Contents

8. Gårding's Inequality	175
9. Friedrichs' Theorem	177
10. The Malgrange-Ehrenpreis Theorem	182
11. Differential Operators with Uniform Strength.	188
12. The Hypoellipticity (Hörmander's Theorem)	189
 VII. Dual Operators	 193
1. Dual Operators	193
2. Adjoint Operators	195
3. Symmetric Operators and Self-adjoint Operators	197
4. Unitary Operators. The Cayley Transform	202
5. The Closed Range Theorem	205
 VIII. Resolvent and Spectrum.	 209
1. The Resolvent and Spectrum	209
2. The Resolvent Equation and Spectral Radius	211
3. The Mean Ergodic Theorem	213
4. Ergodic Theorems of the Hille Type Concerning Pseudo-resolvents	215
5. The Mean Value of an Almost Periodic Function	218
6. The Resolvent of a Dual Operator.	224
7. Dunford's Integral	225
8. The Isolated Singularities of a Resolvent	228
 IX. Analytical Theory of Semi-groups.	 231
1. The Semi-group of Class (C_0)	232
2. The Equi-continuous Semi-group of Class (C_0) in Locally Convex Spaces. Examples of Semi-groups	234
3. The Infinitesimal Generator of an Equi-continuous Semi-group of Class (C_0)	237
4. The Resolvent of the Infinitesimal Generator A	240
5. Examples of Infinitesimal Generators	242
6. The Exponential of a Continuous Linear Operator whose Powers are Equi-continuous	244
7. The Representation and the Characterization of Equi-continuous Semi-groups of Class (C_0) in Terms of the Corresponding Infinitesimal Generators	246
8. Contraction Semi-groups and Dissipative Operators	250
9. Equi-continuous Groups of Class (C_0) . Stone's Theorem	251
10. Holomorphic Semi-groups	254
11. Fractional Powers of Closed Operators	259
12. The Convergence of Semi-groups. The Trotter-Kato Theorem	269
13. Dual Semi-groups. Phillips' Theorem	272
 X. Compact Operators	 274
1. Compact Sets in B -spaces	274
2. Compact Operators and Nuclear Operators	277

Contents

3. The Rellich-Gårding Theorem	281
4. Schauder's Theorem	282
5. The Riesz-Schauder Theory	283
6. Dirichlet's Problem	286
Appendix to Chapter X. The Nuclear Space of A. GROTHENDIECK	289
 XI. Normed Rings and Spectral Representation	 294
1. Maximal Ideals of a Normed Ring	295
2. The Radical. The Semi-simplicity	298
3. The Spectral Resolution of Bounded Normal Operators	302
4. The Spectral Resolution of a Unitary Operator	306
5. The Resolution of the Identity	309
6. The Spectral Resolution of a Self-adjoint Operator	313
7. Real Operators and Semi-bounded Operators. Friedrichs' Theorem	316
8. The Spectrum of a Self-adjoint Operator. Rayleigh's Principle and the Krylov-Weinstein Theorem. The Multiplicity of the Spectrum	319
9. The General Expansion Theorem. A Condition for the Absence of the Continuous Spectrum	323
10. The Peter-Weyl-Neumann Theorem	326
11. Tannaka's Duality Theorem for Non-commutative Compact Groups	332
12. Functions of a Self-adjoint Operator	338
13. Stone's Theorem and Bochner's Theorem	345
14. A Canonical Form of a Self-adjoint Operator with Simple Spectrum	347
15. The Defect Indices of a Symmetric Operator. The Generalized Resolution of the Identity	349
16. The Group-ring L^1 and Wiener's Tauberian Theorem	354
 XII. Other Representation Theorems in Linear Spaces	 362
1. Extremal Points. The Krein-Milman Theorem	362
2. Vector Lattices	364
3. B -lattices and F -lattices	369
4. A Convergence Theorem of BANACH	370
5. The Representation of a Vector Lattice as Point Functions	372
6. The Representation of a Vector Lattice as Set Functions	375
 XIII. Ergodic Theory and Diffusion Theory	 379
1. The Markov Process with an Invariant Measure	379
2. An Individual Ergodic Theorem and Its Applications	383
3. The Ergodic Hypothesis and the H -theorem	389
4. The Ergodic Decomposition of a Markov Process with a Locally Compact Phase Space	393
5. The Brownian Motion on a Homogeneous Riemannian Space	398
6. The Generalized Laplacian of W. FELLER	403
7. An Extension of the Diffusion Operator	408
8. Markov Processes and Potentials	410
9. Abstract Potential Operators and Semi-groups	411

Contents

XIV. The Integration of the Equation of Evolution	418
1. Integration of Diffusion Equations in $L^2(R^m)$	419
2. Integration of Diffusion Equations in a Compact Riemannian Space	425
3. Integration of Wave Equations in a Euclidean Space R^m	427
4. Integration of Temporally Inhomogeneous Equations of Evolution in a B -space	430
5. The Method of TANABE and SOBOLEVSKI	438
6. Non-linear Evolution Equations 1 (The Kōmura-Kato Approach)	445
7. Non-linear Evolution Equations 2 (The Approach through the Crandall-Liggett Convergence Theorem)	454
Supplementary Notes	466
Bibliography	469
Index	487
Notation of Spaces	501

0. Preliminaries

It is the purpose of this chapter to explain certain notions and theorems used throughout the present book. These are related to *Set Theory*, *Topological Spaces*, *Measure Spaces* and *Linear Spaces*.

1. Set Theory

Sets. $x \in X$ means that x is a *member* or *element* of the set X ; $x \notin X$ means that x is not a member of the set X . We denote the set consisting of all x possessing the property P by $\{x; P\}$. Thus $\{y; y = x\}$ is the set $\{x\}$ consisting of a single element x . The *void set* is the set with no members, and will be denoted by \emptyset . If every element of a set X is also an element of a set Y , then X is said to be a *subset* of Y and this fact will be denoted by $X \subseteq Y$, or $Y \supseteq X$. If \mathfrak{X} is a set whose elements are sets X , then the set of all x such that $x \in X$ for some $X \in \mathfrak{X}$ is called the *union* of sets X in \mathfrak{X} ; this union will be denoted by $\bigcup_{X \in \mathfrak{X}} X$. The *intersection* of the sets X in \mathfrak{X} is the set of all x which are elements of every $X \in \mathfrak{X}$; this intersection will be denoted by $\bigcap_{X \in \mathfrak{X}} X$. Two sets are *disjoint* if their intersection is void. A family of sets is disjoint if every pair of distinct sets in the family is disjoint. If a sequence $\{X_n\}_{n=1,2,\dots}$ of sets is a disjoint family, then the union $\bigcup_{n=1}^{\infty} X_n$ may be written in the form of a sum $\sum_{n=1}^{\infty} X_n$.

Mappings. The term *mapping*, *function* and *transformation* will be used synonymously. The symbol $f: X \rightarrow Y$ will mean that f is a single-valued function whose *domain* is X and whose *range* is contained in Y ; for every $x \in X$, the function f assigns a uniquely determined element $f(x) = y \in Y$. For two mappings $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, we can define their *composite mapping* $gf: X \rightarrow Z$ by $(gf)(x) = g(f(x))$. The symbol $f(M)$ denotes the set $\{f(x); x \in M\}$ and $f(M)$ is called the *image* of M under the mapping f . The symbol $f^{-1}(N)$ denotes the set $\{x; f(x) \in N\}$ and $f^{-1}(N)$ is called the *inverse image* of N under the mapping f . It is clear that

$$Y_1 = f(f^{-1}(Y_1)) \text{ for all } Y_1 \subseteq f(X), \text{ and } X_1 \subseteq f^{-1}(f(X_1)) \text{ for all } X_1 \subseteq X.$$

If $f: X \rightarrow Y$, and for each $y \in f(X)$ there is only one $x \in X$ with $f(x) = y$, then f is said to have an *inverse* (mapping) or to be *one-to-one*. The inverse mapping then has the domain $f(X)$ and range X ; it is defined by the equation $x = f^{-1}(y) = f^{-1}(\{y\})$.

The domain and the range of a mapping f will be denoted by $D(f)$ and $R(f)$, respectively. Thus, if f has an inverse then

$$f^{-1}(f(x)) = x \text{ for all } x \in D(f), \text{ and } f(f^{-1}(y)) = y \text{ for all } y \in R(f).$$

The function f is said to map X *onto* Y if $f(X) = Y$ and *into* Y if $f(X) \subseteq Y$. The function f is said to be an *extension* of the function g and g a *restriction* of f if $D(f)$ contains $D(g)$, and $f(x) = g(x)$ for all x in $D(g)$.

Zorn's Lemma

Definition. Let P be a set of elements a, b, \dots . Suppose there is a binary relation defined between certain pairs (a, b) of elements of P , expressed by $a < b$, with the properties:

$$\begin{cases} a < a, \\ \text{if } a < b \text{ and } b < a, \text{ then } a = b, \\ \text{if } a < b \text{ and } b < c, \text{ then } a < c \text{ (transitivity).} \end{cases}$$

Then P is said to be *partially ordered* (or *semi-ordered*) by the relation $<$.

Examples. If P is the set of all subsets of a given set X , then the set inclusion relation ($A \subseteq B$) gives a partial ordering of P . The set of all complex numbers $z = x + iy$, $w = u + iv, \dots$ is partially ordered by defining $z < w$ to mean $x \leq u$ and $y \leq v$.

Definition. Let P be a partially ordered set with elements a, b, \dots . If $a < c$ and $b < c$, we call c an *upper bound* for a and b . If furthermore $c < d$ whenever d is an upper bound for a and b , we call c the *least upper bound* or the *supremum* of a and b , and write $c = \sup(a, b)$ or $a \vee b$. This element of P is unique if it exists. In a similar way we define the *greatest lower bound* or the *infimum* of a and b , and denote it by $\inf(a, b)$ or $a \wedge b$. If $a \vee b$ and $a \wedge b$ exist for every pair (a, b) in a partially ordered set P , P is called a *lattice*.

Example. The totality of subsets M of a fixed set B is a lattice by the partial ordering $M_1 < M_2$ defined by the set inclusion relation $M_1 \subseteq M_2$.

Definition. A partially ordered set P is said to be *linearly ordered* (or *totally ordered*) if for every pair (a, b) in P , either $a < b$ or $b < a$ holds. A subset of a partially ordered set is itself partially ordered by the relation which partially orders P ; the subset might turn out to be linearly ordered by this relation. If P is partially ordered and S is a subset of P , an $m \in P$ is called an *upper bound* of S if $s < m$ for every $s \in S$. An $m \in P$ is said to be *maximal* if $p \in P$ and $m < p$ together imply $m = p$.

Zorn's Lemma. Let P be a non-empty partially ordered set with the property that every linearly ordered subset of P has an upper bound in P . Then P contains at least one maximal element.

It is known that Zorn's lemma is equivalent to Zermelo's axiom of choice in set theory.

2. Topological Spaces

Open Sets and Closed Sets

Definition. A system τ of subsets of a set X defines a *topology* in X if τ contains the void set, the set X itself, the union of every one of its subsystems, and the intersection of every one of its finite subsystems. The sets in τ are called the *open sets* of the *topological space* (X, τ) ; we shall often omit τ and refer to X as a topological space. Unless otherwise stated, we shall assume that a topological space X satisfies *Hausdorff's axiom of separation*:

For every pair (x_1, x_2) of distinct points x_1, x_2 of X , there exist disjoint open sets G_1, G_2 such that $x_1 \in G_1, x_2 \in G_2$.

A *neighbourhood* of the point x of X is a set containing an open set which contains x . A neighbourhood of the subset M of X is a set which is a neighbourhood of every point of M . A point x of X is an *accumulation point* or *limit point* of a subset M of X if every neighbourhood of x contains at least one point $m \in M$ different from x .

Definition. Any subset M of a topological space X becomes a topological space by calling "open" the subsets of M which are of the form $M \cap G$ where G 's are open sets of X . The induced topology of M is called the *relative topology* of M as a subset of the topological space X .

Definition. A set M of a topological space X is *closed* if it contains all its accumulation points. It is easy to see that M is closed iff¹ its complement $M^c = X - M$ is open. Here $A - B$ denotes the totality of points $x \in A$ not contained in B . If $M \subseteq X$, the intersection of all closed subsets of X which contain M is called the *closure* of M and will be denoted by M^a (the superscript "a" stands for the first letter of the German: abgeschlossene Hülle).

Clearly M^a is closed and $M \subseteq M^a$; it is easy to see that $M = M^a$ iff M is closed.

Metric Spaces

Definition. If X, Y are sets, we denote by $X \times Y$ the set of all ordered pairs (x, y) where $x \in X$ and $y \in Y$; $X \times Y$ will be called the *Cartesian product* of X and Y . X is called a *metric space* if there is defined a func-

¹ iff is the abbreviation for "if and only if".

tion d with domain $X \times X$ and range in the real number field R^1 such that

$$\begin{cases} d(x_1, x_2) \geq 0 \text{ and } d(x_1, x_2) = 0 \text{ iff } x_1 = x_2, \\ d(x_1, x_2) = d(x_2, x_1), \\ d(x_1, x_3) \leq d(x_1, x_2) + d(x_2, x_3) \text{ (the triangle inequality).} \end{cases}$$

d is called the *metric* or the *distance function* of X . With each point x_0 in a metric space X and each positive number r , we associate the set $S(x_0; r) = \{x \in X; d(x, x_0) < r\}$ and call it the *open sphere* with centre x_0 and radius r . Let us call "open" the set M of a metric space X iff, for every point $x_0 \in M$, M contains a sphere with centre x_0 . Then the totality of such "open" sets satisfies the axiom of open sets in the definition of the topological space.

Hence a metric space X is a topological space. It is easy to see that a point x_0 of X is an accumulation point of M iff, to every $\varepsilon > 0$, there exists at least one point $m \neq x_0$ of M such that $d(m, x_0) < \varepsilon$. The n -dimensional Euclidean space R^n is a metric space by

$$d(x, y) = \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{1/2}, \text{ where } x = (x_1, \dots, x_n) \text{ and } y = (y_1, \dots, y_n).$$

Continuous Mappings

Definition. Let $f: X \rightarrow Y$ be a mapping defined on a topological space X into a topological space Y . f is called *continuous at a point* $x_0 \in X$ if to every neighbourhood U of $f(x_0)$ there corresponds a neighbourhood V of x_0 such that $f(V) \subseteq U$. The mapping f is said to be *continuous* if it is continuous at every point of its domain $D(f) = X$.

Theorem. Let X, Y be topological spaces and f a mapping defined on X into Y . Then f is continuous iff the inverse image under f of every open set of Y is an open set of X .

Proof. If f is continuous and U an open set of Y , then $V = f^{-1}(U)$ is a neighbourhood of every point $x_0 \in X$ such that $f(x_0) \in U$, that is, V is a neighbourhood of every point x_0 of V . Thus V is an open set of X . Let, conversely, for every open set $U \ni f(x_0)$ of Y , the set $V = f^{-1}(U)$ be an open set of X . Then, by the definition, f is continuous at $x_0 \in X$.

Compactness

Definition. A system of sets $G_\alpha, \alpha \in A$, is called a *covering* of the set X if X is contained as a subset of the union $\bigcup_{\alpha \in A} G_\alpha$. A subset M of a topological space X is called *compact* if every system of open sets of X which covers M contains a finite subsystem also covering M .

In view of the preceding theorem, a *continuous image of a compact set is also compact*.

Proposition 1. Compact subsets of a topological space are necessarily closed.

Proof. Let there be an accumulation point x_0 of a compact set M of a topological space X such that $x_0 \notin M$. By Hausdorff's axiom of separation, there exist, for any point $m \in M$, disjoint open sets G_{m, x_0} and $G_{x_0, m}$ of X such that $m \in G_{m, x_0}$, $x_0 \in G_{x_0, m}$. The system $\{G_{m, x_0}; m \in M\}$ surely covers M . By the compactness of M , there exists a finite subsystem $\{G_{m_i, x_0}; i = 1, 2, \dots, n\}$ which covers M . Then $\bigcap_{i=1}^n G_{x_0, m_i}$ does not intersect M . But, since x_0 is an accumulation point of M , the open set $\bigcap_{i=1}^n G_{x_0, m_i} \ni x_0$ must contain a point $m \in M$ distinct from x_0 . This is a contradiction, and M must be closed.

Proposition 2. A closed subset M_1 of a compact set M of a topological space X is compact.

Proof. Let $\{G_\alpha\}$ be any system of open sets of X which covers M_1 . M_1 being closed, $M_1^c = X - M_1$ is an open set of X . Since $M_1 \subseteq M$, the system of open sets $\{G_\alpha\}$ plus M_1^c covers M , and since M is compact, a properly chosen finite subsystem $\{G_{\alpha_i}; i = 1, 2, \dots, n\}$ plus M_1^c surely covers M . Thus $\{G_{\alpha_i}; i = 1, 2, \dots, n\}$ covers M_1 .

Definition. A subset of a topological space is called *relatively compact* if its closure is compact. A topological space is said to be *locally compact* if each point of the space has a compact neighbourhood.

Theorem. Any locally compact space X can be embedded in another compact space Y , having just one more point than X , in such a way that the relative topology of X as a subset of Y is just the original topology of X . This Y is called a *one point compactification* of X .

Proof. Let y be any element distinct from the points of X . Let $\{U\}$ be the class of all open sets in X such that $U^c = X - U$ is compact. We remark that X itself $\in \{U\}$. Let Y be the set consisting of the points of X and the point y . A set in Y will be called open if either (i) it does not contain y and is open as a subset of X , or (ii) it does contain y and its intersection with X is a member of $\{U\}$. It is easy to see that Y thus obtained is a topological space, and that the relative topology of X coincides with its original topology.

Suppose $\{V\}$ be a family of open sets which covers Y . Then there must be some member of $\{V\}$ of the form $U_0 \cup \{y\}$, where $U_0 \in \{U\}$. By the definition of $\{U\}$, U_0^c is compact as a subset of X . It is covered by the system of sets $V \cap X$ with $V \in \{V\}$. Thus some finite subsystem: $V_1 \cap X, V_2 \cap X, \dots, V_n \cap X$ covers U_0^c . Consequently, V_1, V_2, \dots, V_n and $U_0 \cup \{y\}$ cover Y , proving that Y is compact.

Tychonov's Theorem

Definition. Corresponding to each α of an index set A , let there be given a topological space X_α . The Cartesian product $\prod_{\alpha \in A} X_\alpha$ is, by defini-