

# 欧几里得原理十三本书

第2卷

## The Thirteen Books of Euclid's Elements

II

[古希腊] 欧几里得 著

世界图书出版公司  
[www.wpcbj.com.cn](http://www.wpcbj.com.cn)

THE THIRTEEN BOOKS  
OF  
EUCLID'S ELEMENTS

TRANSLATED FROM THE TEXT OF HEIBERG  
WITH INTRODUCTION AND COMMENTARY

BY  
T. L. HEATH, C.B., Sc.D.,  
SOMETIME FELLOW OF TRINITY COLLEGE, CAMBRIDGE



VOLUME II  
BOOKS III—IX

图书在版编目 (C I P) 数据

欧几里得原理十三本书=The Thirteen Books of Euclid's Elements. 第2卷: 英文 / (古希腊) 欧几里得 (Euclid) 著. —影印本. —北京: 世界图书出版公司北京公司, 2015. 2  
ISBN 978-7-5100-9409-5

I. ①欧… II. ①欧… III. ①欧氏几何—英文 IV. ①O184

中国版本图书馆 CIP 数据核字 (2015) 第 037269 号

欧几里得原理十三本书 第2卷

The Thirteen Books of Euclid's Elements II

---

作 者: [古希腊] 欧几里得 (Euclid)

策 划: 杜玉娟

出版统筹: 李艺琳

责任编辑: 张明德

封面设计: 张瑞玲

出版发行: 世界图书出版公司北京公司 <http://www.wpcbj.com.cn>

地 址: 北京市朝内大街 137 号 (邮编 100010 电话 010-64077922)

销 售: 各地新华书店及外文书店

印 刷: 北京财经印刷厂

开 本: 170mm×230mm

印 张: 27.75

字 数: 132 千

版 次: 2015 年 5 月第 1 版 2015 年 5 月第 1 次印刷

书 号: 978-7-5100-9409-5

定 价: 60.00 (元)

---

版权所有 翻印必究

(如发现印装质量问题, 请与本公司联系调换)





# CONTENTS OF VOLUME II

	PAGE
BOOK III. DEFINITIONS . . . . .	I
PROPOSITIONS . . . . .	6
BOOK IV. DEFINITIONS . . . . .	78
PROPOSITIONS . . . . .	80
BOOK V. INTRODUCTORY NOTE . . . . .	112
DEFINITIONS . . . . .	113
PROPOSITIONS . . . . .	138
BOOK VI. INTRODUCTORY NOTE . . . . .	187
DEFINITIONS . . . . .	188
PROPOSITIONS . . . . .	191
BOOK VII. DEFINITIONS . . . . .	277
PROPOSITIONS . . . . .	296
BOOK VIII. . . . .	345
BOOK IX. . . . .	384
GREEK INDEX TO VOL. II. . . . .	427
ENGLISH INDEX TO VOL. II. . . . .	431

## BOOK III.

### DEFINITIONS.

1. **Equal circles** are those the diameters of which are equal, or the radii of which are equal.
2. A straight line is said to **touch a circle** which, meeting the circle and being produced, does not cut the circle.
3. **Circles** are said to **touch one another** which, meeting one another, do not cut one another.
4. In a circle straight lines are said to be **equally distant from the centre** when the perpendiculars drawn to them from the centre are equal.
5. And that straight line is said to be **at a greater distance** on which the greater perpendicular falls.
6. A **segment of a circle** is the figure contained by a straight line and a circumference of a circle.
7. An **angle of a segment** is that contained by a straight line and a circumference of a circle.
8. An **angle in a segment** is the angle which, when a point is taken on the circumference of the segment and straight lines are joined from it to the extremities of the straight line which is the **base of the segment**, is contained by the straight lines so joined.
9. And, when the straight lines containing the angle cut off a circumference, the angle is said to **stand upon** that circumference.

10. A sector of a circle is the figure which, when an angle is constructed at the centre of the circle, is contained by the straight lines containing the angle and the circumference cut off by them.

11. Similar segments of circles are those which admit equal angles, or in which the angles are equal to one another.

#### DEFINITION 1.

ἴσοι κύκλοι εἰσίν, ὧν αἱ διάμετροι ἴσαι εἰσίν, ἢ ὧν αἱ ἐκ τῶν κέντρων ἴσαι εἰσίν.

Many editors have held that this should not have been included among definitions. Some, e.g. Tartaglia, would call it a *postulate*; others, e.g. Borelli and Playfair, would call it an *axiom*; others again, as Billingsley and Clavius, while admitting it as a *definition*, add explanations based on the mode of constructing a circle; Simson and Pfeiderer hold that it is a *theorem*. I think however that Euclid would have maintained that it is a definition in the proper sense of the term; and certainly it satisfies Aristotle's requirement that a "definitional statement" (ὀριστικὸς λόγος) should not only state the *fact* (τὸ ὅτι) but should indicate the *cause* as well (*De anima* II. 2, 413 a 13). The equality of circles with equal radii can of course be proved by superposition, but, as we have seen, Euclid avoided this method wherever he could, and there is nothing technically wrong in saying "By *equal circles* I mean circles with equal radii." No flaw is thereby introduced into the system of the *Elements*; for the definition could only be objected to if it could be proved that the equality predicated of the two circles in the definition was not the same thing as the equality predicated of other equal figures in the *Elements* on the basis of the Congruence-Axiom, and, needless to say, this cannot be proved because it is not true. The *existence* of equal circles (in the sense of the definition) follows from the existence of equal straight lines and I. Post. 3.

The Greeks had no distinct word for *radius*, which is with them, as here, the (straight line drawn) from the centre ἢ ἐκ τοῦ κέντρου (εὐθεΐα); and so definitely was the expression appropriated to the radius that ἐκ τοῦ κέντρου was used without the article as a predicate, just as if it were one word. Thus, e.g., in III. 1 ἐκ κέντρου γάρ means "for they are radii": cf. Archimedes, *On the Sphere and Cylinder* II. 2, ἢ BE ἐκ τοῦ κέντρου ἐστὶ τοῦ...κύκλου, BE is a radius of the circle.

#### DEFINITION 2.

Εὐθεΐα κύκλον ἐφάπτεσθαι λέγεται, ἥτις ἀπομένῃ τοῦ κύκλου καὶ ἐκβαλλομένη οὐ τέμνει τὸν κύκλον.

Euclid's phraseology here shows the regular distinction between ἀπτεσθαι and its compound ἐφάπτεσθαι, the former meaning "to meet" and the latter "to touch." The distinction was generally observed by Greek geometers from Euclid onwards. There are however exceptions so far as ἀπτεσθαι is concerned; thus it means "to touch" in Eucl. IV. Def. 5 and sometimes in Archimedes. On the other hand, ἐφάπτεσθαι is used by Aristotle in certain



cases where the orthodox geometrical term would be ἀπτεσθαι. Thus in *Meteorologica* III. 5 (376 b 9) he says a certain circle *will pass through all the angles* (ἀπασῶν ἐφάψεται τῶν γωνιῶν), and (376 a 6) M *will lie on a given* (circular) *circumference* (δεδομένης περιφερείας ἐφάψεται τὸ M). We shall find ἀπτεσθαι used in these senses in Book IV. Deff. 2, 6 and Deff. 1, 3 respectively. The latter of the two expressions quoted from Aristotle means that *the locus of M is a given circle*, just as in Pappus ἀψεται τὸ σημεῖον θέσει δεδομένης εὐθείας means that *the locus of the point is a straight line given in position*.

## DEFINITION 3.

Κύκλοι ἐφάπτεσθαι ἀλλήλων λέγονται οἷτινες ἀπτόμενοι ἀλλήλων οὐ τέμνουσιν ἀλλήλους.

Todhunter remarks that different opinions have been held as to what is, or should be, included in this definition, one opinion being that it only means that the circles do not cut in the neighbourhood of the point of contact, and that it must be shown that they do not cut elsewhere, while another opinion is that the definition means that the circles do not cut at all. Todhunter thinks the latter opinion correct. I do not think this is proved; and I prefer to read the definition as meaning simply that the circles meet at a point but do not cut *at that point*. I think this interpretation preferable for the reason that, although Euclid does practically assume in III. 11—13, without stating, the theorem that circles touching at one point do not intersect anywhere else, he has given us, before reaching that point in the Book, means for proving for ourselves the truth of that statement. In particular, he has given us the propositions III. 7, 8 which, taken as a whole, give us more information as to the general nature of a circle than any other propositions that have preceded, and which can be used, as will be seen in the sequel, to solve any doubts arising out of Euclid's unproved assumptions. Now, as a matter of fact, the propositions are not used in any of the genuine proofs of the theorems in Book III.; III. 8 is required for the second proof of III. 9 which Simson selected in preference to the first proof, but the first proof only is regarded by Heiberg as genuine. Hence it would not be easy to account for the appearance of III. 7, 8 at all unless as affording means of answering possible *objections* (cf. Proclus' explanation of Euclid's reason for inserting the second part of I. 5).

*External* and *internal* contact are not distinguished in Euclid until III. 11, 12, though the *figure* of III. 6 (not the *enunciation* in the original text) represents the case of internal contact only. But the definition of touching circles here given must be taken to imply so much about *internal* and *external* contact respectively as that (a) a circle touching another internally must, immediately before "meeting" it, have passed through points *within* the circle that it touches, and (b) a circle touching another externally must, immediately before meeting it, have passed through points *outside* the circle which it touches. These facts must indeed be admitted if *internal* and *external* are to have any meaning at all in this connexion, and they constitute a minimum admission necessary to the proof of III. 6.

## DEFINITION 4.

Ἐν κύκλῳ ἴσον ἀπέχειν ἀπὸ τοῦ κέντρου εὐθείαι λέγονται, ὅταν αἱ ἀπὸ τοῦ κέντρου ἐπ' αὐτὰς κάβητοι ἀγόμεναι ἴσαι ὦσιν.

## DEFINITION 5.

Μεῖζον δὲ ἀπέχειν λέγεται, ἐφ' ἣν ἡ μείζων κάθετος πίπτει.

## DEFINITION 6.

Τμήμα κύκλου ἐστὶ τὸ περιεχόμενον σχῆμα ὑπὸ τε εὐθείας καὶ κύκλου περιφερείας.

## DEFINITION 7.

Τμήματος δὲ γωνία ἐστὶν ἡ περιεχομένη ὑπὸ τε εὐθείας καὶ κύκλου περιφερείας.

This definition is only interesting historically. The *angle of a segment*, being the “angle” formed by a straight line and a “circumference,” is of the kind described by Proclus as “mixed.” A particular “angle” of this sort is the “*angle of a semicircle*,” which we meet with again in III. 16, along with the so-called “horn-like angle” (κερατοειδής), the supposed “angle” between a tangent to a circle and the circle itself. The “angle of a semicircle” occurs once in Pappus (VII. p. 670, 19), but it there means scarcely more than the *corner* of a semicircle regarded as a point to which a straight line is directed. Heron does not give the definition of the *angle of a segment*, and we may conclude that the mention of it and of the *angle of a semicircle* in Euclid is a survival from earlier text-books rather than an indication that Euclid considered either to be of importance in elementary geometry (cf. the note on III. 16 below).

We have however, in the note on I. 5 above (Vol. I. pp. 252—3), seen evidence that the *angle of a segment* had played some part in geometrical proofs up to Euclid's time. It would appear from the passage of Aristotle there quoted (*Anal. prior.* I. 24, 41 b 13 sqq.) that the theorem of I. 5 was, in the text-books immediately preceding Euclid, proved by means of the equality of the two “*angles of*” any one segment. This latter property must therefore have been regarded as more elementary (for whatever reason) than the theorem of I. 5; indeed the definition as given by Euclid practically implies the same thing, since it speaks of only *one* “angle of a segment,” namely “*the angle contained by a straight line and a circumference of a circle.*” Euclid abandoned the actual use of the “angle” in question, but no doubt thought it unnecessary to break with tradition so far as to strike the definition out also.

## DEFINITION 8.

Ἐν τμήματι δὲ γωνία ἐστίν, ὅταν ἐπὶ τῆς περιφερείας τοῦ τμήματος ληθῇ τι σημεῖον καὶ ἀπ' αὐτοῦ ἐπὶ τὰ πέρατα τῆς εὐθείας, ἥ ἐστι βᾶσις τοῦ τμήματος, ἐπιζευχθῶσιν εὐθείαι, ἡ περιεχομένη γωνία ὑπὸ τῶν ἐπιζευχθεισῶν εὐθειῶν.

## DEFINITION 9.

Ὅταν δὲ αἱ περιέχουσιν τὴν γωνίαν εὐθεῖαι ἀπολαμβάνωσιν τινὰ περιφέρειαν, ἐπ' ἐκείνης λέγεται βεβηκέναι ἡ γωνία.

## DEFINITION 10.

Τομεὺς δὲ κύκλου ἐστίν, ὅταν πρὸς τῷ κέντρῳ τοῦ κύκλου συσταθῇ γωνία, τὸ περιεχόμενον σχῆμα ὑπὸ τε τῶν τὴν γωνίαν περιεχουσῶν εὐθειῶν καὶ τῆς ἀπολαμβανομένης ὑπ' αὐτῶν περιφερείας.

A scholiast says that it was the *shoemaker's knife*, σκυτοτομικὸς τομεὺς, which suggested the name τομεὺς for a sector of a circle. The derivation of the name from a resemblance of shape is parallel to the use of ἀρβηλος (also a *shoemaker's knife*) to denote the well known figure of the Book of Lemmas partly attributed to Archimedes.


A wider definition of a sector than that given by Euclid is found in a Greek scholiast (Heiberg's Euclid, Vol. v. p. 260) and in an-Nairizi (ed. Curtze, p. 112). "There are two varieties of sectors; the one kind have the angular vertices at the centres, the other at the circumferences. Those others which have their vertices neither at the circumferences nor at the centres, but at some other points, are for that reason not called sectors but sector-like figures (τομοειδῆ σχήματα)." The exact agreement between the scholiast and an-Nairizi suggests that Heron was the authority for this explanation.

The *sector-like figure* bounded by an arc of a circle and two lines drawn from its extremities to meet at any point actually appears in Euclid's book *On divisions* (περὶ διαιρέσεων) discovered in an Arabic ms. and edited by Woepcke (cf. Vol. i. pp. 8—10 above). This treatise, alluded to by Proclus, had for its object the division of figures such as triangles, trapezia, quadrilaterals and circles, by means of straight lines, into parts equal or in given ratios. One proposition e.g. is, *To divide a triangle into two equal parts by a straight line passing through a given point on one side*. The proposition (28) in which the *quasi-sector* occurs is, *To divide such a figure by a straight line into two equal parts*. The solution in this case is given by Cantor (*Gesch. d. Math.* 1<sub>3</sub>, pp. 287—8).

If  $ABCD$  be the given figure,  $E$  the middle point of  $BD$  and  $EC$  at right angles to  $BD$ , the broken line  $AEC$  clearly divides the figure into two equal parts.

Join  $AC$ , and draw  $EF$  parallel to it meeting  $AB$  in  $F$ .

Join  $CF$ , when it is seen that  $CF$  divides the figure into two equal parts.



## DEFINITION 11.

Ὅμοια τμήματα κύκλων ἐστὶ τὰ δεχόμενα γωνίας ἴσας, ἥ ἐν οἷς αἱ γωνίαι ἴσαι ἀλλήλαις εἰσὶν.

De Morgan remarks that the use of the word *similar* in "similar segments" is an anticipation, and that similarity of *form* is meant. He adds that the definition is a theorem, or would be if "similar" had taken its final meaning.

# BOOK III. PROPOSITIONS.

## PROPOSITION I.

*To find the centre of a given circle.*

Let  $ABC$  be the given circle;  
thus it is required to find the centre of the circle  $ABC$ .

Let a straight line  $AB$  be drawn  
5 through it at random, and let it be bisected  
at the point  $D$ ;

from  $D$  let  $DC$  be drawn at right angles  
to  $AB$  and let it be drawn through to  $E$ ;  
let  $CE$  be bisected at  $F$ ;

10 I say that  $F$  is the centre of the circle  
 $ABC$ .

For suppose it is not, but, if possible,  
let  $G$  be the centre,  
and let  $GA, GD, GB$  be joined.

15 Then, since  $AD$  is equal to  $DB$ ,  
and  $DG$  is common,


the two sides  $AD, DG$  are equal to the two sides  
 $BD, DG$  respectively;

and the base  $GA$  is equal to the base  $GB$ , for they are  
20 radii;

therefore the angle  $ADG$  is equal to the angle  $GDB$ . [1. 8]

But, when a straight line set up on a straight line makes  
the adjacent angles equal to one another, each of the equal  
angles is right; [1. Def. 10]

25 therefore the angle  $GDB$  is right,



But the angle  $FDB$  is also right ;  
therefore the angle  $FDB$  is equal to the angle  $GDB$ , the  
greater to the less : which is impossible.

Therefore  $G$  is not the centre of the circle  $ABC$ .

30 Similarly we can prove that neither is any other point  
except  $F$ .

Therefore the point  $F$  is the centre of the circle  $ABC$ .

PORISM. From this it is manifest that, if in a circle a  
straight line cut a straight line into two equal parts and at  
35 right angles, the centre of the circle is on the cutting straight  
line.

Q. E. F.


12. For suppose it is not. This is expressed in the Greek by the two words  $M\eta\ \gamma\acute{\alpha}\rho$ , but such an elliptical phrase is impossible in English.

17. the two sides  $AD$ ,  $DG$  are equal to the two sides  $BD$ ,  $DG$  respectively. As before observed, Euclid is not always careful to put the equals in corresponding order. The text here has " $GD$ ,  $DB$ ."

Todhunter observes that, when, in the construction,  $DC$  is said to be *produced* to  $E$ , it is assumed that  $D$  is within the circle, a fact which Euclid first demonstrates in III. 2. This is no doubt true, although the word  $\deltaι\acute{\eta}\chi\theta\omega$ , "let it be *drawn through*," is used instead of  $\epsilon\kappa\beta\epsilon\beta\lambda\acute{\eta}\sigma\theta\omega$ , "let it be *produced*." And, although it is not necessary to assume that  $D$  is within the circle, it is necessary for the success of the construction that the straight line drawn through  $D$  at right angles to  $AB$  shall meet the circle in two points (and no more): an assumption which we are not entitled to make on the basis of what has gone before only.

Hence there is much to be said for the alternative procedure recommended by De Morgan as preferable to that of Euclid. De Morgan would first prove the fundamental theorem that "the line which bisects a chord perpendicularly must contain the centre," and then make III. 1, III. 25 and IV. 5 immediate corollaries of it. The fundamental theorem is a direct consequence of the theorem that, if  $P$  is any point equidistant from  $A$  and  $B$ , then  $P$  lies on the straight line bisecting  $AB$  perpendicularly. We then take any two chords  $AB$ ,  $AC$  of the given circle and draw  $DO$ ,  $EO$  bisecting them perpendicularly. Unless  $BA$ ,  $AC$  are in one straight line, the straight lines  $DO$ ,  $EO$  must meet in some point  $O$  (see note on IV. 5 for possible methods of proving this). And, since both  $DO$ ,  $EO$  must contain the centre,  $O$  must be the centre.

This method, which seems now to be generally preferred to Euclid's, has the advantage of showing that, in order to find the centre of a circle, it is sufficient to know three points on the circumference. If therefore two circles have three points in common, they must have the same centre and radius, so that two circles cannot have three points in common without coinciding entirely. Also, as indicated by De Morgan, the same construction enables us (1) to draw the complete circle of which a segment or arc only is given (III. 25), and (2) to circumscribe a circle to any triangle (IV. 5).



But, if the Greeks had used this construction for finding the centre of a circle, they would have considered it necessary to add a proof that no other point than that obtained by the construction can be the centre, as is clear both from the similar *reductio ad absurdum* in III. 1 and also from the fact that Euclid thinks it necessary to prove as a separate theorem (III. 9) that, if a point within a circle be such that three straight lines (at least) drawn from it to the circumference are equal, that point must be the centre. In fact, however, the proof amounts to no more than the remark that the two perpendicular bisectors can have no more than one point common.

And even in De Morgan's method there is a yet unproved assumption. In order that  $DO$ ,  $EO$  may meet, it is necessary that  $AB$ ,  $AC$  should not be in one straight line or, in other words, that  $BC$  should not pass through  $A$ . This results from III. 2, which therefore, strictly speaking, should precede.

To return to Euclid's own proposition III. 1, it will be observed that the demonstration only shows that the centre of the circle cannot lie on either side of  $CD$ , so that it must lie on  $CD$  or  $CD$  produced. It is however taken for granted rather than proved that the centre must be the middle point of  $CE$ . The proof of this by *reductio ad absurdum* is however so obvious as to be scarcely worth giving. The same consideration which would prove it may be used to show that a circle cannot have more than one centre, a proposition which, if thought necessary, may be added to III. 1 as a corollary.

Simson observed that the proof of III. 1 could not but be by *reductio ad absurdum*. At the beginning of Book III. we have nothing more to base the proof upon than the definition of a circle, and this cannot be made use of unless we assume some point to be the centre. We cannot however assume that the point found by the construction is the centre, because that is the thing to be proved. Nothing is therefore left to us but to assume that some other point is the centre and then to prove that, whatever other point is taken, an absurdity results; whence we can infer that the point found is the centre.

The Porism to III. 1 is inserted, as usual, parenthetically before the words  $\delta\pi\epsilon\rho\ \epsilon\delta\epsilon\iota\ \pi\omicron\iota\eta\sigma\alpha\iota$ , which of course refer to the problem itself.


### PROPOSITION 2.

*If on the circumference of a circle two points be taken at random, the straight line joining the points will fall within the circle.*

Let  $ABC$  be a circle, and let two points  $A$ ,  $B$  be taken at random on its circumference;

I say that the straight line joined from  $A$  to  $B$  will fall within the circle.

For suppose it does not, but, if possible, let it fall outside, as  $AEB$ ; let the centre of the circle  $ABC$  be taken [III. 1], and let it be  $D$ ; let  $DA$ ,  $DB$  be joined, and let  $DFE$  be drawn through.




Then, since  $DA$  is equal to  $DB$ ,  
 the angle  $DAE$  is also equal to the angle  $DBE$ . [I. 5]  
 And, since one side  $AEB$  of the triangle  $DAE$  is produced,  
 the angle  $DEB$  is greater than the angle  $DAE$ . [I. 16]  
 But the angle  $DAE$  is equal to the angle  $DBE$ ;  
 therefore the angle  $DEB$  is greater than the angle  $DBE$ .  
 And the greater angle is subtended by the greater side; [I. 19]  
 therefore  $DB$  is greater than  $DE$ .  
 But  $DB$  is equal to  $DF$ ;  
 therefore  $DF$  is greater than  $DE$ ,  
 the less than the greater: which is impossible.  
 Therefore the straight line joined from  $A$  to  $B$  will not  
 fall outside the circle.  
 Similarly we can prove that neither will it fall on the  
 circumference itself;  
 therefore it will fall within.  
 Therefore etc.

Q. E. D.

The *reductio ad absurdum* form of proof is not really necessary in this case, and it has the additional disadvantage that it requires the destruction of two hypotheses, namely that the chord is (1) outside, (2) on the circle. To prove the proposition directly, we have only to show that, if  $E$  be any point on the straight line  $AB$  between  $A$  and  $B$ ,  $DE$  is less than the radius of the circle. This may be done by the method shown above, under I. 24, for proving what is assumed in that proposition, namely that, in the figure of the proposition,  $F$  falls below  $EG$  if  $DE$  is not greater than  $DF$ . The assumption amounts to the following proposition, which De Morgan would make to precede I. 24: "Every straight line drawn from the vertex of a triangle to the base is less than the greater of the two sides, or than either if they be equal." The case here is that in which the two sides are equal; and, since the angle  $DAB$  is equal to the angle  $DBA$ , while the exterior angle  $DEA$  is greater than the interior and opposite angle  $DBA$ , it follows that the angle  $DEA$  is greater than the angle  $DAE$ , whence  $DE$  must be less than  $DA$  or  $DB$ .

Camerer points out that we may add to this proposition the further statement that all points on  $AB$  *produced* in either direction are outside the circle. This follows from the proposition (also proved by means of the theorems that the exterior angle of a triangle is greater than either of the interior and opposite angles and that the greater angle is subtended by the greater side) which De Morgan proposes to introduce after I. 21, namely,

"The perpendicular is the shortest straight line that can be drawn from a



given point to a given straight line, and of others that which is nearer to the perpendicular is less than the more remote, and the converse; also not more than two equal straight lines can be drawn from the point to the line, one on each side of the perpendicular."

The fact that not more than two equal straight lines can be drawn from a given point to a given straight line not passing through it is proved by Proclus on I. 16 (see the note to that proposition) and can alternatively be proved by means of I. 7, as shown above in the note on I. 12. It follows that

*A straight line cannot cut a circle in more than two points:*

a proposition which De Morgan would introduce here after III. 2. The proof given does not apply to a straight line *passing through the centre*; but that such a line only cuts the circle in two points is self-evident.

### PROPOSITION 3.

*If in a circle a straight line through the centre bisect a straight line not through the centre, it also cuts it at right angles; and if it cut it at right angles, it also bisects it.*

Let  $ABC$  be a circle, and in it let a straight line  $CD$   
5 through the centre bisect a straight line  
 $AB$  not through the centre at the point  
 $F$ ;

I say that it also cuts it at right angles.

For let the centre of the circle  $ABC$   
10 be taken, and let it be  $E$ ; let  $EA$ ,  $EB$   
be joined.

Then, since  $AF$  is equal to  $FB$ ,  
and  $FE$  is common,

two sides are equal to two sides;

15 and the base  $EA$  is equal to the base  $EB$ ;

therefore the angle  $AFE$  is equal to the angle  $BFE$ . [I. 8]


But, when a straight line set up on a straight line makes  
the adjacent angles equal to one another, each of the equal  
angles is right; [I. Def. 10]

20 therefore each of the angles  $AFE$ ,  $BFE$  is right.

Therefore  $CD$ , which is through the centre, and bisects  
 $AB$  which is not through the centre, also cuts it at right  
angles.

Again, let  $CD$  cut  $AB$  at right angles;

25 I say that it also bisects it, that is, that  $AF$  is equal to  $FB$ .





For, with the same construction,

since  $EA$  is equal to  $EB$ ,

the angle  $EAF$  is also equal to the angle  $EBF$ . [I. 5]

But the right angle  $AFE$  is equal to the right angle  $BFE$ ,  
 30 therefore  $EAF$ ,  $EBF$  are two triangles having two angles  
 equal to two angles and one side equal to one side, namely  
 $EF$ , which is common to them, and subtends one of the equal  
 angles;

therefore they will also have the remaining sides equal to  
 35 the remaining sides; [I. 26]

therefore  $AF$  is equal to  $FB$ .

Therefore etc.

Q. E. D.

26. with the same construction, τῶν αὐτῶν κατασκευασθέντων.

This proposition asserts the two *partial* converses (cf. note on I. 6) of the Porism to III. 1. De Morgan would place it next to III. 1.

#### PROPOSITION 4.


*If in a circle two straight lines cut one another which are not through the centre, they do not bisect one another.*

Let  $ABCD$  be a circle, and in it let the two straight lines  $AC$ ,  $BD$ , which are not through the centre, cut one another at  $E$ ;

I say that they do not bisect one another.

For, if possible, let them bisect one another, so that  $AE$  is equal to  $EC$ , and  $BE$  to  $ED$ ;

let the centre of the circle  $ABCD$  be taken [III. 1], and let it be  $F$ ; let  $FE$  be joined.



Then, since a straight line  $FE$  through the centre bisects a straight line  $AC$  not through the centre,

it also cuts it at right angles; [III. 3]

therefore the angle  $FEA$  is right.

Again, since a straight line  $FE$  bisects a straight line  $BD$ ,

it also cuts it at right angles; [III. 3]

therefore the angle  $FEB$  is right.