


Electronic and Optical Properties of Conjugated Polymers

Second Edition

WILLIAM BARFORD

*Physical and Theoretical Chemistry Laboratory
Department of Chemistry
University of Oxford
United Kingdom*



OXFORD
UNIVERSITY PRESS

OXFORD
UNIVERSITY PRESS

Great Clarendon Street, Oxford, OX2 6DP,
United Kingdom

Oxford University Press is a department of the University of Oxford.
It furthers the University's objective of excellence in research, scholarship,
and education by publishing worldwide. Oxford is a registered trade mark of
Oxford University Press in the UK and in certain other countries

© William Barford 2005, 2013

The moral rights of the author have been asserted

First Edition published in 2005

Second Edition published in 2013

Impression: 1

All rights reserved. No part of this publication may be reproduced, stored in
a retrieval system, or transmitted, in any form or by any means, without the
prior permission in writing of Oxford University Press, or as expressly permitted
by law, by licence or under terms agreed with the appropriate reprographics
rights organization. Enquiries concerning reproduction outside the scope of the
above should be sent to the Rights Department, Oxford University Press, at the
address above

You must not circulate this work in any other form
and you must impose this same condition on any acquirer

British Library Cataloguing in Publication Data

Data available

ISBN 978-0-19-967746-7

Printed and bound by
CPI Group (UK) Ltd, Croydon, CR0 4YY

Links to third party websites are provided by Oxford in good faith and
for information only. Oxford disclaims any responsibility for the materials
contained in any third party website referenced in this work.

INTERNATIONAL SERIES OF MONOGRAPHS
ON PHYSICS

SERIES EDITORS

J. BIRMAN
S. F. EDWARDS
R. FRIEND
M. REES
D. SHERRINGTON
G. VENEZIANO

CITY UNIVERSITY OF NEW YORK
UNIVERSITY OF CAMBRIDGE
UNIVERSITY OF CAMBRIDGE
UNIVERSITY OF CAMBRIDGE
UNIVERSITY OF OXFORD
CERN, GENEVA

INTERNATIONAL SERIES OF MONOGRAPHS ON PHYSICS

159. W. Barford: *Electronic and optical properties of conjugated polymers, Second edition*
158. F. Strocchi: *An introduction to non-perturbative foundations of quantum field theory*
157. K. H. Bennemann, J. B. Ketterson: *Novel superfluids, Volume 2*
156. K. H. Bennemann, J. B. Ketterson: *Novel superfluids, Volume 1*
155. C. Kiefer: *Quantum gravity, Third edition*
154. L. Mestel: *Stellar magnetism, Second edition*
153. R. A. Klemm: *Layered superconductors, Volume 1*
152. E. L. Wolf: *Principles of electron tunneling spectroscopy, Second edition*
151. R. Blinc: *Advanced ferroelectricity*
150. L. Berthier, G. Biroli, J.-P. Bouchaud, W. van Saarloos, L. Cipelletti: *Dynamical heterogeneities in glasses, colloids, and granular media*
149. J. Wesson: *Tokamaks, Fourth edition*
148. H. Asada, T. Futamase, P. Hogan: *Equations of motion in general relativity*
147. A. Yaouanc, P. Dalmas de Réotier: *Muon spin rotation, relaxation, and resonance*
146. B. McCoy: *Advanced statistical mechanics*
145. M. Bordag, G. L. Klimchitskaya, U. Mohideen, V. M. Mostepanenko: *Advances in the Casimir effect*
144. T. R. Field: *Electromagnetic scattering from random media*
143. W. Götze: *Complex dynamics of glass-forming liquids—a mode-coupling theory*
142. V. M. Agranovich: *Excitations in organic solids*
141. W. T. Grandy: *Entropy and the time evolution of macroscopic systems*
140. M. Alcubierre: *Introduction to 3 + 1 numerical relativity*
139. A. L. Ivanov, S. G. Tikhodeev: *Problems of condensed matter physics—quantum coherence phenomena in electron-hole and coupled matter-light systems*
138. I. M. Vardavas, F. W. Taylor: *Radiation and climate*
137. A. F. Borghesani: *Ions and electrons in liquid helium*
135. V. Fortov, I. Iakubov, A. Khrapak: *Physics of strongly coupled plasma*
134. G. Fredrickson: *The equilibrium theory of inhomogeneous polymers*
133. H. Suhl: *Relaxation processes in micromagnetics*
132. J. Terning: *Modern supersymmetry*
131. M. Mariño: *Chern-Simons theory, matrix models, and topological strings*
130. V. Gantmakher: *Electrons and disorder in solids*
129. W. Barford: *Electronic and optical properties of conjugated polymers*
128. R. E. Raab, O. L. de Lange: *Multipole theory in electromagnetism*
127. A. Larkin, A. Varlamov: *Theory of fluctuations in superconductors*
126. P. Goldbart, N. Goldenfeld, D. Sherrington: *Stealing the gold*
125. S. Atzeni, J. Meyer-ter-Vehn: *The physics of inertial fusion*
123. T. Fujimoto: *Plasma spectroscopy*
122. K. Fujikawa, H. Suzuki: *Path integrals and quantum anomalies*
121. T. Giamarchi: *Quantum physics in one dimension*
120. M. Warner, E. Terentjev: *Liquid crystal elastomers*
119. L. Jacak, P. Sitko, K. Wieczorek, A. Wojs: *Quantum Hall systems*
117. G. Volovik: *The Universe in a helium droplet*
116. L. Pitaevskii, S. Stringari: *Bose-Einstein condensation*
115. G. Dissertori, I. G. Knowles, M. Schmelling: *Quantum chromodynamics*
114. B. DeWitt: *The global approach to quantum field theory*
113. J. Zinn-Justin: *Quantum field theory and critical phenomena, Fourth edition*
112. R. M. Mazo: *Brownian motion—fluctuations, dynamics, and applications*
111. H. Nishimori: *Statistical physics of spin glasses and information processing—an introduction*
110. N. B. Kopnin: *Theory of nonequilibrium superconductivity*
109. A. Aharoni: *Introduction to the theory of ferromagnetism, Second edition*
108. R. Dobbs: *Helium three*
107. R. Wigmans: *Calorimetry*
106. J. Kübler: *Theory of itinerant electron magnetism*
105. Y. Kuramoto, Y. Kitaoka: *Dynamics of heavy electrons*
104. D. Bardin, G. Passarino: *The standard model in the making*
103. G. C. Branco, L. Lavoura, J. P. Silva: *CP Violation*
102. T. C. Choy: *Effective medium theory*
101. H. Araki: *Mathematical theory of quantum fields*
100. L. M. Pismen: *Vortices in nonlinear fields*
99. L. Mestel: *Stellar magnetism*
98. K. H. Bennemann: *Nonlinear optics in metals*
94. S. Chikazumi: *Physics of ferromagnetism*
91. R. A. Bertlmann: *Anomalies in quantum field theory*
90. P. K. Gosh: *Ion traps*
87. P. S. Joshi: *Global aspects in gravitation and cosmology*
86. E. R. Pike, S. Sarkar: *The quantum theory of radiation*
83. P. G. de Gennes, J. Prost: *The physics of liquid crystals*
73. M. Doi, S. F. Edwards: *The theory of polymer dynamics*
69. S. Chandrasekhar: *The mathematical theory of black holes*
51. C. Möller: *The theory of relativity*
46. H. E. Stanley: *Introduction to phase transitions and critical phenomena*
32. A. Abragam: *Principles of nuclear magnetism*
27. P. A. M. Dirac: *Principles of quantum mechanics*
23. R. E. Peierls: *Quantum theory of solids*

Preface

Since the discovery of the light emitting properties of the phenyl-based organic semiconductors in 1990 there has been a huge growth of interest in conjugated polymers. The potential device applications are enormous, ranging from optical switching to solar cells and light emitting devices. These new developments are a direct consequence of the active research in the 1980s on conjugated nonelectroluminescent polymers, such as *trans*-polyacetylene. Polyacetylene has particularly interesting and unusual low-lying electronic excitations, and has attracted much experimental and theoretical interest, culminating in the award of the Nobel prize for chemistry in 2000 for research in this field.

The progress in our understanding of the fundamental physics of conjugated polymers, which provides a crucial underpinning to the technological applications, has also been large. This progress has been driven by experimental, theoretical, and computational developments. A number of very careful and elegant linear and nonlinear optical spectroscopies over the last two decades have established the energies and symmetries of the excited states. Meanwhile, computational advances have been driven by the development of sophisticated numerical techniques, coupled with cheaper and more powerful computers. One of these numerical techniques is the density matrix renormalization group (DMRG) method. This method is highly suited for solving correlated one-dimensional problems.

Conjugated polymers behave as quasi-one-dimensional systems owing to their strong intramolecular interactions and rather weak intermolecular interactions. As a consequence, electron-electron interactions are weakly screened, and thus both electron-electron interactions and electron-nuclear coupling are fundamentally important in determining the electronic behaviour. Electronic interactions play a crucial role in determining the nature of electronic excitations as they completely change the noninteracting electronic description. Moreover, the coupling of these correlated electronic states to the nuclei is also a delicate and complicated problem. Together, electronic interactions and electron-nuclear coupling determine the relative energetic ordering of the electronic states, and this, in turn, largely determines the optical properties of conjugated polymers. This understanding of the origin and nature of the electronic states helps us to explain why some conjugated polymers, for example poly(*para*-phenylene), are electroluminescent, while others, for example *trans*-polyacetylene, are not.

One of the key aims of this book is to explain how electron-electron interactions and electron-nuclear coupling determine the types and character of the low-lying electronic states. Since these effects are complicated, our strategy will be to start with the simplest approximation of noninteracting electrons and gradually develop the full description. At each step care will be taken to explain how electron-electron interactions and electron-nuclear coupling modify the predictions of the simpler approximations.

We will see that one of the reasons why understanding the electronic properties of conjugated polymers is such a challenge is because the electronic potential energy is comparable to the electronic kinetic energy. In other words, the relevant parameter regime is intermediate between the weak and strong electron-electron interaction limits. A useful strategy is therefore to tackle these systems from both the weak and strong coupling extremes. In fact, light emitting polymers lie on the weak-coupling side of the intermediate regime, whereas nonelectroluminescent polymers (such as *trans*-polyacetylene) lie on the strong-coupling side.

We focus on semiempirical models of π -conjugated systems. There are two advantages to this strategy over studying *ab initio* models. First, reduced basis models in one dimension can be solved essentially exactly via the DMRG method for very large systems. Thus, there is no need to make approximations in the method which might obscure or prejudice an understanding of the physics. Second, being approximate, reduced basis models retain some symmetries not present in the *ab initio* models. In particular, electron-hole symmetry is particularly useful in characterizing neutral excited states. We also remark that although they are semiempirical, π -electron models are carefully parametrized so that they also provide accurate predictions of excited state energies.

Solving very large systems by the DMRG method reveals the physics of conjugated polymers not present in conjugated molecules, namely that when the size of the chain (or more precisely, the conjugation length) exceeds the spatial extent of the internal structure of the excited states a quasi-particle description becomes appropriate. Conjugated polymers exhibit a wealth of different quasi-particles: solitons, excitons, magnons, polarons, etc. It is an aim of this book to explain the origin and physical consequences of these quasi-particles. In particular, the description of an exciton as two independent quasi-particles will prove to be a very useful concept.

Having established how two key effects in conjugated polymers (electronic interactions and electron-nuclear coupling) affect their electronic properties, the third key effect will be discussed, namely, disorder. Most conjugated polymer systems are subject to conformational and environmental disorder. Disorder localizes charges and excitons, and determines their energetic and spatial distributions. An understanding of the role of disorder is a necessary prerequisite for the development of theories of charge and energy transport.

Nonlinear optical measurements provide the most direct probe of the electronic states. Conversely, the nonlinear susceptibilities can be calculated if there exists a theoretical understanding of the excited states. We describe the theory of linear and nonlinear optical processes, and recast the so-called essential states model in terms of the primary excitons.

We demonstrate how our theoretical understanding of excited states enables us to make a consistent interpretation of experimental results. Two chapters draw these themes together in discussing *trans*-polyacetylene, and the technologically important phenyl-based light emitting polymers.

Once an understanding of these intramolecular processes is established, a final aim of this book will be to explain electronic processes arising from intermolecular interactions. Thus, energy transfer and migration, and excited state complexes involving two

or more polymer chains are described. Another important consequence of interchain interactions is dispersion interactions, which significantly modify the energy of some intrachain excitations. An understanding of this effect is crucial to the interpretation of optical experiments.

Throughout we apply models to, and develop theories for, conjugated polymers with relatively simple chemical structures, e.g., *trans*-polyacetylene and poly(*para*-phenylene). These polymers serve as model systems to understand the generic electronic properties of most classes of conjugated polymers.

The book is therefore organized as follows. Chapter 1 gives a brief overview of the electronic properties of conjugated polymers. Our basic models for describing these properties are semiempirical π -electron models. So, Chapter 2 introduces and motivates these models. Next, we consider the solution of these models in various limits: noninteracting electrons with fixed geometry in Chapter 3, noninteracting electrons with electron-nuclear coupling in Chapter 4, and interacting electrons with fixed geometry in Chapter 5. Chapter 6 is devoted to a discussion of excitons, as these are so important in determining the photophysical properties of conjugated polymers. The electronic states of interacting electrons with electron-nuclear coupling are described Chapter 7. The next two chapters illustrate the applicability of the concepts thus developed via a discussion of the experimental and theoretical investigations of *trans*-polyacetylene and light emitting polymers in Chapters 8 and 9, respectively. The roles of disorder and electron-nuclear coupling in localizing excitons are then described in Chapter 10, where the fundamental concept of a chromophore is given a quantitative definition. Chapter 11 introduces the nonlinear optical spectroscopies that are used to identify the excited states of conjugated polymers discussed in Chapters 8 and 9. Finally, Chapter 12 describes excitonic processes in conjugated polymers.

This book was written with two kinds of readers in mind, the first being experimentalists who wish to understand and interpret their experimental data in terms of the fundamental electronic and optical properties of conjugated polymers. The other type of readers are theoretical and computational chemists and physicists who want both to understand the fundamental properties of conjugated polymers and to develop models and perform calculations of their own. For these readers there are a number of appendices containing material too technical for the main chapters.

Acknowledgements

My interest in conjugated polymers began in 1995 when Donal Bradley established a polymer optoelectronics research group in the Department of Physics and Astronomy at the University of Sheffield. At the same time three theorists then at Sheffield, Robert Bursill, Gillian Gehring, and Tao Xiang, were using Steve White's highly-accurate density matrix renormalization group (DMRG) method to solve one-dimensional correlated electron problems. The opportunity to apply the DMRG method to models of conjugated polymers at the same time as developing a collaboration with the experimental group was too great to miss.

Since then I have learnt a great deal about conjugated polymers from a large number of people. Donal Bradley, Paul Lane (both then in Sheffield) and Simon Martin are three experimentalists who patiently answered many questions on the optoelectronic properties of conjugated polymers.

Collaborations and/or conversations with David Beljonne, Peter Bobbert, Richard Friend, Neil Greenham, Anna Köhler, Mikhail Yu Lavrentiev, David Lidzey, Sumit Mazumdar, Carlos Silva, and David Yaron are all greatly appreciated.

Last, but not least, my sincere thanks to my coworker since 1995, Robert Bursill. It was he who principally developed the DMRG programs that we have used to study correlated models of conjugated polymers. This book was inspired by those studies.

The first edition of this book was started in May 2002. Without the benefit of a research fellowship from the Leverhulme Trust in 2003–04 it is unlikely that it would ever have been completed. I am very grateful to the Leverhulme Trust for the opportunity to be relieved from my teaching duties for one year. I would also like to thank the Engineering and Physical Sciences Research Council, the Gordon Godfrey Bequest of the University of New South Wales, and the Royal Society for sponsoring my research in this subject. Thanks too to the University of New South Wales, and the Cavendish Laboratory and Clare Hall, Cambridge for their hospitality in 2003–04.

I would like to thank Sonke Adlung at Oxford University Press for his enthusiasm for the book proposal, and his colleagues at OUP and Mark Fox in Sheffield for their help over the preparation of the manuscript. Finally, I am very grateful to Simon Martin and Sumit Mazumdar for providing valuable critical feedback on the draft manuscript. As always, however, all errors and omissions are the sole responsibility of the author.

The second edition of this book would not have been possible without the output of undergraduate and graduate students that I have supervised over the last six years in Oxford. They are: Igor Boczarow, Christopher Menelaou, Nattapong Paiboonvorachat, Christopher Pearson, David Trembath, Oliver Tozer, Alec Ward, and Thomas Wharram, to whom I am very grateful. I am also grateful to a former post-doc, Dmitry Makhov, for his contributions to this new edition. Finally, I thank John Freeman for his help with some of the figures.

This book is dedicated to my mother and father.

Contents

1	Introduction to conjugated polymers	1
2	π-electron theories of conjugated polymers	7
2.1	Introduction	7
2.2	The many body Hamiltonian	7
2.3	The Born-Oppenheimer approximation	8
2.4	Second quantization of the Born-Oppenheimer Hamiltonian	10
2.5	sp^n hybridization	12
2.6	π -electron models	14
2.7	Electron-nuclear coupling	17
2.8	Summary of π -electron models	19
2.9	Symmetries and quantum numbers	22
3	Noninteracting electrons	26
3.1	Introduction	26
3.2	The noninteracting (Hückel) Hamiltonian	26
3.3	The ethylene dimer	27
3.4	Undimerized chains	30
3.5	Dimerized chains	33
3.6	The ground state and electron-hole excitations	36
3.7	Symmetries	38
3.8	Bond order	41
4	Electron-nuclear coupling I: Noninteracting electrons	44
4.1	Introduction	44
4.2	The Peierls model	45
4.3	The dimerized ground state	46
4.4	Self-consistent equations for $\{\Delta_n\}$	48
4.5	Solitons	50
4.6	Soliton-antisoliton pair production	54
4.7	Nondegenerate systems	57
4.8	The continuum limit of the Su-Schrieffer-Heeger model	59
4.9	Polarons	61
4.10	Dynamics of the Su-Schrieffer-Heeger model	62
4.11	Self-trapping	63
4.12	Concluding remarks	63
5	Interacting electrons	65
5.1	Introduction	65
5.2	The weak-coupling limit	70

5.3	The strong-coupling limit	71
5.4	The phase diagram of the undoped Pariser-Parr-Pople model	74
5.5	The valence bond method	75
6	Excitons in conjugated polymers	78
6.1	Introduction	78
6.2	The weak-coupling limit	79
6.3	The strong-coupling limit	93
6.4	The intermediate-coupling regime	96
6.5	Concluding remarks	98
7	Electron-nuclear coupling II: Interacting electrons	101
7.1	Introduction	101
7.2	The Pariser-Parr-Pople-Peierls model	102
7.3	Dimerization and optical gaps	103
7.4	Excited states and soliton structures	107
7.5	Polarons	113
7.6	Extrinsic dimerization	113
7.7	Quantum phase transition	114
7.8	Concluding remarks	118
8	Linear polyenes and <i>trans</i>-polyacetylene	120
8.1	Introduction	120
8.2	Predictions from the Pariser-Parr-Pople-Peierls model	123
8.3	Role of nuclear zero-point fluctuations	128
8.4	Character of the excited states of <i>trans</i> -polyacetylene	129
8.5	Other theoretical approaches	131
9	Light emitting polymers	132
9.1	Introduction	132
9.2	Poly(<i>para</i> -phenylene)	137
9.3	Poly(<i>para</i> -phenylene vinylene)	150
9.4	Other theoretical approaches	152
9.5	Exciton binding energies	158
9.6	The excited states of light emitting polymers	159
9.7	Electronic coupling to nuclear degrees of freedom	161
10	Exciton localization in disordered polymers	168
10.1	Introduction	168
10.2	Definition of the exciton conjugation length	169
10.3	Origins of disorder	170
10.4	Localization of vertical excitations	171
10.5	Dynamical localization	184
10.6	Concluding remarks	189
11	Optical processes in conjugated polymers	192
11.1	Introduction	192
11.2	Linear optical processes	193

11.3	Evaluation of the transition dipole moments	194
11.4	Nonlinear optical processes	204
11.5	Size-dependencies of $\chi^{(n)}$	210
11.6	Photophysical processes in conjugated polymers	211
12	Excitonic processes in conjugated polymers	212
12.1	Introduction	212
12.2	Exciton transfer	212
12.3	Exciton diffusion in the condensed phase	226
12.4	Excited molecular complexes	231
12.5	Second order dispersion interactions	232
13	Epilogue	240
Appendix A	Dirac bra-ket operator representation of one-particle Hamiltonians	242
A.1	The Hückel Hamiltonian	242
A.2	The Frenkel exciton Hamiltonian	243
Appendix B	Electron-hole symmetry and average occupation number	245
Appendix C	Single-particle eigensolutions of a periodic polymer chain	247
C.1	Dimerized chain	248
C.2	Poly(<i>para</i> -phenylene)	249
Appendix D	The Holstein model	250
D.1	The model	251
D.2	General solutions	253
D.3	Variational calculation	254
D.4	Optical intensities	254
Appendix E	Derivation of the effective-particle Schrödinger equation	257
Appendix F	Hydrogenic solutions of the effective-particle exciton models	262
F.1	The weak-coupling limit	262
F.2	The strong-coupling limit	265
Appendix G	Valence-bond description of benzene	267
Appendix H	Derivation of the Frenkel exciton Hamiltonian	270
Appendix I	Evaluation of the electronic transition dipole moments	277
I.1	The weak-coupling limit	277
I.2	The strong-coupling limit	280

Appendix J Spin-orbit coupling in π-conjugated polymers	281
J.1 Spin-orbit coupling of π -electrons	281
J.2 Symmetry restrictions on spin-orbit coupling	282
J.3 Spin-orbit coupling and spatial wavefunctions	283
Appendix K Derivation of the line dipole approximation	284
Appendix L Direct configuration interaction-singles calculations for the Pariser-Parr-Pople model	287
L.1 Hartree-Fock Solutions	287
L.2 Direct CI-singles method	288
Appendix M Density matrix renormalization group method	289
M.1 Introduction to the real-space method	289
M.2 Local Hilbert space truncation	293
References	294
Index	302

1

Introduction to conjugated polymers

Research into the electronic, optical, and magnetic properties of conjugated polymers began in the 1970s after a number of seminal experimental achievements. First, the synthesis of polyacetylene thin films (Itô *et al.* 1974) and the subsequent success in doping these polymers to create conducting polymers (Chiang *et al.* 1977) established the field of synthetic metals. Second, the synthesis of the phenyl-based polymers and the discovery of electroluminescence under low voltages in these systems (Burroughes *et al.* 1990) established the field of polymer optoelectronics.

The electronic and optical properties of conjugated polymers, coupled with their mechanical properties and intrinsic processing advantages, means that they are particularly attractive materials for the electronics industry. There are many potential applications including, light emitting devices, nonlinear optical devices, photovoltaic devices, plastic field-effect transistors, and electromagnetic shielding. The discovery and development of conductive polymers was recognized by the award of the Nobel prize for chemistry in 2000 to Heeger (2000), MacDiarmid (2000), and Shirakawa (2000).

A conjugated polymer is a carbon-based macromolecule through which the valence π -electrons are delocalized.¹ *Trans*-polyacetylene, illustrated in Fig. 1.1, is a linear polyene, whose ground state structure is composed of alternating long and short bonds. Also shown in Fig. 1.1 are two other linear polyenes, *cis*-polyacetylene and polydiacetylene. The light emitting polymers, for example, poly(*para*-phenylene) (or PPP) and poly(*para*-phenylene vinylene) (or PPV), are characterized by containing a phenyl ring in their repeat units. PPP and PPV are illustrated in Fig. 1.2.

As well as their many important technological applications, conjugated polymers are also active components in many biological photophysical processes, for example, as light collectors in photosynthesis, and in the vision mechanism via photoisomerization. Charge transport in organic molecules is also an important component of cellular function. Many of the concepts developed in this book are therefore applicable to these biological systems.

Conjugated polymers exhibit electronic properties that are quite different from those observed in the corresponding inorganic metals or semiconductors. These unusual electronic properties may essentially be attributed to fact that conjugated polymers behave as quasi-one-dimensional systems owing to their strong intramolecular electronic interactions and relatively weak intermolecular electronic interactions. Weak intermolecular electronic interactions (arising from poor electronic wavefunction over-


¹Conjugate from the Latin *conjugatus*, meaning to join or unite.

2 Introduction to conjugated polymers

trans-polyacetylene



cis-polyacetylene



polydiacetylene



Fig. 1.1 The carbon backbone of some linear polyenes. The hydrogen atoms are not shown. More detailed chemical structures are illustrated in Chapter 2.

lap) coupled to strong dissipation (or dephasing) mechanisms means that quantum mechanical coherence is generally confined to a single chain, or at most a few other chains. Electronic wavefunctions are therefore typically localized on single chains, or to pairs of chains in the case of excited state complexes. This quasi-one-dimensionality also means that electron-electron interactions are weakly screened. Thus, electronic correlations are important in determining the character of the electronic states. Another important factor in determining the character of the electronic states is that the electrons and nuclei are strongly coupled. As for electronic interactions, the effects of electron-nuclear coupling are enhanced in low dimensions.

Much early theoretical work on conjugated oligomers and polymers treated electron-electron and electron-nuclear interactions independently. In the 1950s the focus was on the role of electron-nuclear interactions in causing a metal-semiconductor transition in one-dimensional metals (Fröhlich 1954; Peierls 1955), and in determining the bond alternation in linear polyenes (Ooshika 1957, 1959; Longuet-Higgins and Salem 1959). It was also realized that a broken-symmetry ground state of bond alternation implies bond defects between different domains of bond alternation (Ooshika 1957; Longuet-Higgins and Salem 1959), and to associated mid-gap electronic states (Pople and Walmsley 1962). Theoretical and experimental investigations into excited states and their associated bond defects (or solitons) grew rapidly after the introduction of a simplified model of electron-nuclear interactions in *trans*-polyacetylene by Su, Schrieffer, and Heeger (Su *et al.* 1979). These developments are reviewed in (Heeger *et al.* 1988).

An alternative point of view, namely that electron-electron interactions are important in determining the electronic properties in conjugated polymers, was advocated by Ovchinnikov and coworkers (Ovchinnikov *et al.* 1973), who argued that electronic