

John Milnor

Dynamics in One Complex Variable

Third Edition

单复变动力系统 第3版



PRINCETON UNIVERSITY PRESS

世界图书出版公司
www.wpcbj.com.cn

For sale only in the People's Republic of China

新书预告

书号	书名	作者
51005065	Theory of Lie Groups	C. Chevalley
51005283	Theory of Probability 3rd ed.	H. Jeffreys
51005285	Using Algebraic Geometry 2nd ed.	D. A. Cox
51005813	A Primer of Real Analytic Functions 2nd ed.	S. G. Krantz
51005827	A Probability Path	S. I. Resnick
51005812	Astrophysics in a Nutshell	D. Maoz
51005841	Classical Potential Theory and Its Probabilistic Counterpart	J. L. Doob
51005828	Classical Topics in Complex Function Theory	R. Remmert
51005838	Condensed Matter in a Nutshell	G. D. Mahan
51005808	Credit Risk: Modeling, Valuation and Hedging	T. R. Bielecki
51005830	Dynamics in One Complex Variable 3rd ed.	J. Milnor
51005839	Efficient Methods for Valuing Interest Rate Derivatives	A. Pelsser
51005823	Elementary Particle Physics in a Nutshell	C. G. Tully
51005836	Elementary Topics in Differential Geometry	J. A. Thorpe
51005837	Galerkin Finite Element Methods for Parabolic Problems 2nd ed.	V. Thomée
51005826	Hamiltonian Methods in the Theory of Solitons	L. D. Faddeev
51005840	Ideals, Varieties, and Algorithms	D. Cox
51005807	Implicit Functions and Solution Mappings	A. L. Dontchev
51005809	Interacting Particle Systems	T. M. Liggett
51005844	Introduction to Applied Nonlinear Dynamical Systems and Chaos	S. Wiggins
51005820	Large Eddy Simulation for Compressible Flows	E. Garnier
51005810	Linear Algebraic Groups 2nd ed.	T. A. Springer
51005832	Malliavin Calculus for Lévy Processes with Applications to Finance	G. Di Nunno
51005843	Mathematical Methods for Financial Markets	M. Jeanblanc
51005831	Multiple Integrals in the Calculus of Variations	C. B. Morrey Jr.
51005824	Nuclear Physics in a Nutshell	C. A. Bertulani
51005811	String Theory in a Nutshell	E. Kiritsis
51005822	Structural Macroeconometrics 2nd ed.	D. N. DeJong
51005842	Variational Methods in Imaging	O. Scherzer

ISBN 978-7-5100-5830-1

定价: 79.00元

ISBN 978-7-5100-5830-1



9 787510 058301 >

上架建议: 数学/影印版

PRINCETON

数学经典教材
(影印版)

Dynamics in One Complex Variable

单复变动力系统

第3版


北京大学出版社

Dynamics in One Complex Variable

THIRD EDITION

by

John Milnor



PRINCETON UNIVERSITY PRESS

PRINCETON AND OXFORD

2006

Copyright © 2006 by Princeton University Press

Second edition © 2000, published by Vieweg Verlag, Wiesbaden, Germany
First edition © 1999, published by Vieweg Verlag, Wiesbaden, Germany

Published by Princeton University Press, 41 William Street,
Princeton, New Jersey 08540

In the United Kingdom: Princeton University Press,
3 Market Place, Woodstock, Oxfordshire OX20 1SY

All Rights Reserved

The Annals of Mathematics Studies are edited by
Phillip A. Griffiths, John N. Mather, and Elias M. Stein

Library of Congress Cataloging-in-Publication Data

Milnor, John Willard, 1931–

Dynamics in one complex variable / John Milnor— 3rd ed.

p. cm. — (Annals of mathematics studies ; no. 160)

Includes bibliographical references and index

ISBN-13: 978-0-691-12487-2 (acid-free paper)

ISBN-10: 0-691-12487-6 (acid-free paper)

ISBN-13: 978-0-691-12488-9 (pbk. : acid-free paper)

ISBN-10: 0-691-12488-4 (pbk. : acid-free paper)

1. Functions of complex variables. 2. Holomorphic mappings.
3. Riemann surfaces. I. Title. II. Series.

QA331.7.M55 2006

515'.93—dc22

2005051060

British Library Cataloging-in-Publication Data is available.

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without permission in writing from the Publisher.

图书在版编目 (CIP) 数据

单复变动力系统: 第3版 = Dynamics in one complex variable 3rd ed. : 英文/(美) 米尔诺 (Milnor, J.) 著. —影印本. —北京: 世界图书出版公司北京公司, 2013. 3
ISBN 978 - 7 - 5100 - 5830 - 1

I. ①单… II. ①米… III. ①变量—动力系统 (数学)—英文 IV. ①019

中国版本图书馆 CIP 数据核字 (2013) 第 035505 号

书 名: Dynamics in One Complex Variable 3rd ed.

作 者: John Milnor

中译名: 单复变动力系统 第3版

责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河市国英印务有限公司

发 行 者: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010 - 64021602, 010 - 64015659

电子信箱: kjb@wpcbj.com.cn

开 本: 16 开

印 张: 20

版 次: 2013 年 3 月

版权登记: 图字: 01 - 2013 - 0906

书 号: 978 - 7 - 5100 - 5830 - 1

定 价: 79.00 元

Annals of Mathematics Studies

Number 160

PREFACE TO THE THIRD EDITION

This book studies the dynamics of iterated holomorphic mappings from a Riemann surface to itself, concentrating on the classical case of rational maps of the Riemann sphere. It is based on introductory lectures given at Stony Brook during the fall term of 1989–90 and in later years. I am grateful to the audiences for a great deal of constructive criticism and to Bodil Branner, Adrien Douady, John Hubbard, and Mitsuhiro Shishikura, who taught me most of what I know in this field. Also, I want to thank a number of individuals for their extremely helpful criticisms and suggestions, especially Adam Epstein, Rodrigo Perez, Alfredo Poirier, Lasse Rempe, and Saeed Zakeri. Araceli Bonifant has been particularly helpful in the preparation of this third edition.

There have been a number of extremely useful surveys of holomorphic dynamics over the years. See the textbooks by Devaney [1989], Beardon [1991], Carleson and Gamelin [1993], Steinmetz [1993], and Berteloot and Mayer [2001], as well as expository articles by Brolin [1965], Douady [1982–83, 1986, 1987], Blanchard [1984], Lyubich [1986], Branner [1989], Keen [1989], Blanchard and Chiu [1991], and Eremenko and Lyubich [1990]. (See the list of references at the end of the book.)

This subject is large and rapidly growing. These lectures are intended to introduce the reader to some key ideas in the field, and to form a basis for further study. The reader is assumed to be familiar with the rudiments of complex variable theory and of 2-dimensional differential geometry, as well as some basic topics from topology. The necessary material can be found for example in Ahlfors [1966], Hocking and Young [1961], Munkres [1975], Thurston [1997], and Willmore [1959]. However, two big theorems will be used here without proof, namely the Uniformization Theorem in §1 and the existence of solutions for the measurable Beltrami equation in Appendix F. (See the references in those sections.)

The basic outline of this third edition has not changed from previous editions, but there have been many improvements and additions. A brief historical survey has been added in §4.1, the definition of Lattès map has been made more inclusive in §7.4, the Écalle-Voronin theory of parabolic points is described in §10.12, the résidu itératif is studied in §12.9, the material on two complex variables in Appendix D has been expanded, and recent results on effective computability have been added in Appendix H. The list of references has also been updated and expanded.

John Milnor
Stony Brook, August 2005

CHRONOLOGICAL TABLE

Following is a list of some of the founders of the field of complex dynamics.

Ernst Schröder	1841–1902
Hermann Amandus Schwarz	1843–1921
Henri Poincaré	1854–1912
Gabriel Kœnigs	1858–1931
Léopold Leau	1868–1940(?)
Lucjan Emil Böttcher	1872– ?
Samuel Lattès	1873–1918
Constantin Carathéodory	1873–1950
Paul Montel	1876–1975
Pierre Fatou	1878–1929
Paul Koebe	1882–1945
Arnaud Denjoy	1884–1974
Gaston Julia	1893–1978
Carl Ludwig Siegel	1896–1981
Hubert Cremer	1897–1983
Herbert Grötzsch	1902–1993
Charles B. Morrey	1907–1984
Lars Ahlfors	1907–1996
Lipman Bers	1914–1993
Irvine Noel Baker	1932–2001
Michael (Michel) R. Herman	1942–2000

Among the many present-day workers in the field, let me mention a few whose work is emphasized in these notes: Adrien Douady (b. 1935), Dennis P. Sullivan (b. 1941), Bodil Branner (b. 1943), John Hamal Hubbard (b. 1945), William P. Thurston (b. 1946), Mary Rees (b. 1953), Jean-Christophe Yoccoz (b. 1955), Curtis McMullen (b. 1958), Mikhail Y. Lyubich (b. 1959), and Mitsuhiro Shishikura (b. 1960).

TABLE OF CONTENTS

List of Figures	vi
Preface to the Third Edition	vii
Chronological Table	viii
Riemann Surfaces	
1. Simply Connected Surfaces	1
2. Universal Coverings and the Poincaré Metric	13
3. Normal Families: Montel's Theorem	30
Iterated Holomorphic Maps	
4. Fatou and Julia: Dynamics on the Riemann Sphere	39
5. Dynamics on Hyperbolic Surfaces	56
6. Dynamics on Euclidean Surfaces	65
7. Smooth Julia Sets	69
Local Fixed Point Theory	
8. Geometrically Attracting or Repelling Fixed Points	76
9. Böttcher's Theorem and Polynomial Dynamics	90
10. Parabolic Fixed Points: The Leau-Fatou Flower	104
11. Cremer Points and Siegel Disks	125
Periodic Points: Global Theory	
12. The Holomorphic Fixed Point Formula	142
13. Most Periodic Orbits Repel	153
14. Repelling Cycles Are Dense in J	156
Structure of the Fatou Set	
15. Herman Rings	161
16. The Sullivan Classification of Fatou Components	167
Using the Fatou Set to Study the Julia Set	
17. Prime Ends and Local Connectivity	174
18. Polynomial Dynamics: External Rays	188
19. Hyperbolic and Subhyperbolic Maps	205
Appendix A. Theorems from Classical Analysis	219
Appendix B. Length-Area-Modulus Inequalities	226
Appendix C. Rotations, Continued Fractions, and Rational Approximation	234
Appendix D. Two or More Complex Variables	246
Appendix E. Branched Coverings and Orbifolds	254
Appendix F. No Wandering Fatou Components	259
Appendix G. Parameter Spaces	266
Appendix H. Computer Graphics and Effective Computation	271
References	277
Index	293

LIST OF FIGURES

- | | |
|---|--|
| 1: Coordinate neighborhoods, 1 | 25: A rational parabolic Julia set, 124 |
| 2: Part of $\mathbb{D} \setminus \{0\}$, isometrically embedded into $\mathbb{R}^3$, 20 | 26: Siegel disks, 127 |
| 3: Cross-ratio and hyperbolic distance, 25 | 27: Classifying irrational numbers, 128 |
| 4: Poincaré neighborhoods in a subset, 34 | 28: Siegel disk with fjords, 132 |
| 5: Five polynomial Julia sets, 42 | 29: Double Mandelbrot set, 145 |
| 6: Four rational Julia sets, 43 | 30: Parabolic cubic parameter space, 151 |
| 7: A family of “rabbits”, 50 | 31: Homoclinic sketch, 156 |
| 8: A cubic Julia set, 52 | 32: A Herman ring, 164 |
| 9: Region with bad boundary, 63 | 33: Snail sketch, 169 |
| 10: Sine Julia set, 66 | 34: A polynomial Julia set, 172 |
| 11: Quasicircle Julia set, 80 | 35: Wandering domains, 172 |
| 12: Spirals in Julia sets, 87 | 36: A Baker domain, 172 |
| 13: Böttcher domains, 94 | 37: Some bad boundaries, 175 |
| 14: Böttcher domain with an extra critical point, 94 | 38: The witch’s broom, 187 |
| 15: Equipotentials for a dendrite, 97 | 39: Rabbit equipotentials and rays, 189 |
| 16: Cantor set equipotentials, 97 | 40: A symmetric comb, 192 |
| 17: Disconnected Julia set, 99 | 41: Two cubic polynomial Julia sets, 196 |
| 18: Parabolic sketch, 105 | 42: Zoom to a repelling point, 199 |
| 19: Parabolic point with 3 basins, 106 | 43: The annulus $f^{-1}(\mathbb{D}_r) \setminus \overline{\mathbb{D}}_r$, 208 |
| 20: Parabolic sector, 107 | 44: Orbifold sketch, 211 |
| 21: Parabolic point with a 7th root of unity as multiplier, 109 | 45: Successive area estimates for some filled Julia sets, 224 |
| 22: Parabolic flower, 112 | 46: An irrational rotation, 234 |
| 23: Écalé–Voronin classification, 117 | 47: Successive close returns, 237 |
| 24: The cauliflower Julia set, 121 | 48: The Gauss map of $(0, 1]$, 245 |
| | 49: The Mandelbrot set, 267 |

RIEMANN SURFACES

§1. Simply Connected Surfaces

The first three sections will present an overview of some background material.

If $V \subset \mathbb{C}$ is an open set of complex numbers, a function $f : V \rightarrow \mathbb{C}$ is called *holomorphic* (or “complex analytic”) if the first derivative

$$z \mapsto f'(z) = \lim_{h \rightarrow 0} (f(z+h) - f(z))/h$$

is defined and continuous as a function from V to $\mathbb{C}$, or equivalently if f has a power series expansion about any point $z_0 \in V$ which converges to f in some neighborhood of z_0 . (See, for example, Ahlfors [1966].) Such a function is *conformal* if the derivative $f'(z)$ never vanishes. Thus our conformal maps must always preserve orientation. It is *univalent* (or *schlicht*) if it is conformal and one-to-one.

By a *Riemann surface* S we mean a connected complex analytic manifold of complex dimension 1. Thus S is a connected Hausdorff space. Furthermore, in some neighborhood U of an arbitrary point of S we can choose a *local uniformizing parameter* (or “coordinate chart”) which maps U homeomorphically onto an open subset of the complex plane $\mathbb{C}$, with the following property: In the overlap $U \cap U'$ between two such neighborhoods, each of these local uniformizing parameters can be expressed as a holomorphic function of the other.




Figure 1. Overlapping coordinate neighborhoods.

By definition, two Riemann surfaces S and S' are *conformally isomorphic* (or *biholomorphic*) if and only if there is a homeomorphism from S onto S' which is holomorphic in terms of the respective local uniformizing parameters. (It is an easy exercise to show that the inverse map $S' \rightarrow S$ must then also be holomorphic.) In the special case $S = S'$, such a conformal isomorphism $S \rightarrow S$ is called a *conformal automorphism* of S .

Although there are uncountably many conformally distinct Riemann surfaces, there are only three distinct surfaces in the simply connected case. (By definition, the surface S is *simply connected* if every map from a circle into S can be continuously deformed to a constant map. Compare §2.) The following result is due to Poincaré and to Koebe.

Theorem 1.1 (Uniformization Theorem). *Any simply connected Riemann surface is conformally isomorphic either*

- (a) *to the plane $\mathbb{C}$ consisting of all complex numbers $z = x + iy$,*
- (b) *to the open disk $\mathbb{D} \subset \mathbb{C}$ consisting of all z with $|z|^2 = x^2 + y^2 < 1$, or*
- (c) *to the Riemann sphere $\hat{\mathbb{C}}$ consisting of $\mathbb{C}$ together with a point at infinity, using $\zeta = 1/z$ as local uniformizing parameter in a neighborhood of the point at infinity.*

This is a generalization of the classical Riemann Mapping Theorem. We will refer to these three cases as the *Euclidean*, *hyperbolic*, and *spherical* cases, respectively. (Compare §2.) The proof of Theorem 1.1 is nontrivial and will not be given here. However, proofs may be found in Koebe [1907], Ahlfors [1973], Beardon [1984], Farkas and Kra [1980], Nevanlinna [1967], and in Springer [1957]. (See also Fisher, Hubbard, and Wittner [1988].) By assuming this result, we will be able to pass more quickly to interesting ideas in holomorphic dynamics.

The Open Disk $\mathbb{D}$. For the rest of this section, we will discuss these three surfaces in more detail. We begin with a study of the unit disk $\mathbb{D}$.

Lemma 1.2 (Schwarz Lemma). *If $f : \mathbb{D} \rightarrow \mathbb{D}$ is a holomorphic map with $f(0) = 0$, then the derivative at the origin satisfies $|f'(0)| \leq 1$. If equality holds, $|f'(0)| = 1$, then f is a rotation about the origin. That is, $f(z) = cz$ for some constant $c = f'(0)$ on the unit circle. On the other hand, if $|f'(0)| < 1$, then $|f(z)| < |z|$ for all $z \neq 0$.*

(The Schwarz Lemma was first proved, in this generality, by Carathéodory.)

Remarks. If $|f'(0)| = 1$, it follows that f is a conformal automorphism of the unit disk. But if $|f'(0)| < 1$ then f cannot be a conformal automorphism of $\mathbb{D}$, since the composition with any $g : (\mathbb{D}, 0) \rightarrow (\mathbb{D}, 0)$ would have derivative $g'(0)f'(0) \neq 1$. The example $f(z) = z^2$ shows that f may map $\mathbb{D}$ onto itself even when $|f(z)| < |z|$ for all $z \neq 0$ in $\mathbb{D}$.

Proof of Lemma 1.2. We use the *Maximum Modulus Principle*, which asserts that a nonconstant holomorphic function cannot attain its maximum absolute value at any interior point of its region of definition. First note that the quotient function $q(z) = f(z)/z$ is well defined and holomorphic throughout the disk $\mathbb{D}$, as one sees by dividing the local power series for f by z . Since $|q(z)| < 1/r$ when $|z| = r < 1$, it follows by the Maximum Modulus Principle that $|q(z)| < 1/r$ for all z in the disk $|z| \leq r$. Since this is true for all $r \rightarrow 1$, it follows that $|q(z)| \leq 1$ for all $z \in \mathbb{D}$. Again by the Maximum Modulus Principle, we see that the case $|q(z)| = 1$, for some z in the open disk, can occur only if the function $q(z)$ is constant. If we exclude this case $f(z)/z \equiv c$, then it follows that $|q(z)| = |f(z)/z| < 1$ for all $z \neq 0$, and similarly that $|q(0)| = |f'(0)| < 1$. $\square$

Here is a useful variant statement.

Lemma 1.2' (Cauchy Derivative Estimate). *If f maps the disk of radius r about z_0 into some disk of radius s , then*

$$|f'(z_0)| \leq s/r.$$

Proof. This follows easily from the Cauchy integral formula (see, for example, Ahlfors [1966]): Set $g(z) = f(z + z_0) - f(z_0)$, so that g maps the disk $\mathbb{D}_r$ centered at the origin to the disk $\mathbb{D}_s$ centered at the origin. Then

$$f'(z_0) = g'(0) = \frac{1}{2\pi i} \oint_{|z|=r_1} \frac{g(z) dz}{z^2}$$

for all $r_1 < r$, and the conclusion follows easily. $\square$

(An alternative proof, based on the Schwarz Lemma, is described in Problem 1-a at the end of this section. With an extra factor of 2 on the right, this inequality would follow immediately from Lemma 1.2 simply by linear changes of variable, since the target disk of radius s must be contained in the disk of radius $2s$ centered at the image $f(z_0)$.)

As an easy corollary, we obtain the following.

Theorem 1.3 (Liouville Theorem). *A bounded function f which is defined and holomorphic everywhere on $\mathbb{C}$ must be constant.*

For in this case we have s fixed but r arbitrarily large, hence f' must be identically zero. $\square$

As another corollary, we see that our three model surfaces really are distinct. There are natural inclusion maps $\mathbb{D} \rightarrow \mathbb{C} \rightarrow \hat{\mathbb{C}}$. Yet it follows from the Maximum Modulus Principle that every holomorphic map $\hat{\mathbb{C}} \rightarrow \mathbb{C}$ must be constant, and from Liouville's Theorem that every holomorphic map $\mathbb{C} \rightarrow \mathbb{D}$ must be constant.

Another closely related statement is the following. Let U be an open subset of $\mathbb{C}$.

Theorem 1.4 (Weierstrass Uniform Convergence Theorem). *If a sequence of holomorphic functions $f_n : U \rightarrow \mathbb{C}$ converges uniformly to the limit function f , then f itself is holomorphic. Furthermore, the sequence of derivatives f'_n converges, uniformly on any compact subset of U , to the derivative f' .*

It follows inductively that the sequence of second derivatives f''_n converges, uniformly on compact subsets, to f'' , and so on.

Proof of Theorem 1.4. Note first that the sequence of first derivatives f'_n , restricted to any compact subset $K \subset U$, converges uniformly. For example, if $|f_n(z) - f_m(z)| < \epsilon$ for $m, n > N$, and if the r -neighborhood of any point of K is contained in U , then it follows from Lemma 1.2' that $|f'_n(z) - f'_m(z)| < \epsilon/r$ for $m, n > N$ and for all $z \in K$. This proves uniform convergence of $\{f'_n\}$ restricted to K to some limit function g , which is necessarily continuous since any uniform limit of continuous functions is continuous. It follows that the integral of f'_n along any path in U converges to the integral of g along this path. Thus $f = \lim f_n$ is an indefinite integral of g , and hence g can be identified with the derivative of f . Thus f has a continuous complex first derivative and therefore is a holomorphic function. $\square$

Conformal Automorphism Groups. For any Riemann surface S , the notation $\mathcal{G}(S)$ will be used for the group consisting of all conformal automorphisms of S . The identity map will be denoted by $I = I_S \in \mathcal{G}(S)$.

We first consider the case of the Riemann sphere $\hat{\mathbb{C}}$ and show that $\mathcal{G}(\hat{\mathbb{C}})$ can be identified with a well-known complex Lie group. Thus $\mathcal{G}(\hat{\mathbb{C}})$ is not only a group, but also a complex manifold, and the product and inverse operations for this group are both holomorphic maps.

Lemma 1.5 (Möbius Transformations). *The group $\mathcal{G}(\hat{\mathbb{C}})$ of all conformal automorphisms of the Riemann sphere is equal to the group of all fractional linear transformations (also called Möbius transformations)*

$$g(z) = (az + b)/(cz + d),$$

where the coefficients are complex numbers with $ad - bc \neq 0$.

Here, if we multiply numerator and denominator by a common factor, then it is always possible to normalize so that the determinant $ad - bc$ is equal to $+1$. The resulting coefficients are well defined up to a simultaneous change of sign. Thus it follows that the group $\mathcal{G}(\hat{\mathbb{C}})$ of conformal automorphisms can be identified with the complex 3-dimensional Lie group $\mathrm{PSL}(2, \mathbb{C})$, consisting of all 2×2 complex matrices with determinant $+1$ modulo the subgroup $\{\pm I\}$. Since the complex dimension is 3, it follows that the real dimension of $\mathrm{PSL}(2, \mathbb{C})$ is 6.

Proof of Lemma 1.5. It is easy to check that $\mathcal{G}(\hat{\mathbb{C}})$ contains this group of fractional linear transformations as a subgroup. After composing the given $g \in \mathcal{G}(\hat{\mathbb{C}})$ with a suitable element of this subgroup, we may assume that $g(0) = 0$ and $g(\infty) = \infty$. But then the quotient $g(z)/z$ is a bounded holomorphic function from $\mathbb{C} \setminus \{0\}$ to itself. (In fact, $g(z)/z$ tends to the nonzero finite value $g'(0)$ as $z \rightarrow 0$. Setting $\zeta = 1/z$ and $G(\zeta) = 1/g(1/\zeta)$, evidently $g(z)/z = \zeta/G(\zeta)$ tends to the nonzero finite value $1/G'(0)$ as $z \rightarrow \infty$.) Setting $z = e^w$, it follows that the composition $w \mapsto g(e^w)/e^w$ is a bounded holomorphic function on $\mathbb{C}$. Hence it takes a constant value c by Liouville's Theorem. Therefore $g(z) = cz$ is linear, and hence g itself is an element of $\mathrm{PSL}(2, \mathbb{C})$. $\square$

Next we will show that both $\mathcal{G}(\mathbb{C})$ and $\mathcal{G}(\mathbb{D})$ can be considered as Lie subgroups of $\mathcal{G}(\hat{\mathbb{C}})$.

Corollary 1.6 (The Affine Group). *The group $\mathcal{G}(\mathbb{C})$ of all conformal automorphisms of the complex plane consists of all affine transformations*

$$f(z) = \lambda z + c$$

with complex coefficients $\lambda \neq 0$ and c .

Proof. First note that every conformal automorphism f of $\mathbb{C}$ extends uniquely to a conformal automorphism of $\hat{\mathbb{C}}$. In fact $\lim_{z \rightarrow \infty} f(z) = \infty$, so the singularity of $1/f(1/\zeta)$ at $\zeta = 0$ is removable. (Compare Ahlfors [1966, p. 124].) It follows that $\mathcal{G}(\mathbb{C})$ can be identified with the subgroup of $\mathcal{G}(\hat{\mathbb{C}})$ consisting of Möbius transformations which fix the point ∞ . Evidently this