

Society for  
Applied Bacteriology  
Symposium Series  
No. 2

# ACTINOMYCETALE

## Characteristics and Practical Importance

edited by  
G. Sykes and F. A. Skinner

THE SOCIETY FOR APPLIED BACTERIOLOGY  
SYMPOSIUM SERIES NO. 2

ACTINOMYCETALES:  
CHARACTERISTICS AND  
PRACTICAL IMPORTANCE

*Edited by*

G. SYKES

AND

F. A. SKINNER

1973



ACADEMIC PRESS · LONDON · NEW YORK

ACADEMIC PRESS INC. (LONDON) LTD  
24-28 OVAL ROAD  
LONDON N.W.1

*U.S. Edition published by*  
ACADEMIC PRESS INC.  
111 FIFTH AVENUE  
NEW YORK, NEW YORK 10003

Copyright © 1973 By the Society for Applied Bacteriology

ALL RIGHTS RESERVED

NO PART OF THIS BOOK MAY BE REPRODUCED IN ANY FORM BY PHOTOSTAT,  
MICROFILM, OR ANY OTHER MEANS, WITHOUT WRITTEN PERMISSION FROM  
THE PUBLISHERS

*Library of Congress Catalog Card Number: LCCCN 72-9777*  
*ISBN: 0-12-679950-4*

Printed in Great Britain by  
The Whitefriars Press Ltd., London and Tonbridge, England

## Contributors

- NINA S. AGRE, *Laboratory for Physiology of Actinomycetes, Institute of Microbiology, USSR Academy of Sciences, Moscow, U.S.S.R.*
- R. W. ATTWELL, *Postgraduate School of Studies in Biological Sciences, University of Bradford, Bradford BD7 1DP, England*
- H. BEERENS, *Institut Pasteur, 20, Boulevard Louis XIV, 59000 Lille, France*
- W. BLYTH, *Department of Botany, University of Edinburgh, Edinburgh EH9 3JH Scotland*
- G. H. BOWDEN, *The London Hospital Medical College. Dental School, Turner Street, London, England*
- R. M. BRADSHAW, *Hartley Botanical Laboratories, University of Liverpool, Liverpool, England*
- CATHERINE L. BULLEN, *Public Health Laboratory, Lewsey Road, Luton, England*
- N. P. BURMAN, *Metropolitan Water Board, Rosebery Avenue, London EC1R 4TP, England*
- M. CATTEAU, *C.E.R.T.I.A., 369, Rue Jules Guesde, Flers Bourg, 59650 Villeneuve d'Ascq, France*
- T. CROSS, *Postgraduate School of Studies in Biological Sciences, University of Bradford, Bradford BD7 1DP, England*
- M. GOODFELLOW, *Department of Microbiology Medical School, University of Newcastle-upon-Tyne, Newcastle-upon-Tyne NE1 7RU, England*
- D. GOTTLIEB, *Department of Plant Pathology, University of Illinois, Urbana, Illinois 61801, U.S.A.*
- J. M. HARDIE, *The London Hospital Medical College, Dental School, Turner Street, London, England*
- D. A. HOPWOOD, *John Innes Institute, Colney Lane, Norwich NOR 70F, England*
- L. V. KALAKOUTSKII, *Laboratory for Physiology of Actinomycetes, Institute of Microbiology, USSR Academy of Sciences, Moscow, U.S.S.R.*
- J. LACEY, *Rothamsted Experimental Station, Harpenden, Herts, England*
- D. H. LAPWOOD, *Rothamsted Experimental Station, Harpenden, Herts, England*
- J. LOSFELD, *U.E.R. Mathématiques, Université des Sciences, 59000 Lille, France*
- F. PONCELET, *C.E.R.T.I.A., 369, Rue Jules Guesde, Flers-Bourg, 59650 Villeneuve d'Ascq, France*
- L. M. POUZHARITSKAJA, *Laboratory for Physiology of Actinomycetes Institute of Microbiology, USSR Academy of Sciences, Moscow, U.S.S.R.*
- G. P. SHARPLES, *Hartley Botanical Laboratories, University of Liverpool, Liverpool, England*

- W. B. TURNER, *I.C.I. Ltd., Pharmaceuticals Division, Alderley Park, Macclesfield, Cheshire, England*
- KATHLEEN WILLIAMS, *Public Health Laboratory, Lewsey Road, Luton, England*
- S. T. WILLIAMS, *Hartley Botanical Laboratories, University of Liverpool, Liverpool, England*
- A. T. WILLIS, *Public Health Laboratory, Lewsey Road, Luton, England*

## Preface

THIS second volume in the Symposium series of the Society for Applied Bacteriology publications is concerned with a review of the characteristics and significance of the Actinomycetales. The Symposium was held at the Loughborough University of Technology, England in July 1972. Each chapter has been contributed by a specialist in his field and so the book as a whole constitutes a unique collection of information on this group of micro-organisms which has long been recognized as having many unusual and interesting features.

As might be expected of such a diverse group of organisms there is much to be learned of their physiological characteristics and several chapters are devoted to such studies. Under this heading there are valuable and comprehensive contributions on the taxonomy and genetics of the group and on the production and germination of their spores. In other chapters the occurrence and significance of the actinomycetes in water, soil and the human digestive tract, and their role in causing potato scab disease is discussed.

It is relevant to stress the medical importance of the actinomycetes both in causing and curing disease. The realization that some streptomycetes could produce antibacterial substances stimulated great interest in the group, especially in the years immediately following the Second World War, and led to the development of the present impressive array of antibiotics of clinical value. The pathogenic properties of the anaerobic actinomycetes have been known for a long time but more recently attention has been directed to the activities of the aerobic types as allergens in pulmonary conditions such as Farmer's Lung disease.

G. SYKES  
3 Grosvenor Court  
Egerton Road  
Weybridge  
Surrey, England

F. A. SKINNER  
Rothamsted Experimental Station  
Harpenden  
Hertfordshire  
England

January 1973

## Contents

### LIST OF CONTRIBUTORS

PREFACE	vii
---------	-----

### General Consideration and Implications of the Actinomycetales

D. GOTTLEIB

Introduction	1
Habitat	2
Soil competition	3
Antibiotics	4
Species pathogenic to man	5
Plant diseases	6
Animal nutrition	6
Food preservation	7
Co-operative taxonomy	7
References	9

### Taxonomy and Classification of the Actinomycetes

T. CROSS and M. GOODFELLOW

Introduction	11
Family Actinomycetaceae	13
Family Actinoplanaceae	30
Family Dermatophilaceae	38
Family Frankiaceae	40
Family Micromonosporaceae	41
Family Mycobacteriaceae	43
Family Nocardiaceae	57
Family Streptomyacetaceae	64
Family Thermoactinomycetaceae	76
Family Thermomonosporaceae	78
Genera in search of a family	88
References	91

### The Fine Structure of the Actinomycetales

S. T. WILLIAMS, G. P. SHARPLES and R. M. BRADSHAW

Introduction	113
Methods for examining fine structure	113
(a) Ultrathin sections	113
(b) Whole cell mounts	118

(c) Carbon replicas of whole cells	114
(d) Carbon replicas of freeze etched cells	114
(e) Negative staining of disrupted cells	114
(f) Scanning electron microscopy	114
General cell contents	114
Mesosomes	115
Cross walls	117
Modifications to basic cell structure	119
Flagella	120
Formation of spores	121
(a) Fragmentation of a sheathless hypha (group 1a)	123
(b) Fragmentation of a sheathed hypha (group 1b)	123
(c) Endogenous spore formation (group 2)	125
Acknowledgements	126
References	127

## Genetics of the Actinomycetales

### D. A. HOPWOOD

Introduction	131
An outline of the conjugation system in <i>Streptomyces coelicolor</i>	
A3(2)	132
(a) The fertility types	132
(b) Zygote genomes	135
(c) The conjugation process	136
(d) Conclusions	138
Other conjugational systems	138
(a) Other streptomycetes	138
(b) <i>Nocardia</i>	140
(c) <i>Micromonospora</i>	141
Transformation systems	141
(a) <i>Thermoactinomyces vulgaris</i>	141
(b) Streptomycetes	142
Actinomycete genetics and industrial problems	143
(a) Mutagenesis	143
(b) The predictive value of the linkage map	145
(c) Ultrafertility	146
(d) Interspecific and inter-strain recombination	147
(e) The desirability of harnessing transduction or transformation in strain improvement	149
Actinomycete genetics and the study of morphogenesis	150
Acknowledgements	150
References	150



**The Streptomyces Spore: Its Distinct Features and Germinal Behaviour**

L. V. KALAKOUTSKII and L. M. POUZHARITSKAJA

Introduction	155
Spore viability and survival	156
Resistance to deleterious agents	157
Spore structure and composition	158
(a) Structure	158
(b) Composition	160
(c) Metabolic activity	164
Germination	165
(a) Effect of environmental factors	165
(b) Germination events	168
Conclusion	170
References	171

**Endospores of Actinomycetes: Dormancy and Germination**

L. V. KALAKOUTSKII and NINA S. AGRE

Introduction	179
The spore	181
Germination stages	182
(a) Effects of low temperatures on activation	182
(b) Other activation treatments	184
(c) What changes occur during activation?	186
(d) Germination	189
(e) Heterogeneity of spore populations	190
Effect of repeated transfer of cultures on the ability of their spores to germinate	191
Concluding remarks	192
References	193

**Germination of Actinomycete Spores**

R. W. ATTWELL and T. CROSS

Introduction	197
Actinomycetes studied	198
Spore production	198
Appearance in microculture	199
Changes in optical density of spore suspensions	199
(a) Influence of suspending medium	200
(b) Ageing	200
(c) Heat treatment	201
(d) Effects of antibiotics	202
Discussion	202

Summary	205
Acknowledgements	206
References	206

## Secondary Metabolism with Special Reference to Actinomycetales

W. B. TURNER

Introduction	209
Relationship of primary and secondary metabolism	210
Biosynthetic origins of the secondary metabolites	210
(a) Incorporation of the intact glucose skeleton	211
(b) Compounds related to the nucleosides	211
(c) The shikimic acid pathway	211
(d) The polyketide and polypropionate routes	214
(e) Terpenes and steroids	215
(f) Compounds derived from amino acids	216
Conclusions	216
References	217

## The Occurrence and Significance of Actinomycetes

N. P. BURMAN

Introduction	219
Methods	219
Occurrence and significance	221
(a) In river water	221
(b) In stored water	222
(c) In slow sand filters	223
(d) In biological sludge blankets	227
(e) In chlorinated water	227
(f) In the distribution system	227
References	229

## Actinomycetes in Soils, Composts and Fodders

J. LACEY

Introduction	231
Methods of assessment and their interpretation	232
Soils	233
(a) Numbers and types of actinomycetes	233
(b) State of actinomycetes in soil	235
(c) Factors influencing actinomycetes in soil	235
Manure and composts	237
(a) Animal manures: mushroom composts	238
(b) Vegetable composts	240
(c) Town wastes	240

Fodders . . . . .	240
(a) Hay . . . . .	240
(b) Cereal grains . . . . .	242
(c) Straw . . . . .	242
Other materials . . . . .	242
(a) Bagasse . . . . .	242
(b) Miscellaneous . . . . .	243
The role of actinomycetes . . . . .	243
(a) Interactions with other organisms . . . . .	244
(b) Antibiotic production . . . . .	245
(c) Degradation of organic matter . . . . .	245
Conclusion . . . . .	246
References . . . . .	246

### ***Streptomyces scabies* and Potato Scab Disease**

D. H. LAPWOOD

The disease . . . . .	253
(a) General . . . . .	253
(b) Symptoms . . . . .	253
The pathogen . . . . .	254
(a) Identity . . . . .	254
(b) Isolation and culture . . . . .	255
Infection and disease development . . . . .	255
Factors affecting the pathogen and disease incidence . . . . .	256
Control measures . . . . .	257
(a) Seedborne inoculum . . . . .	258
(b) Soilborne inoculum . . . . .	258
(c) Prevention of infection . . . . .	258
References . . . . .	259

### **Farmer's Lung Disease**

W. BLYTH

Introduction . . . . .	261
Incidence . . . . .	262
Organisms known to cause farmer's lung and their occurrence . . . . .	263
Inhalation, retention and elimination of respiratory allergens . . . . .	263
Sputum culture . . . . .	265
Common types of respiratory hypersensitivity . . . . .	265
Clinical features . . . . .	266
Histopathological features . . . . .	267

Serological evidence . . . . .	268
(a) Mouldy hay and thermophilic actinomycetes . . . . .	268
(b) Fungal antigens . . . . .	271
Relationships between serology and pathogenesis . . . . .	271
Treatment and control . . . . .	273
References . . . . .	273

### Commensal and Pathogenic *Actinomyces* Species in Man

G. H. BOWDEN and J. M. HARDIE

Introduction . . . . .	277
Early studies on the nature of the pathogen in actinomycosis . . . . .	277
<i>Actinomyces</i> as commensal organisms in man . . . . .	279
<i>Actinomyces</i> as pathogens . . . . .	281
(a) In man . . . . .	281
(b) Experimental infections . . . . .	283
Some characteristics of <i>Actinomyces</i> species . . . . .	284
(a) Morphology . . . . .	285
(b) Colonial . . . . .	286
(c) Cellular . . . . .	286
(d) Physiological . . . . .	286
Serology of <i>Actinomyces</i> species . . . . .	290
References . . . . .	295

### A Rapid Method for the Identification of *Bifidobacterium* Species Using 50 Characters

M. CATTEAU, F. PONCELET, H. BEERENS and J. LOSFELD

Introduction . . . . .	301
Matériel et Méthodes . . . . .	301
(a) Souches . . . . .	301
(b) Préparation des cultures . . . . .	302
(c) Encemement des galeries . . . . .	303
(d) Exploitation des résultats . . . . .	303
Résultats . . . . .	303
Discussion et Conclusions . . . . .	309
Références . . . . .	309

### The Significance of *Bifidobacteria* in the Intestinal Tract of Infants

CATHERINE L. BULLEN, A. T. WILLIS and KATHLEEN WILLIAMS

Introduction . . . . .	311
The composition of breast milk and cows' milk . . . . .	313
(a) Important constituents . . . . .	313

Some properties of breast milk and cows' milk . . . . .	314
(a) Coagulability . . . . .	314
(b) Buffering capacity of breast milk and cows' milk . . . . .	314
Growth of <i>Escherichia coli</i> and anaerobic Lactobacilli in breast milk and cows' milk . . . . .	314
Growth of anaerobic lactobacilli and <i>E. coli</i> in lactose broth . . . . .	316
Continuous culture of anaerobic lactobacilli and <i>E. coli</i> in breast milk and cows' milk . . . . .	316
Decrease of buffering capacity of cows' milk . . . . .	318
Growth in breast milk and cows' milk preparations . . . . .	319
The origin of anaerobic lactobacilli in the infant intestine . . . . .	319
Anaerobic lactobacilli in the vaginae of pregnant women and the mouths of newborn infants . . . . .	320
Discussion . . . . .	321
Acknowledgements . . . . .	324
References . . . . .	324

Techniques for the Isolation and Characterization of <i>Actinomyces</i> and <i>Bifidobacterium</i> Species: Report of a Panel Discussion . . . . .	327
---	-----

Techniques for the Isolation and Identification of Aerobic Actino- mycetales: Summary of the Discussion Session . . . . .	335
--	-----

# General Consideration and Implications of the Actinomycetales

D. GOTTLIEB

*Department of Plant Pathology, University of Illinois,  
Urbana, Illinois 61801, U.S.A.*

## CONTENTS

1. Introduction	1
2. Habitat	2
3. Soil competition	3
4. Antibiotics	4
5. Species pathogenic to man	5
6. Plant diseases	6
7. Animal nutrition	6
8. Food preservation	7
9. Co-operative taxonomy	7
10. References	9

## 1. Introduction

THE ACTINOMYCETES have always been a strange group of organisms to the bacterial taxonomists. The number of investigators studying this group of microbes has always been small and for the most part these organisms have been neglected by medical bacteriologists, physiologists and biochemists. Such affairs were to be expected when one recalls that until recently not even their taxonomic position was certain (Breed, Murray & Smith, 1957). Bacteriologists considered them as bacteria and mycologists generally considered them as fungi. The diseases caused by them have been described in books on medical mycology (e.g. Conant *et al.*, 1954). This era is now over and the actinomycetes are generally accepted as bacteria. They have no nuclear membrane, are sensitive to lysozyme for the most part and to the common antibacterial agents; there is a similarity to the type of bacterial flagella when these organelles are present and the types of cell wall resemble those of bacteria. The hyphal diameters are much smaller than those of fungi and are close to those of the bacteria.

Genera in actinomycetes consist of varied groups of bacteria whose common feature is the formation of hyphae at some stage of development. This tendency is tenuous and often requires imagination to believe in it. In some species of the genus *Mycobacterium* hyphae are never seen, whereas other species have incipient hyphae only in young cultures. An even worse situation exists in the various genera of the Actinomycetaceae; these form microcolonies which have very transitory filaments that are extremely difficult to see and the dominant forms are diphtheroid, bifid or even rod shaped. Hyphal development is more

pronounced in the nocardias but they, too, fragment. The time at which this occurs varies with the species and it determines the types of colony which form. Colonies in which the filaments break up early tend to be soft or mucoid whereas those whose filaments break up after a few days have time for a branched mycelium to form and develop a leathery texture: they later fragment to bacillary or coccoid elements. In the Streptomycetaceae, filament development is strong and fragmentation is rarely found. Examples are in the *Streptomyces*, *Micromonospora* and *Actinoplanes* spp.

Reproduction is usually asexual, though sexual processes have been shown to occur by genetic analyses. In the nonhyphal forms, asexual reproduction is by fragmentation or perhaps even by the usual fission of single cells. Where stable hyphae are produced, vegetative reproduction is by well formed spores resembling fungal arthrospores, borne either free or in sporangia, as in the Actinoplanaceae. The free spores are in the form of sporophores and may consist of 1, 2 or many spored chains arising singly or verticillate from primary hyphae. The long chains are straight, looped or spiral. Except for the Actinomycetaceae, which contain aerobic or micro-aerophilic genera, the actinomycetes are generally aerobic.

## 2. Habitat

Members of the actinomycetes which live a saprophytic existence can be found with greater or less frequency in most ecological settings, soil having the greatest population density. Numerically they are less dominant than the other bacteria and more prominent than the fungi. The numbers vary a great deal but it is not uncommon in fertile soils from many different parts of the world to find  $-5 \times 10^6$  propagules/g of soil (Taber, 1960; Gottlieb, unpublished). They occur in greatest numbers in the top few inches of the soil and decrease with depth (Waksman & Purvis, 1932). The sensitivity of *Streptomyces* spp. to acids at pH 5.0 or less precludes high populations in such soils (Taber, 1960). Similarly, the writer has not found them in soils from the Arctic tundra of northern Sweden or in soils of the Great Salt Desert of the U.S.A. High densities of spores can also be found on potato or sweet potato that have been infected by pathogenic *Streptomyces* spp. *Streptomyces* spp. are also common on most plant parts and, being dustborne, can be isolated from almost all artifacts and natural materials (Grein & Meyer, 1958). They are found in marine littoral and associated with *Micromonospora* spp., especially in fresh water lakes and in bottom muds. By their nature thermophilic forms, are found commonly in composts of manures. Other types are favoured by particular oxygen environments; *Actinomyces israeli*, for example, is anaerobic, whilst a number of other species are micro-aerophilic.

The streptomycetes have 3 important roles in the soil. One is their function in decomposing the organic matter of the soil (Waksman & Lomanitz, 1925;

Reynolds, 1954; Williams, 1966). Another is their effect on soil structure in binding clay particles by the hyphal threads to impart a granular viable structure that is conducive to crop production. The actinomycetes are undoubtedly responsible in part for the earthy odour of soil (Gerber & Lechevalier, 1965).

### 3. Soil Competition

The third interesting role of actinomycetes in soil is their ability to produce antibiotics which are inimical to other soil microbes. This is on the assumption that antibiotics are produced in normal untreated soils in the same way as they are produced in special laboratory media. This results in an intriguing hypothesis to explain some of the competitive effects in soil, resulting in the increase of some species and loss of other species among the microbial components of the soil. Competition certainly occurs between organisms in soil, but it is difficult to demonstrate that the actinomycetes produce in untreated soils antibiotics at levels high enough to inhibit other soil organisms. There are data to show that actinomycetes, especially members of the genus *Streptomyces*, can produce antibiotics in some sterilized soils, with or without added metabolites, and even in nonsterile soils, but the question remains: which of these conditions represents the natural state of the soil? Two main considerations are (1) natural soils do not have the necessary metabolites in concentrations necessary to stimulate and allow antibiotic production and (2) the necessary metabolites are normal in soil and are part of the natural life cycles in soils. An added difficulty is that of assessing the 'clinical' level of antibiotic concentration in soil.

A few examples clarify the situation. The addition of *S. griseus* and *Bacillus subtilis* to normal or treated sterile soil resulted in a decrease in the population of the *Bacillus* but there was no detectable amount of streptomycin. Moreover if streptomycin had been produced it would have been adsorbed by the negatively charged clay component of the soil to render it inactive (Siminoff & Gottlieb, 1951). On the other hand *S. venezuelae* produces chloromycetin, a neutral molecule. This actinomycete produced the antibiotic in trace amounts in sterile, but not in nonsterile soil, and the amounts were increased by adding green alfalfa hay or tryptone to the soil (Gottlieb & Siminoff, 1952). The antibiotic was not found in field plots that had been inoculated with the streptomycete nor in any of the 91 soils that were collected in field surveys, including areas from which the organism had originally been isolated (Ehrlich, Anderson, Coffey & Gottlieb, 1952, 1953). Actidione, another neutral antibiotic produced by *S. griseus* gave similar results (Gottlieb, Siminoff & Martin, 1952). The results of studies with *S. rimosus* and *S. aureofaciens*, which produce the amphoteric compounds terramycin and aureomycin, respectively, are different (Martin & Gottlieb, 1952). The actinomycetes inhibited *B. polymyxa* in soil even though relatively high concentrations of the antibiotics did not. In this case the



inactivation or removal from soil of the antibiotics was presumably due to the pH value of the soil.

There appears to be no correlation between the ability of an organism to inhibit the growth of other organisms in soil and its ability to produce an antibiotic that inhibits them. As a part of one of our studies, 15 actinomycetes were isolated that produced an antibiotic which inhibited *Rhizoctonia solani* and 15 others that did not. The actinomycetes were separately added to sterile soil 2 weeks before, at the same time as, and 2 weeks after, the *Rhizoctonia*. After 1 month in pots the relative populations of *Rhizoctonia* in the soils were determined by the numbers of radish and pea seedlings that did not emerge or were damped off, and there were no significant differences between groups. Some isolates that produced antibiotics antagonized the fungus, whereas others did not. Similarly some that did not produce antibiotics inhibited the test fungus whereas others did not.

#### 4. Antibiotics

It is because of their ability to produce antibiotics that the actinomycetes are perhaps best known. Except for a few that are produced by fungi, e.g. penicillin and cephalosporin, and a few that are produced by bacteria, e.g. bacitracin and polymyxin, all other antibiotics that are medically useful and have a wide application are synthesized by actinomycetes. A compilation of the microbial sources of antibiotics discovered in the U.S.A. and Japan between 1953 and 1970 reveal that c. 85% are produced by actinomycetes, 11% by fungi and 4.5% by bacteria. These differences may not reflect the actual proclivities of the different groups to produce antibiotics for similar ratios might be obtained if more effort was spent on the last groups. Yet the decision, if it were so made, to spend most time on the actinomycetes is in my experience the result of the fact that the actinomycetes have shown the greatest potential for producing new and useful antibiotics. The exact number of antibiotics that are known is difficult to even estimate. About 2700 substances have been reported but I would estimate that c. 50% of them are so poorly characterized that their uniqueness is suspect. The most commonly reported new antibiotics are those with antibacterial activity, the proportion in Japan being 48% and in the U.S.A. 56%; next are antifungal agents, in Japan, 32% and in the U.S.A. 24%. Antitumour antiviral and antiprotozoal antibiotics comprise the remainder. These figures probably reflect not so much the innate ability of microbes to produce such agents but the amount of research effort that has been placed in the search for antibiotics of different therapeutic abilities.

Another means of determining the importance of antibiotic production is by the amount and value of the product that is produced. With these data one can get into ethical difficulties such as the value of saving one life against the value