


DISCRETE MATHEMATICS AND ITS APPLICATIONS

Graph Polynomials



Yongtang Shi
Matthias Dehmer
Xueliang Li
Ivan Gutman



CRC Press
Taylor & Francis Group

A CHAPMAN & HALL BOOK

Graph Polynomials

Yongtang Shi
Matthias Dehmer
Xueliang Li
Ivan Gutman



CRC Press

Taylor & Francis Group

Boca Raton London New York

CRC Press is an imprint of the
Taylor & Francis Group, an **informa** business
A CHAPMAN & HALL BOOK

CRC Press
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

© 2017 by Taylor & Francis Group, LLC
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works

Printed by CPI on sustainably sourced paper
Version Date: 20161019

International Standard Book Number-13: 978-1-4987-5590-0 (Hardback)

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Library of Congress Cataloging-in-Publication Data

Names: Shi, Yongtang.
Title: Graph polynomials / [edited by] Yongtang Shi [and three others].
Description: Boca Raton : CRC Press, [2017] | Series: Discrete mathematics and its applications | Includes bibliographical references and index.
Identifiers: LCCN 2016031173 | ISBN 9781498755900 (hardback) | ISBN 9781315367996 (e-book) | ISBN 9781498755917 (e-book)
Subjects: LCSH: Graph theory. | Combinatorial analysis. | Polynomials.
Classification: LCC QA166 .G69878 2017 | DDC 511/.5--dc23
LC record available at <https://lccn.loc.gov/2016031173>

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>

and the CRC Press Web site at
<http://www.crcpress.com>

Graph Polynomials

DISCRETE MATHEMATICS AND ITS APPLICATIONS

R. B. J. T. Allenby and Alan Slomson, How to Count: An Introduction to Combinatorics, Third Edition

Craig P. Bauer, Secret History: The Story of Cryptology

Juergen Bierbrauer, Introduction to Coding Theory

Katalin Bimbó, Combinatory Logic: Pure, Applied and Typed

Katalin Bimbó, Proof Theory: Sequent Calculi and Related Formalisms

Donald Bindner and Martin Erickson, A Student's Guide to the Study, Practice, and Tools of Modern Mathematics

Francine Blanchet-Sadri, Algorithmic Combinatorics on Partial Words

Miklós Bóna, Combinatorics of Permutations, Second Edition

Miklós Bóna, Handbook of Enumerative Combinatorics

Miklós Bóna, Introduction to Enumerative and Analytic Combinatorics, Second Edition

Jason I. Brown, Discrete Structures and Their Interactions

Richard A. Brualdi and Dragoš Cvetković, A Combinatorial Approach to Matrix Theory and Its Applications

Kun-Mao Chao and Bang Ye Wu, Spanning Trees and Optimization Problems

Charalambos A. Charalambides, Enumerative Combinatorics

Gary Chartrand and Ping Zhang, Chromatic Graph Theory

Henri Cohen, Gerhard Frey, et al., Handbook of Elliptic and Hyperelliptic Curve Cryptography

Charles J. Colbourn and Jeffrey H. Dinitz, Handbook of Combinatorial Designs, Second Edition

Abhijit Das, Computational Number Theory

Matthias Dehmer and Frank Emmert-Streib, Quantitative Graph Theory: Mathematical Foundations and Applications

Martin Erickson, Pearls of Discrete Mathematics

Martin Erickson and Anthony Vazzana, Introduction to Number Theory

Titles (continued)

Steven Furino, Ying Miao, and Jianxing Yin, Frames and Resolvable Designs: Uses, Constructions, and Existence

Mark S. Gockenbach, Finite-Dimensional Linear Algebra

Randy Goldberg and Lance Riek, A Practical Handbook of Speech Coders

Jacob E. Goodman and Joseph O'Rourke, Handbook of Discrete and Computational Geometry, Second Edition

Jonathan L. Gross, Combinatorial Methods with Computer Applications

Jonathan L. Gross and Jay Yellen, Graph Theory and Its Applications, Second Edition

Jonathan L. Gross, Jay Yellen, and Ping Zhang Handbook of Graph Theory, Second Edition

David S. Gunderson, Handbook of Mathematical Induction: Theory and Applications

Richard Hammack, Wilfried Imrich, and Sandi Klavžar, Handbook of Product Graphs, Second Edition

Darrel R. Hankerson, Greg A. Harris, and Peter D. Johnson, Introduction to Information Theory and Data Compression, Second Edition

Darel W. Hardy, Fred Richman, and Carol L. Walker, Applied Algebra: Codes, Ciphers, and Discrete Algorithms, Second Edition

Daryl D. Harms, Miroslav Kraetzl, Charles J. Colbourn, and John S. Devitt, Network Reliability: Experiments with a Symbolic Algebra Environment

Silvia Heubach and Toufik Mansour, Combinatorics of Compositions and Words

Leslie Hogben, Handbook of Linear Algebra, Second Edition

Derek F. Holt with Bettina Eick and Eamonn A. O'Brien, Handbook of Computational Group Theory

David M. Jackson and Terry I. Visentin, An Atlas of Smaller Maps in Orientable and Nonorientable Surfaces

Richard E. Klima, Neil P. Sigmon, and Ernest L. Stitzinger, Applications of Abstract Algebra with Maple™ and MATLAB®; Second Edition

Richard E. Klima and Neil P. Sigmon, Cryptology: Classical and Modern with Maplets

Patrick Knupp and Kambiz Salari, Verification of Computer Codes in Computational Science and Engineering

William Kocay and Donald L. Kreher, Graphs, Algorithms, and Optimization

Donald L. Kreher and Douglas R. Stinson, Combinatorial Algorithms: Generation Enumeration and Search

Hang T. Lau, A Java Library of Graph Algorithms and Optimization

C. C. Lindner and C. A. Rodger, Design Theory, Second Edition

San Ling, Huaxiong Wang, and Chaoping Xing, Algebraic Curves in Cryptography

Nicholas A. Loehr, Bijective Combinatorics

Toufik Mansour, Combinatorics of Set Partitions

Titles (continued)

Toufik Mansour and Matthias Schork, Commutation Relations, Normal Ordering, and Stirling Numbers

Alasdair McAndrew, Introduction to Cryptography with Open-Source Software

Elliott Mendelson, Introduction to Mathematical Logic, Fifth Edition

Alfred J. Menezes, Paul C. van Oorschot, and Scott A. Vanstone, Handbook of Applied Cryptography

Stig F. Mjølsnes, A Multidisciplinary Introduction to Information Security

Jason J. Molitierno, Applications of Combinatorial Matrix Theory to Laplacian Matrices of Graphs

Richard A. Mollin, Advanced Number Theory with Applications

Richard A. Mollin, Algebraic Number Theory, Second Edition

Richard A. Mollin, Codes: The Guide to Secrecy from Ancient to Modern Times

Richard A. Mollin, Fundamental Number Theory with Applications, Second Edition

Richard A. Mollin, An Introduction to Cryptography, Second Edition

Richard A. Mollin, Quadratics

Richard A. Mollin, RSA and Public-Key Cryptography

Carlos J. Moreno and Samuel S. Wagstaff, Jr., Sums of Squares of Integers

Gary L. Mullen and Daniel Panario, Handbook of Finite Fields

Goutam Paul and Subhamoy Maitra, RC4 Stream Cipher and Its Variants

Dingyi Pei, Authentication Codes and Combinatorial Designs

Kenneth H. Rosen, Handbook of Discrete and Combinatorial Mathematics

Yongtang Shi, Matthias Dehmer, Xueliang Li, and Ivan Gutman, Graph Polynomials

Douglas R. Shier and K.T. Wallenius, Applied Mathematical Modeling: A Multidisciplinary Approach

Alexander Stanoyevitch, Introduction to Cryptography with Mathematical Foundations and Computer Implementations

Jörn Steuding, Diophantine Analysis

Douglas R. Stinson, Cryptography: Theory and Practice, Third Edition

Roberto Tamassia, Handbook of Graph Drawing and Visualization

Roberto Togneri and Christopher J. deSilva, Fundamentals of Information Theory and Coding Design

W. D. Wallis, Introduction to Combinatorial Designs, Second Edition

W. D. Wallis and J. C. George, Introduction to Combinatorics

Jiacun Wang, Handbook of Finite State Based Models and Applications

Lawrence C. Washington, Elliptic Curves: Number Theory and Cryptography, Second Edition

Preface

Graph polynomials have been developed for measuring structural information of networks using combinatorial graph invariants and for characterizing graphs. Various problems in graph theory and discrete mathematics can be treated and solved in a rather efficient manner by making use of polynomials. Various graph polynomials have been proven useful in discrete mathematics, engineering, information sciences, mathematical chemistry, and related disciplines.

In general, graph polynomials encode graph-theoretical information of the underlying graph in various ways. A particular graph polynomial may be interesting if it encodes useful graph parameters. An example is the Wiener polynomial whose coefficients are based on distances in a graph. Then for simple graph classes, the Wiener polynomial provides a simple and compact characterization of these graphs which can be analyzed quantitatively. In addition, graph polynomials and their zeros have been a valuable source for investigating various problems in discrete mathematics and related areas. It seems that graph polynomials were first introduced by J.J. Sylvester in 1878, and further studied by J. Petersen. Until now, there have been plenty of graph polynomials, such as the chromatic polynomial, characteristic polynomial, matching polynomial, Tutte polynomial, Whitney (rank) polynomial, Jones polynomial, Hosoya polynomial, Wiener polynomial, distance polynomial, edge-difference polynomial, independence polynomial, adjacency polynomial, flow polynomial, knot polynomial, topological transition polynomial, interlace polynomial, permanent polynomial, homomorphism polynomial, and so on. While studying these polynomials, it is crucial to study their particular properties such as location of zeros, interpretation of zeros, and so forth.

The main goal of this book is to present how graph polynomials characterize graph parameters efficiently, by emphasizing theoretical and practical problems. In the past few decades, many graph polynomials have been studied and plenty of theoretical and practical approaches have been developed. The topics addressed in this book cover a broad range of concepts and methods in terms of graph polynomials. The topics range from analyzing mathematical properties of graph polynomials to applying the polynomials in several application areas. By covering this broad range of topics, the book aims to fill a gap in contemporary literature in disciplines such as applied mathematics, information sciences, and mathematical chemistry.

Many colleagues, whether consciously or unconsciously, have provided us with input, help, and support before and during the preparation of this book. In particular, we thank Helge Mowshowitz, Frank Emmert-Streib, Zengqiang Chen, Bo Hu, Shailesh Tripathi, Abbtin Trinks, and Guihai Yu, and we apologize to all whose names have been inadvertently omitted. Also, we thank our editors, Sunil Nair and Alexander Edwards from CRC Press/Taylor & Francis Group, who have always been available and helpful. Last but not least, Yongtang Shi, Matthias Dehmer, and Xueliang Li thank the National Natural Science Foundation of China and Nankai University for their support. Matthias Dehmer thanks the Austrian Science Funds (project no. P26142) for supporting this work. Matthias Dehmer also thanks his sister, Marion Dehmer-Sehn, who passed away in 2012, for all her mental support.

To date, no book dedicated exclusively to graph polynomials has been published. Therefore, we hope this book will broaden the scope of the scientists who deal with topics related to graph polynomials rooted in graph theory, discrete mathematics, algebra, chemical graph theory, applied mathematics, computer science, information sciences, and related disciplines. Finally, we hope this book conveys the enthusiasm and joy we have for this field and inspires fellow researchers in their own practical or theoretical work.

Yongtang Shi

Tianjin

Matthias Dehmer

Munich

Xueliang Li

Tianjin

Ivan Gutman

Kragujevac

Contributors

Yvan Le Borgne

Centre National de la Recherche
Scientifique, Laboratoire Bordelais de
Recherche en Informatique, Université
Bordeaux 1
Talence, France

Walter Carballosa

Department of Mathematics
and Statistics
Florida International University
Miami, Florida

Mark Dukes

Department of Computer and
Information Sciences
University of Strathclyde
Glasgow, United Kingdom

Philip Feinsilver

Department of Mathematics
Southern Illinois University
Carbondale, Illinois

Ivan Gutman

Faculty of Science
University of Kragujevac
Kragujevac, Serbia

Tomer Kotek

Institut für Informationssysteme
Vienna University of Technology
Vienna, Austria

Bodo Lass

Centre National de la Recherche
Scientifique UMR 5208,
Institut Camille Jordan
Univ Lyon, Université Claude Bernard
Lyon 1
Villeurbanne, France

Wei Li

Department of Applied Mathematics
School of Science
Northwestern Polytechnical University
Xi'an, People's Republic of China

Xueliang Li

Center for Combinatorics
Nankai University
Tianjin, People's Republic of China

Shunyi Liu

College of Science
Chang'an University
Xi'an, People's Republic of China

Klas Markström

Department of Mathematics and
Mathematical Statistics
Umeå University
Umeå, Sweden

John P. McSorley

Department of Mathematics
Southern Illinois University
Carbondale, Illinois

Ada Morse

Department of Mathematics and Statistics
University of Vermont
Burlington, Vermont

José M. Rodríguez

Department of Mathematics
Carlos III University of Madrid
Madrid, Spain

Yongtang Shi

Center for Combinatorics
Nankai University
Tianjin, People's Republic of China

José M. Sigarreta

Faculty of Mathematics
Autonomous University of Guerrero
Acapulco, Mexico

William Staton

Department of Mathematics
University of Mississippi
Oxford, Mississippi

Yadira Torres-Nuñez

Department of Mathematics
Miami Dade College
Miami, Florida

Bing Wei

Department of Mathematics
University of Mississippi
Oxford, Mississippi

Tingzeng Wu

School of Mathematics and Statistics
Qinghai Nationalities University
Xining, People's Republic of China

Heping Zhang

School of Mathematics and Statistics
Lanzhou University
Lanzhou, People's Republic of China

Contents

Preface	ix
Contributors	xi
1 The Interlace Polynomial	1
<i>Ada Morse</i>	
2 Independence Polynomials of k-Trees and Compound Graphs	25
<i>William Staton and Bing Wei</i>	
3 New Aspects of the Abelian Sandpile Model on Graphs and Their Polynomials	41
<i>Mark Dukes and Yvan Le Borgne</i>	
4 Second Quantization of Recurrences	61
<i>Philip Feinsilver and John P. McSorley</i>	
5 A Survey on the Matching Polynomial	77
<i>Ivan Gutman</i>	
6 On the Permanental Polynomials of Graphs	101
<i>Wei Li, Shunyi Liu, Tingzeng Wu, and Heping Zhang</i>	
7 From the Ising and Potts Models to the General Graph Homomorphism Polynomial	123
<i>Klas Markström</i>	
8 Derivatives and Real Roots of Graph Polynomials	139
<i>Xueliang Li and Yongtang Shi</i>	
9 Logic-Based Computation of Graph Polynomials	151
<i>Tomer Kotek</i>	
10 Alliance Polynomial	177
<i>Walter Carballosa, José M. Rodríguez, José M. Sigarreta, and Yadira Torres-Nuñez</i>	
1 Graph Polynomials and Set Functions	207
<i>Bodo Lass</i>	
Index	239

Chapter 1

The Interlace Polynomial

Ada Morse

University of Vermont

Burlington, VT

1.1	Introduction	1
1.2	The Interlace Polynomial of a Graph	2
1.2.1	Preliminary definitions	2
1.2.2	Evaluations of the interlace polynomial	7
1.3	Connections to Other Polynomials	8
1.3.1	The Martin and Tutte polynomials	9
1.3.2	Isotropic systems and the Tutte–Martin polynomials	11
1.3.3	Isotropic matroids and the Tutte polynomial	13
1.4	Generalizations	14
1.4.1	Square matrices	15
1.4.2	Delta-matroids	16
1.5	Conclusion	20
	References	21

1.1 Introduction

The interlace polynomial of a graph arises in a number of settings both theoretical (e.g., isotropic systems) and applied (e.g., DNA sequencing by hybridization). We begin with the most straightforward, that of a recursive method for counting Eulerian circuits in two-in, two-out digraphs arising from an application in DNA sequencing. The interlace polynomial of a simple graph is obtained by generalizing the recursion used to solve this counting problem. We then discuss a closed form for the polynomial in terms of its adjacency matrix, the structure of which suggests definitions for analogous polynomials as well as a two-variable generalization. Another context in which the interlace polynomial arises is in isotropic systems, where it appears as a specialization of the Tutte–Martin polynomials, a connection we follow by way of the Martin polynomials of 4-regular graphs. Finally, we review generalizations of the polynomial to square matrices and delta-matroids.

In the context of counting Eulerian circuits in two-in, two-out digraphs, the interlace polynomial arose through Arratia et al.’s work on DNA sequencing [2]. In DNA sequencing by hybridization, the goal is to reconstruct a string of DNA knowing only information about its shorter substrings. The problem is to determine, from knowledge about the shorter substrings, whether a unique reconstruction exists.

More precisely, if $A = a_1a_2 \cdots a_m$ is a sequence consisting of m base pairs, the l -spectrum of A is the multiset containing all l -tuples consisting of l consecutive base pairs in A . Given the knowledge of the l -spectrum, the goal is to determine the number $k_l(m)$ of sequences of base pairs of length m having that l -spectrum.

In [2], the authors associate with a given l -spectrum its *de Bruijn graph*: a two-in two-out digraph D such that the Eulerian circuits of D are in bijection with sequences of base pairs having that l -spectrum. The problem, then, is to count the number of Eulerian circuits of D . This approach led to the discovery of a recursive formula for computing the number of Eulerian circuits of D based on an associated interlace graph. In [3], Arratia et al. generalized this recursion to define the interlace polynomial of an arbitrary simple graph.

The Eulerian circuits and cycle decompositions of 4-regular graphs have been an area of significant interest among graph theorists for many years, and approaches using graph polynomials have frequently proved fruitful [36,35,33,29]. The Martin polynomial [36,35], in particular, is closely related to the interlace polynomial as it counts, for any k , the number of k -component circuit partitions of a 4-regular graph.

This connection can be made explicit and, indeed, generalized. In a series of papers in the 1980s–1990s, Bouchet introduced the notion of an isotropic system to unify aspects of the study of 4-regular graphs and binary matroids [8,12,14], including a generalization of the Martin polynomials to this area [14]. Shortly after the discovery of the interlace polynomial, it was noticed that the interlace polynomial can be found as a specialization of the (restricted) Tutte–Martin polynomial of an isotropic system [13,1].

A connection between the interlace polynomial and the Tutte polynomial can be found by way of the Martin polynomial. However, this connection only captures the Tutte polynomial $t(G; x, y)$ for plane graphs when $x = y$, and so does not provide any strong link between the interlace polynomial and the many specializations of the Tutte polynomial, such as the chromatic polynomial. In recent work, Traldi has introduced a matroid associated with a graph, called its *isotropic matroid*, such that the interlace polynomial(s) of the graph can be recovered from parameterized Tutte polynomial(s) of its isotropic matroid [44].

Many generalizations of the interlace polynomial have been obtained. In [4], Arratia et al. defined a two-variable interlace polynomial of which the single-variable polynomial is a specialization. In doing so, they discovered, concurrently with Aigner and van der Holst [1], a closed form for the single-variable interlace polynomial in terms of its adjacency matrix. This closed form has a natural extension to arbitrary square matrices, and using a delta-matroid associated with the adjacency matrix of a graph, Brijder and Hoogeboom obtained a generalization of the interlace polynomial to delta-matroids [22]. In each case, the recursive definition of the interlace polynomial has also been generalized.

1.2 The Interlace Polynomial of a Graph

We begin by defining the interlace polynomial recursively by way of counting Eulerian circuits in two-in, two-out digraphs, and then discuss a closed form, an analogous polynomial, and a two-variable generalization. We conclude with selected evaluations of the interlace polynomial.

1.2.1 Preliminary definitions

We first establish some standard definitions and notation to be used throughout the chapter. Formally speaking, a *graph* G is a triple $(V(G), E(G), \phi)$ where $V(G)$ is a finite set of *vertices*, $E(G)$ is a finite set of *edges*, and ϕ is a function from $E(G)$ to $\{\{a, b\} : a, b \in V(G)\}$. Let $e \in E(G)$. If $\phi(e) = \{a\}$ is a singleton set, then e is said to be a *loop* and a is called a *looped vertex*. If $\phi(e) = \{a, b\}$, then we say e is an edge *between* a and b , a and b are

adjacent, a and b are the endpoints of e , and both a and b are incident to e . In general, we will suppress the function ϕ , define an edge by its endpoints, and define the graph G by the pair $(V(G), E(G))$. The *degree* of a vertex is the number of edges to which it is incident, counting loops twice. A graph is said to be k -regular if every vertex has degree k . If a and b are vertices with more than one edge between them, then we say there is a *multiple edge* between a and b . A graph is *simple* if it contains no loops or multiple edges. A *walk* in G is an alternating sequence of vertices and edges $v_1e_1v_2e_2\cdots v_ke_kv_{k+1}$ such that e_i is incident to v_i and v_{i+1} for each i . A walk is *closed* if its first and last vertices are the same. A closed walk repeating no edges is a *circuit*. A circuit in which no vertices are repeated is called a *cycle*. A *path* is a walk beginning and ending at distinct vertices and containing no repeated vertices. A graph is *connected* if for any vertices a and b there is a path from a to b . A *component* of a graph is a maximal connected subgraph. We denote by $k(G)$ the number of components of G . For $T \subseteq V(G)$ the *subgraph induced by T* , denoted $G[T]$, is the graph (T, E') where E' is the set of edges in $E(G)$ with both endpoints in T . If $v \in V(G)$, define $G \setminus v$ to be the subgraph of G induced by $V(G) \setminus \{v\}$.

The *adjacency matrix* of a graph G , denoted $A(G)$, is the $|V(G)| \times |V(G)|$ matrix over $GF(2)$ with rows and columns indexed by $V(G)$ defined by setting $A(G)_{ab} = 1$ if a and b are adjacent and 0 otherwise. For $T \subseteq V(G)$, define the *rank* $r(G[T])$ to be the matrix rank $r(A(G[T]))$, and the *nullity* $n(G[T])$ to be the matrix nullity $n(A(G[T]))$. By convention, $n(G[\emptyset]) = 0$. Note that for a graph G , the nullity and rank of G are sometimes defined by $n(G) = |E(G)| - |V(G)| + k(G)$ and $r(G) = |V(G)| - k(G)$.

A *digraph*, informally speaking, is a graph where each edge is given a direction, that is, the function ϕ has codomain $V(G) \times V(G)$. If a directed edge e goes from a to b , then a is called the *tail* of e and b is called the *head* of e . The indegree of a vertex in a digraph is the number of edges for which the vertex is the head, the outdegree the number for which it is the tail. A *two-in, two-out digraph* is a 4-regular digraph such that each vertex has both indegree and outdegree equal to 2. A walk $v_1e_1v_2e_2\cdots v_ke_kv_{k+1}$ in a digraph is *directed* or *consistently oriented* if e_i is directed from v_i to v_{i+1} for each i .

A *planar graph* is a graph that can be embedded in the plane (i.e., drawn in the plane by associating vertices to points and edges to curves between their endpoints, such that no two edges intersect other than at a shared endpoint). A *plane graph* is a planar embedding of a (planar) graph.

For sets A and B , the *symmetric difference* of A and B is $A\Delta B = (A \cup B) \setminus (A \cap B)$.

We can now begin defining the interlace polynomial by way of counting Eulerian circuits in two-in, two-out digraphs.

Definition 1.2.1 (interlace graph) Let G be a two-in, two-out digraph. A *Eulerian circuit* of G is a closed, directed walk of G containing each edge exactly once. Given a Eulerian circuit C of G , we say that vertices a and b are *interlaced* if the cycle visits them in the order $\dots a \dots b \dots a \dots b \dots$ and *noninterlaced* otherwise. The *interlace graph* or *circle graph* of C , denoted $H(C)$, is the graph whose vertices are the vertices of G with an edge between two vertices if they are interlaced in C (see Figure 1.1b and c).

Interlace graphs have been extensively studied [11,9,15,26,41,40] and were characterized by Bouchet in [9,15]. A particular focus of the area, due to a Gauss problem, has been characterizations of the interlace graphs arising from Eulerian cycles in plane 4-regular graphs [26,41,40].

There is a natural operation defined on Eulerian circuits of two-in, two-out digraphs in terms of this interlace relation.

Definition 1.2.2 At each vertex of a two-in, two-out digraph G , there are two possible (orientation consistent) pairings of in-edges and out-edges. For a pair of vertices a and b




FIGURE 1.1: Transpositions of Eulerian circuits and the interlace graph: (a) a two-in, two-out digraph G ; (b) a Eulerian cycle C in G ; (c) the interlace graph $H(C)$; (d) the Eulerian circuit C^{ab} ; (e) the interlace graph $H(C^{ab})$.

interlaced in a Eulerian circuit C of G , define the transposition C^{ab} to be the Eulerian circuit obtained by switching the pairing of edges at a and b (see Figure 1.1b and d).

The Eulerian circuits of G form a single orbit under the action of transposition, the proof of which can be found in [3] but was known previously in more general form in [39,47]. Observation of the effect on the interlace relation by performing the above operation to a Eulerian circuit leads to a corresponding definition for interlace graphs, presented here for graphs in general.

Definition 1.2.3 Let G be any graph. Let $v \in V(G)$. For any pair of vertices $a, b \in V(G)$, partition the remaining vertices of G into the following sets: (1) vertices adjacent to a and not b , (2) vertices adjacent to b and not a , (3) vertices adjacent to both a and b , and (4) vertices adjacent to neither a nor b . Define the pivot G^{ab} to be the graph obtained by inserting all possible edges between the first three of these sets, and deleting those that were already present in G (see Figure 1.2). Denote by G_{ab} the graph G with the labels of the vertices a and b swapped.

Although the above definition of pivot is attributable to Arratia et al. [2], the idea of the pivot appeared in the earlier work of Kotzig [34] on local complementations and the graph $(G^{ab})_{ab}$ is defined by Bouchet in [12] as the *complementation of G along the edge ab* . The precise connection to both is as follows:

Definition 1.2.4 Let G be a graph. For $v \in V(G)$, define the *open neighborhood* of v to be $N(v) = \{w \in V(G) \setminus \{v\} : w \text{ is adjacent to } v\}$. Note that $v \notin N(v)$ even if v is a looped vertex. We define the *local complement* $G * v$ to be the graph obtained from G