


Chemical Biology of Enzymes for Biotechnology and Pharmaceutical Applications

Enzyme Technologies

*Pluripotent Players in Discovering
Therapeutic Agents*

Edited by

Hsiu-Chiung Yang • Wu-Kuang Yeh • James R. McCarthy



ENZYME TECHNOLOGIES

Pluripotent Players in Discovering Therapeutic Agents

Edited by

HSIU-CHIUNG YANG


Eli Lilly and Company
Indianapolis, Indiana

WU-KUANG YEH

Indiana University Purdue University Indianapolis
Indianapolis, Indiana

JAMES R. McCARTHY

Indiana University Purdue University Indianapolis
Indianapolis, Indiana



WILEY

Copyright © 2014 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permission>.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Yang, Hsiu-Chiung.

Enzyme technologies : pluripotent players in discovering therapeutic agents / by Hsiu-Chiung Yang, Wu-Kuang Yeh, and J.R. McCarthy.

pages cm. – (Chemical biology of enzymes for biotechnology and pharmaceutical applications)

Includes index.

ISBN 978-0-470-28626-5 (hardback)

1. Enzyme inhibitors. 2. Enzymes—Pharmacokinetics 3. Drug design.

I. Yeh, Wu-Kuang, 1942– II. McCarthy, J. R. III. Title.

QP601.5.M33 2014

615.3–dc23

2013013968

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

ENZYME TECHNOLOGIES

**CHEMICAL BIOLOGY OF ENZYMES FOR
BIOTECHNOLOGY AND PHARMACEUTICAL
APPLICATIONS**

(A Series Consisting of Three Volumes)

Volume I. *Enzyme Technologies: Metagenomics, Evolution, Biocatalysis, and Biosynthesis*
Editors: Wu-Kuang Yeh, Hsiu-Chiung Yang, and James R. McCarthy

Volume II. *Enzyme Technologies: Pluripotent Players in Discovering Therapeutic Agents*
Editors: Hsiu-Chiung Yang, Wu-Kuang Yeh, and James R. McCarthy

Volume III. *Enzyme Technologies: Two Facets of Modern Structure-Based Design*
Editors: James R. McCarthy, Hsiu-Chiung Yang, and Wu-Kuang Yeh

CONTRIBUTORS

Herve Aloysius, MS, Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

Michael Beck, MD, University Medical Center, Children's Hospital, Mainz, Germany

Rathnam Chaguturu, PhD, Del Shankel Structural Biology Center, High Throughput Screening Laboratory, Lawrence, Kansas; SRI International, Harrisonburg, Virginia

Yu Chen, MS, Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

Ozlem Goker-Alpan, PhD, LSD Research and Treatment Unit, Center for Clinical Trials, Fairfax, Virginia

Longqin Hu, PhD, Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

Daigo Inoyama, BS, Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

Jutta Keller, MD, Department of Internal Medicine, Israelitic Hospital in Hamburg, Hamburg, Germany

Ivan S. Krylov, PhD, Department of Chemistry, University of Southern California, Los Angeles, California

Ley Nadine Lachawan, PhD, LSD Research and Treatment Unit, Center for Clinical Trials, Fairfax, Virginia

Peter Layer, MD, PhD, FACC, AGAF, Department of Internal Medicine, Israelitic Hospital in Hamburg, Hamburg, Germany

Gerald H. Lushington, PhD, Molecular Graphics and Modeling Laboratory, University of Kansas, Lawrence, Kansas; LiS Consulting, Lawrence, Kansas

James R. McCarthy, PhD, Department of Chemistry and Chemical Biology, Indiana University Purdue University Indianapolis, Indianapolis, Indiana

James McGee, PhD, Quantitative Biology, Eli Lilly and Company, Indianapolis, Indiana

Charles E. McKenna, PhD, Department of Chemistry, University of Southern California, Los Angeles, California

Shujaath Mehdi, PhD, Immunoinflammation Therapeutic Strategy Unit, Sanofi Pharmaceuticals, Bridgewater, New Jersey

Taichi Ohshiro, PhD, Department of Microbial Chemistry, Graduate School of Pharmaceutical Sciences, Kitasato University, Tokyo, Japan; Section on Lipid Sciences, Department of Pathology, Wake Forest University School of Medicine, Winston-Salem, North Carolina

Richard G. Peterson, PhD, PreClinOmics, Inc., Indianapolis, Indiana

Anuradha Roy, PhD, Del Shankel Structural Biology Center, High Throughput Screening Laboratory, Lawrence, Kansas

Henrike von Schassen, MD, Department of Internal Medicine, Israelitic Hospital in Hamburg, Hamburg, Germany

Hiroshi Tomoda, PhD, Department of Microbial Chemistry, Graduate School of Pharmaceutical Sciences, Kitasato University, Tokyo, Japan

Yanhui Yang, PhD, Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

Wu-Kuang Yeh, PhD, Indiana University Purdue University Indianapolis, Indianapolis, Indiana

Wei Zheng, PhD, National Center for Advancing Translational Sciences, National Institutes of Health, Bethesda, Maryland

PREFACE

The human genome was predicted to contain approximately 2742 genes that encode enzymes, which corresponds to 9.5% of the genome (HumanCyc version 7.5). These predicted enzymes can be subdivided into 1653 metabolic enzymes and 1089 nonmetabolic enzymes (including enzymes whose substrates are macromolecules, such as protein kinases and DNA polymerases). Enzymes play an important role in human physiology and the pathophysiology of disease. Understanding the function of an enzyme presents a significant opportunity for finding therapeutic agents. Therefore, a comprehensive understanding of all aspects of enzyme technology is critical in discovering therapeutic agents targeting enzymes, therapeutic enzymes, and enzyme-based applications, enabling research in drug discovery, chemistry, material science, and a vast number of other fields in science and technology.

This series on Chemical Biology of Enzymes for Biotechnology and Pharmaceutical Applications consists of three volumes. Volume I, *Enzyme Technologies: Metagenomics, Evolution, Biocatalysis, and Biosynthesis*, was published in 2010. Volume II, *Enzyme Technologies in Drug Discovery*, as listed in Volume I, has now been changed to *Enzyme Technologies: Pluripotent Players in Discovering Therapeutic Agents*. This book is intended both for biotech and pharmaceutical scientists in academic research institutes and industry as a comprehensive reference material for all common applications of enzyme technology in drug discovery. This book will be useful for biotechnology, biochemistry, molecular biology, and medicinal chemistry faculty members for teaching or conducting research in the field of enzyme technology. It is also a practical handbook for industrial scientists to study various aspects of enzyme technology and discover new treatments for unmet medical needs.

This book is divided into three parts: Part A: Enzymes – Essential Workhorses in Pharmaceutical Research; Part B: Enzymes – Indispensable Tools for Improving Druggability; and Part C: Enzymes – Powerful Weapons for Correcting Nature’s Errors. Part A consists of four chapters. Chapter 1, by Dr. A. Roy et al., discusses the principles of assay development and cutting-edge technologies available for protease assays, using proteases as a prototype. Chapter 2, by Drs. Ohshiro and Tomoda, provides a case study on the design and development of selective enzyme inhibitors, using lipid metabolizing enzymes as a prototype. There is a belief that covalent enzyme inhibitors (also called “irreversible inhibitors”) are not desirable for drug candidates. Chapter 3, by Dr. Mehdi, describes methods for characterizing covalent inhibitors and their therapeutic applications and explains how enzyme kinetics has been applied in drug discovery. Chapter 4, by Drs. Yeh and Peterson, provides a comprehensive coverage on various technologies that have been applied for *in vitro* enzymatic assays, as well as on common *in vivo* models to assess preclinical drug discovery for metabolic diseases. After going through this part, readers will have a better understanding as to how to select the best enzyme targets for drug discovery, the steps involved in designing enzyme inhibitors for therapeutic agents, and methods for evaluating selective enzyme inhibitors.

Part B consists of three chapters. It explains the principles of improving drug-gability and provides examples on how to utilize the properties of enzymes for designing therapeutic agents, specifically prodrugs. Chapter 5, by Dr. Hu et al., provides a comprehensive review of enzymes that are being, or can be, used to design prodrugs to improve druggability of existing drug molecules. Chapters 6 and 7 summarize case studies on the design of two successful prodrugs. Chapter 6, by Dr. McCarthy, provides a detailed approach and explains the hypothesis and rationale for the design and synthesis of a Gemcitabine prodrug. Chapter 7, by Drs. McKenna and Krylov, presents several examples of successful prodrug approaches that explicitly depend on enzyme-mediated activation. After going through this part, readers will have a complete understanding on how to select best target molecules for the prodrug approach to improve druggability and how to design successful prodrugs.

Part C consists of three chapters and provides a different viewpoint as to how enzymes can be used in pharmaceutical applications. There are many types of genetic in-born error disorders. Some of these disorders are involved in either deficiency or malfunctioning of specific enzymes. In such cases, functional enzymes can be reintroduced into patients. With recent advances in protein technology, there are several successful examples, such as the production of high-quality proteins and optimal methods for the delivery of these large molecules to the body. However, further improvements on either manufacturing or delivery of enzymes for therapeutics are still required. Chapter 8, by Dr. Beck, deals with Hunter’s syndrome, while Chapter 9, by Dr. Lachawan et al., discusses enzyme replacement therapy for Fabry disease. Chapter 10, by Dr. von Schassen et al., analyzes the level and activity of pancreatic enzymes as a means of diagnosing patients with pancreatic dysfunction, for example, Hunter’s syndrome. After

going through this part, readers will have an insight into how enzymes can be applied as therapeutic agents or diagnostic tools. Furthermore, readers may be able to identify possible enzyme targets to treat genetic disorders that still do not have effective medication.

*Eli Lilly and Company
Indiana University Purdue University Indianapolis
Indianapolis, Indiana*

HSIU-CHIUNG YANG
WU-KUANG YEH
JAMES R. MCCARTHY

CONTENTS

CONTRIBUTORS	vii
PREFACE	ix
PART A ENZYMES – ESSENTIAL WORKHORSES IN PHARMACEUTICAL RESEARCH	 1
1 Assay Technologies for Proteases	3
<i>Anuradha Roy, Gerald H. Lushington, James McGee, and Rathnam Chaguturu</i>	
2 Discovery and Development of Isozyme-Selective Inhibitors Involved in Lipid Metabolism	55
<i>Taichi Ohshiro and Hiroshi Tomoda</i>	
3 Covalent Enzyme Inhibition in Drug Discovery and Development	81
<i>Shujaath Mehdi</i>	
4 Preclinomics: Enzyme Assays and Rodent Models for Metabolic diseases	131
<i>Wu-Kuang Yeh and Richard G. Peterson</i>	
PART B ENZYMES – INDISPENSABLE TOOLS FOR IMPROVING DRUGGABILITY	 163
5 Enzymes and Targeted Activation of Prodrugs	165
<i>Yanhui Yang, Yu Chen, Herve Aloysius, Daigo Inoyama, and Longqin Hu</i>	

6	Evolution of an Orally Active Prodrug of Gemcitabine	237
	<i>James R. McCarthy</i>	
7	Enzymatically Activated Phosphate and Phosphonate Prodrugs	253
	<i>Ivan S. Krylov and Charles E. McKenna</i>	
PART C	ENZYMES – POWERFUL WEAPONS FOR CORRECTING NATURE’S ERRORS	301
8	Treatment Options for Mucopolysaccharidosis Type II (Hunter’s Syndrome)	303
	<i>Michael Beck</i>	
9	Enzyme Replacement Therapy for Fabry Disease	321
	<i>Ley Nadine Lacbawan, Wei Zheng, and Ozlem Goker-Alpan</i>	
10	Methods and Principles of Pancreatic Function Tests	335
	<i>Henrike von Schassen, Jutta Keller, and Peter Layer</i>	
	Index	341

PART A

ENZYMES – ESSENTIAL WORKHORSES IN PHARMACEUTICAL RESEARCH

1

ASSAY TECHNOLOGIES FOR PROTEASES

ANURADHA ROY

*Del Shankel Structural Biology Center, High Throughput Screening Laboratory,
Lawrence, Kansas*

GERALD H. LUSHINGTON

*Molecular Graphics and Modeling Laboratory, University of Kansas, Lawrence, Kansas; LiS
Consulting, Lawrence, Kansas*

JAMES MCGEE

Quantitative Biology, Eli Lilly and Company, Indianapolis, Indiana

RATHNAM CHAGUTURU

*Del Shankel Structural Biology Center, High Throughput Screening Laboratory,
Lawrence, Kansas; SRI International, Harrisonburg, Virginia*

I. INTRODUCTION

Proteases are ubiquitously expressed enzymes which catalyze hydrolysis of peptide bonds and work under a wide range of conditions using diverse catalytic mechanisms [1]. Proteases specifically cleave protein substrates either from the N- or C-termini (aminopeptidases and carboxypeptidases, respectively) or in the middle of the molecule (endopeptidases) [2]. Proteolytic enzymes modulate many physiological processes ranging from nonspecific hydrolysis of dietary proteins to highly specific and regulated proteolysis in cell cycle regulation, tissue remodeling,

Enzyme Technologies: Pluripotent Players in Discovering Therapeutic Agents, First Edition.

Edited by Hsiu-Chiung Yang, Wu-Kuang Yeh, and James R. McCarthy.

© 2014 John Wiley & Sons, Inc. Published 2014 by John Wiley & Sons, Inc.

blood coagulation, blood pressure control, angiogenesis, apoptosis, inflammation, ovulation, fertilization, and embryonic development [3,4]. Over 500 proteases each from humans, rat, mouse, and chimpanzee have been annotated and compiled in the Degradome database (<http://degradome.uniovi.es>) [5,6]. Information on all known proteases and their substrates/inhibitors is listed in the MEROPS database [7]. Based on the amino acid or metal that catalyzes the nucleophilic attack on substrate peptide bonds, the proteases are classified into five major types: aspartic (Asp), metallo-, cysteine (Cys), serine (Ser), and threonine (Thr) proteases. Aspartic and metalloproteases use an activated water molecule as a nucleophile to attack the peptide bond of the substrate, whereas in Cys, Ser, and Thr proteases, a catalytic amino acid residue (Cys, Ser, or Thr, respectively) serves as a nucleophile (Fig. 1). As a result, acyl-enzyme intermediates are formed only in the reactions catalyzed by Ser/Thr and Cys peptidases. Within each class of protease type are several enzymes that may have overlapping or distinct substrate recognition sites. Rawlings and Barrett proposed a classification of proteases into families based on amino acid sequence similarity, and families with similar three-dimensional folding are assembled into clans, indicating common ancestry [7,8]. The focus of this article is mainly on mammalian proteases and retroviral proteases which are of significant therapeutic relevance.

While pepsin in gastric juices digests a variety of proteins with broad specificity, renin is an example of Asp protease that shows high substrate specificity. Most proteases bind their substrates in fairly similar manner, first elucidated for papain by Schechter and Berger [9–11]. The catalytic site is flanked on one or both sides by sites that confer specificity of substrate binding to the protease and accommodate a side chain of an amino acid residue of the substrate. The enzymatic binding sites toward the N-terminus of the substrate are the non-prime side designated as S1, S2, ..., S_n from the catalytic site, and the residues C-terminal to the cleavage site are the prime side designated as S1', S2', ..., S_n' [8,12,13]. The amino acid residues in the protein substrate which correspond to their respective subsites are numbered P1, P2, ..., P_n and P1', P2', ..., P_n' (Fig. 1). Only few of the substrate binding sites have stringent specificities. For instance, site S1 confers specificity for Ser proteases and caspases, whereas the site S2, a hydrophobic subsite, defines specificity for the papain family of Cys proteases. In addition to the sites close to the catalytic site of the enzyme, distant sites on the enzyme may also contribute to the binding of substrates to the protease [9]. The specificity and biological activity of caspases are also determined by S4, which is distant from the catalytic site [14]. Proteolytic processing is being recognized as a mechanism for regulation of enzymatic activities, localization, and fate of proteins that are activated by limited and specific hydrolysis of peptide bonds. Dysregulation of proteolytic activity, structure, or expression results in major pathologies in the areas of cardiovascular diseases, cancer, neurodegenerative disorders, osteoporosis, diabetes type II, pancreatitis, inflammation, arthritis, and infectious diseases [4]. A large number of marketed drugs target the proteolytic enzymes that are involved in pathogenesis of various diseases [15] (Table 1). Although only a relatively small number of proteases are currently targeted for




FIGURE 1 Schematic representation of binding of substrate to a protease site. (a) The binding sites of the protease are numbered on either side of the scissile bond, with the non-primed sites (S1, S2, ..., Sn) located toward the amino-terminus of the substrate and S1' ... Sn' or the primed subsites toward the carboxy-terminus. (b) Structure of the peptide bond which is hydrolyzed by proteases and the two basic catalytic mechanisms for all types of protease hydrolysis. In Ser, Cys, and Thr proteases, an amino acid at the active site serves as the nucleophile forming a transient covalent intermediate, whereas in metallo- and Asp proteases, an active water molecule functions as nucleophile (adapted from Reference [13]). The base in covalent catalysis is usually a His, and in non-covalent intermediate, Asp/Glu and zinc (metalloproteinases) serve as acids and bases. The proteases are also classified as endo- and/or exo-proteinases based on their ability to cleave within or at the amino-/carboxy-terminus of the peptide chain.