

# ADVANCES IN PROTEIN CHEMISTRY

EDITED BY

**C. B. ANFENSEN, JR.**

*Laboratory of Cellular Physiology  
National Heart Institute  
Bethesda, Maryland*

**M. L. ANSON**

*London, England*

**KENNETH BAILEY**

*University of Cambridge  
Cambridge, England*

**JOHN T. EDSALL**

*Biological Laboratories  
Harvard University  
Cambridge, Massachusetts*

VOLUME 15



1960

ACADEMIC PRESS • New York and London

Copyright © 1960, by Academic Press Inc.

ALL RIGHTS RESERVED

NO PART OF THIS BOOK MAY BE REPRODUCED IN ANY FORM  
BY PHOTOSTAT, MICROFILM, OR ANY OTHER MEANS,  
WITHOUT WRITTEN PERMISSION FROM THE PUBLISHERS.

ACADEMIC PRESS INC.

111 FIFTH AVENUE  
NEW YORK 3, N. Y.

*United Kingdom Edition*

Published by

ACADEMIC PRESS INC. (LONDON) LTD.

17 OLD QUEEN STREET, LONDON, S.W. 1-

*Library of Congress Catalog Card Number 44-3853*

PRINTED IN THE UNITED STATES OF AMERICA

## CONTRIBUTORS TO VOLUME 15

- TH. BÜCHER, *Institut für Physiologische Chemie, Philipps-Universität, Marburg, Germany*
- HALVOR N. CHRISTENSEN, *Department of Biological Chemistry, The University of Michigan, Ann Arbor, Michigan*
- J. CRAVIOTO, *Group for Research on Infantile Malnutrition, Hospital Infantil de Mexico, Mexico City, Mexico, D. F.*
- R. CZOK, *Institut für Physiologische Chemie, Philipps-Universität, Marburg, Germany*
- KURT FELIX,\* *Institut für Vegetative Physiologie, Frankfurt, Germany*
- D. W. KUPKE, *Department of Biochemistry, School of Medicine, University of Virginia, Charlottesville, Virginia*
- JOAN M. L. STEPHEN, *Medical Research Council of Great Britain, Tropical Metabolism Research Unit, University College of the West Indies, Jamaica, W. I.*
- J. C. WATERLOW, *Medical Research Council of Great Britain, Tropical Metabolism Research Unit, University College of the West Indies, Jamaica, W. I.*

\* Deceased, August 2, 1960.

## CONTENTS

CONTRIBUTORS TO VOLUME 15.....	v
PREFACE.....	vii

### Protamines

KURT FELIX

I. Historical.....	1
II. Preparation of Protamines.....	2
III. Composition of the Protamines.....	12
IV. Occurrence of Protamines in Other Animals.....	17
V. General Properties of Protamines.....	20
VI. Heterogeneity of the Protamines.....	22
VII. Molecular Weights of Protamines.....	30
VIII. Amino Acid Sequences in Protamines.....	32
IX. Behavior of Protamines on Treatment with Proteolytic Enzymes.....	42
X. Protamine Compounds.....	43
XI. Are Protamines the Only Proteins in the Cell Nuclei of Fish Sperm?...	47
XII. Development of Protamines during Spermatogenesis.....	48
XIII. Biological Significance of Protamines.....	52
References.....	53

### Osmotic Pressure

D. W. KUPKE

I. Introduction.....	57
II. Fundamental Considerations.....	58
III. Applications.....	71
IV. Experimental Method.....	90
V. Summary.....	126
References.....	126

### Protein Malnutrition in Man

J. G. WATERLOW, J. CRAVIOTO, AND JOAN M. L. STEPHEN

I. Introduction.....	131
II. Clinical and Social Aspects.....	138
III. Biochemical Aspects.....	162
IV. General Conclusion.....	226
References.....	227

## Reactive Sites and Biological Transport

HALVOR N. CHRISTENSEN

I. Introduction .....	239
II. Nature of the Reactive Site .....	243
III. How Is Energy Supplied to Active Transport Reactions? .....	290
IV. Speculation as to the Basis of Humoral Control of Transport .....	296
V. Summary .....	304
References .....	305

## Crystallized Enzymes from the Myogen of Rabbit Skeletal Muscle

R. CZOK AND TH. BÜCHER

I. Introduction .....	315
II. Myogen .....	316
III. Preparative Procedures .....	323
IV. Enzymes .....	350
V. Cellular Physiology .....	395
References .....	407
AUTHOR INDEX .....	417
SUBJECT INDEX .....	436



# PROTAMINES

By KURT FELIX

Institut für Vegetative Physiologie, Frankfurt, Germany

I. Historical.....	1
II. Preparation of Protamines.....	2
A. Procedures of Kossel.....	3
B. Procedure of Ando and Yamasaki.....	6
C. Procedure of Block <i>et al.</i> .....	7
D. Procedure of Felix <i>et al.</i> .....	7
III. Composition of the Protamines.....	12
IV. Occurrence of Protamines in Other Animals.....	17
V. General Properties of Protamines.....	20
VI. Heterogeneity of the Protamines.....	22
A. Fractionation by Countercurrent Distribution According to Craig.....	22
B. Fractionation According to Solubility of the Picrates and Sulfates.....	24
C. Fractionation by Electrophoresis.....	25
D. Fractionation by Chromatography.....	26
VII. Molecular Weights of Protamines.....	30
VIII. Amino Acid Sequences in Protamines.....	32
A. Amino Terminal End Groups.....	33
B. Carboxyl Terminal End Groups.....	34
C. Internal Amino Acid Sequences in Protamines.....	36
IX. Behavior of Protamines on Treatment with Proteolytic Enzymes.....	42
X. Protamine Compounds.....	43
A. Compounds of Protamines in General.....	43
B. Compounds of Protamines with Nucleic Acids.....	43
XI. Are Protamines the Only Proteins in the Cell Nuclei of Fish Sperm?.....	47
XII. Development of Protamines during Spermatogenesis.....	48
XIII. Biological Significance of Protamines.....	52
References.....	53

## I. HISTORICAL

The first protamine was isolated by Miescher (1874) from salmon sperm. He extracted the fat-free sperm with dilute (1-2%) hydrochloric acid, neutralized the excess acid, and treated the solution with platinum chloride. All the protamine precipitated as the double salt of platinum and was free of sulfur and phosphorus. According to Piccard (1874) it still contained traces of guanine and hypoxanthine. The precipitate was insoluble in water, but soluble in an excess of hydrochloric acid. A similar platinum salt was obtained by him from the acid extract of carp sperm. While he

considered salmon sperm protamine to be a simple nitrogenous base, he observed that the corresponding material from carp sperm gave reactions for protein.

The protamine of salmon, later named salmine, which can be extracted with hydrochloric acid and precipitated with platinum chloride, corresponds to about 26.8 % of the dried sperm (Miescher, 1874).

About 20 years later, Kossel (1896) showed that the protamines are made up of amino acids and are therefore proteins. He repeated Miescher's experiments and obtained the same material from salmon sperm. However, since he was unable to obtain sufficient amounts of starting material for further investigations, he worked primarily on the sperm of sturgeon (*Acipenser sturio*) which were obtainable in adequate amounts from the Baltic Sea. He also simplified the preparation to some extent. The mature testes were minced, shaken with water, strained through a cloth, and the turbid liquid was acidified with a few drops of acetic acid, thereby conglomerating particulate elements so that they could be filtered off easily. The residue on the filter was extracted several times with alcohol and ether and then dried. The dried material was shaken with 5 volumes of 1 % sulfuric acid and any insoluble material was filtered off. The extraction was repeated three times with the same amount of sulfuric acid. The protamine sulfate was precipitated from the filtrate with a threefold excess of alcohol. The sulfate is very soluble in water and may be washed further by repeatedly redissolving it and reprecipitating it with alcohol. The salmine sulfate from salmon sperm was precipitated as the picrate from aqueous solution by sodium picrate and then reconverted into the sulfate. The protamine from sturgeon sperm (sturine) was quite similar to salmine, but differed from it in composition and in some of its reactions (Kossel, 1896). Among the hydrolysis products of sturine Kossel discovered histidine (Kossel, 1898). Along with his co-workers and students, he continued to isolate further protamines from the sperms of several species of fish.

## II. PREPARATION OF PROTAMINES

The preparation of protamines continues, in general, to follow the method outlined by Miescher and Kossel, which is based on the fact that the protamines occur as salts of nucleic acids in the nuclei of fish spermatozoa. The nucleic acid salts can be dissociated by treatment with dilute sulfuric acid (1 ml of concentrated  $\text{H}_2\text{SO}_4$  made up to 100 ml with water) or with 1 % hydrochloric acid. In view of the possibility that, during extraction, the nucleic acid may be partially broken down and some of its bases may go into solution, the protamine must be precipitated and washed. Although the sulfate of the protamine can be precipitated with 3 volumes of alcohol, it is not insoluble enough to be washed thoroughly. To precipitate the chloride, acetone is needed.

The combination with heavy metals has not proven successful, since the metals have to be removed as their sulfides and the precipitates occlude considerable amounts of nitrogenous material. Besides platinum chloride, mercuric nitrate in nitric acid solution can be used for precipitation of protamines, or alkaloid precipitants may be used. The most suitable products are the picrates since they are highly insoluble and can be washed thoroughly, even with 0.1 *N* sodium hydroxide which will remove any purine picrate that may also have precipitated. Precipitation is carried out, not with alcoholic picric acid, but with aqueous sodium picrate which reacts almost exclusively with the strongly basic protamines and with very few other bases. The picrates are taken up in sulfuric acid and the picric acid is removed by extraction with ether or toluene. The sulfates are then reprecipitated with alcohol.

The best starting material for the preparation of protamines is freshly drawn milt from fish that are ready to spawn, since it contains only one type of cells, the spermatozoa. If sufficient milt cannot be obtained, the mature testes are used as described above in the historical introduction. Occasionally these contain other cells besides the sperm, such as earlier stages of sperm and connective tissue cells; the more immature the testes, the less homogeneous the cellular material. The protamines obtained therefrom may still be contaminated with other proteins or their decomposition products. It is important to disperse the minced testes in 4 or 5 volumes of water and to shake the mixture vigorously. The material is then strained through muslin or a wire sieve. The extract is acidified with acetic acid until it just gives a blue color with Congo red paper. The precipitate is collected on fluted filter paper, freed from fatty material, and dried as described above. To prepare the protamine from the dried material, Kossel (1929) describes two procedures. The first of these, based on extraction with acid, is preferred by most authors. The original directions are as follows.

#### *A. Procedures of Kossel*

(1) Approximately 100 gm of the powdery material are shaken with 500 ml of 1 % sulfuric acid for  $\frac{1}{2}$  hr and then filtered. The residue on the filter is extracted several times, until an aliquot of the sulfuric acid extract gives no visible precipitate with alcohol. The extraction should take no longer than 1 day; if the sperm material is left in contact with the sulfuric acid for a longer period, some decomposition of nucleic acids and formation of interfering substances will result. The protamine sulfate is precipitated from the sulfuric acid extract by addition of 3 volumes of alcohol and the supernatant is decanted or siphoned off. The precipitate is then dissolved in a little hot water and the alcohol precipitation is repeated. The total amount of precipitate obtained from 100 gm of air-dried sperm mate-



rial is then dissolved in about 1 or 2 liters of hot water. On cooling, a small amount of the sulfate separates out as a yellow or brownish oil. This most insoluble fraction of the protamine sulfate is separated from the supernatant liquid and the latter evaporated to a small volume. The bulk of the oil is then obtained by means of a separatory funnel, the middle fraction of the oil being the most pure.

The protamine sulfate, after being dissolved in warm water, is purified further by precipitation with sodium picrate, followed by the removal of picric acid by shaking the well-washed precipitate with toluene in the presence of an excess of sulfuric acid, and then protamine sulfate is precipitated from the sulfuric acid solution with alcohol. The alcohol precipitation is then repeated. The consistency of the precipitate is dependent upon the degree of acidity of the solution; if the amount of acid is inadequate, a turbid solution is obtained. In that case, flocculation can be effected by careful, dropwise addition of sulfuric acid. The formation of a sticky precipitate indicates that too much acid is present and solution in water and precipitation with alcohol must be repeated. The protamine sulfate should come down as a powdery, pure white precipitate. Then this is washed with alcohol and ether and dried in a desiccator. The yield, for mature herring testes, should be 15-20 % of the air-dried sperm material.

When larger quantities are to be prepared, the amount of alcohol may be reduced by evaporating the sulfuric acid extracts to a smaller volume. However, it should be noted that the extract contains small amounts of other tissue constituents which are removed by the alcohol precipitation described above, but which, during evaporation, will partially decompose and adhere to the preparation.

Then the sulfuric acid extracts are treated as follows: the combined extracts are neutralized with barium hydroxide and evaporated in a water bath. The extracts are reduced to a volume suitable for the separation of the oil, which is then purified with picrate as described above.

This method of preparation may be used for most protamines. With the protamines of carp sperm, however, the oily separation of the sulfate does not occur.

(2) In the second procedure, the dried sperm material is treated with cupric chloride solution. The protamines pass into solution, while the nucleic acids form insoluble copper salts. The protamines are precipitated from the filtrate with sodium picrate. This method was developed by Nelson-Gerhardt (1919) of Kossel's Institute, making use of data by Schmiedeberg (1899) and Malenik (1908). Since the method is still used by some authors, the details of the procedure are described below.

One hundred grams of dried sperm material, prepared as described above, are incubated in an oven with a solution of 100 gm of cupric chloride in 1

liter of water. Incubation is continued for 3 days, with occasional shaking. Then the supernatant liquid is filtered off with suction. The residue is resuspended in water three times and filtered with suction. Washing of the residue is continued until an aliquot of the filtrate, when tested with concentrated sodium picrate solution, does not show a precipitate.

Concentrated sodium picrate solution is added, with stirring, to the combined filtrates and washings until the precipitate sticks together and settles out quickly. The precipitate is filtered off, washed with a little water to which some sodium picrate has been added, and dissolved while still wet, with slight heating, in a mixture of 1 volume of acetone and 3 volumes of water. If possible, the solution is filtered and the filtrate treated with half the volume of alcohol. Sulfuric acid (20% v/v) is then added dropwise, with stirring, until no further precipitate forms. An excess of sulfuric acid will render the precipitate smeary and soluble and therefore must be avoided. The supernatant liquid is decanted through a filter and the residue treated with absolute alcohol, thus being made hard and crumbly. The precipitate is decanted several times with alcohol and finally with ether, filtered off, and washed with ether. Then the ether is removed in a desiccator. Further purification of the preparation may be accomplished, making use of its resistance to the action of pepsin. An aqueous solution of 10 gm of protamine sulfate is digested for 24 hr at 37°C in about 250 ml of water to which 0.1 gm of commercial pepsin and 0.5 gm of HCl have been added. The digestion mixture is neutralized with sodium carbonate and the protamine picrate is precipitated, as described above, and converted to the sulfate, which is then made to solidify, and is redissolved as in procedure (1).

#### 1. *Conversion of Protamine Picrate to Protamine Ester Hydrochloride*

From picrate suspended in dilute sulfuric acid, picric acid can be removed only by repeated extraction with toluene or ether. This removal may be accomplished faster by suspending the dry picrate in 100 volumes of anhydrous 0.2 N hydrochloric acid in methyl alcohol and letting it stand for 24 hr, with occasional stirring. The residue is filtered off with suction, dried, shaken with fresh methyl alcoholic hydrochloric acid (about one-fourth the amount used before), allowed to stand 24 hr, filtered as before and the residue discarded. The combined filtrates are treated with 3 volumes of anhydrous ether, allowed to stand overnight in the cold room, and the protamine methyl ester hydrochloride precipitate is filtered off with suction. The crude product is redissolved in the 0.2 N methyl alcoholic hydrochloric acid, reprecipitated with 3 volumes of anhydrous ether, and finally dried with phosphorus pentoxide under vacuum (Felix and Dirr, 1929; Felix and Mager, 1937c; Rauen *et al.*, 1952).

## 2. Purification of Protamine Sulfate by Refractionation (*Umscheiden*)

Waldschmidt-Leitz and co-workers employ a "refractionation" procedure for the purification of clupeine sulfate. The alcohol and ether dried testes are extracted with cupric chloride (as described in Section II, A), precipitated with sodium picrate, the picrate converted to the sulfate, and the latter precipitated with acetone. A saturated solution of the sulfate at 30–40°C (about 7%) is prepared and this is cooled to 0°C, with the resultant separation of the clupeine sulfate as an oily liquid. This separation is repeated many times over a period of several weeks. (Rauen *et al.*, 1952; Waldschmidt-Leitz *et al.*, 1931; Waldschmidt-Leitz and Vohr, 1954; Waldschmidt-Leitz and Mindemann, 1956; Waldschmidt-Leitz and Gudernatsch, 1957.)

### B. Procedure of Ando and Yamasaki

Ando *et al.* (1957c) and Yamasaki (1958) recommend the following modification of Kossel's procedure.

Frozen herring milt (410 gm) is finely powdered and stirred into 1.6 liters of 0.05 *M* sodium citrate, pH 7. Then it is strained through cheesecloth and the filtrate is centrifuged for 10 min at 4000 rpm. The pellet is suspended again in 1.6 liters of the citrate solution, stirred for 30 min, and then centrifuged at a lower speed (1300 rpm) for 2 min. The brown sediment (tissue debris) is discarded and the supernatant is centrifuged for 5 min at 4000 rpm.

The wet sperm heads are stirred vigorously in 4 liters of 2 *M* NaCl containing 0.05 *M* sodium citrate. This dissolves the nucleoclupeine. While stirring, 400 ml of a saturated  $\text{CuSO}_4$  solution are added to the mixture, thus precipitating the nucleic acids as their copper salts. Stirring is continued for additional 30 min and the mixture is allowed to stand overnight in the cold room. The copper salts of the nucleic acids are then filtered off and washed with water. From the combined filtrates and washings (6 liters, the clupeine is precipitated as its picrate by treatment with 0.125 *M* sodium picrate (400 ml).

Ten grams of the picrate are dissolved at room temperature in 100 ml of aqueous acetone solution (67% w/v) and the insoluble material is filtered off. At this point, fractionation can be accomplished by cooling, the less soluble picrate coming down as an oil. Both picrates are converted to the sulfates by addition of 2 *N* sulfuric acid, and the sulfates are precipitated with alcohol. Then they are dissolved in water and reprecipitated with alcohol. By means of an anion exchanger (Amberlite IRA-400, Cl form), they may be converted to the hydrochlorides.

### C. Procedure of Block *et al.*

Block and Bolling recommend a different procedure. From a suspension of spermatozoa, the heads are precipitated with 1 % citric acid, extracted with 0.2 % hydrochloric acid, and the extract is heated to the boiling point. The pH is adjusted to 8 with ammonia and the precipitate is filtered off. The protamine is precipitated from the filtrate with the optimal amount of 33 % metaphosphoric acid, and the protamine metaphosphate is converted to the sulfate with 1 N sulfuric acid. The sulfate is then precipitated in the usual manner with alcohol or acetone and dried. This method, while perhaps producing a larger yield, might also cause some hydrolysis (Block *et al.*, 1949; Rauen *et al.*, 1953).

### D. Procedure of Felix *et al.*

Recent procedures use purified spermatozoa nuclei as the starting material. From these, nucleoprotamines are prepared, and from them, the protamines are obtained.

#### 1. Preparation of Fish Sperm Nuclei

a. *Principle.* On treating the filtrate from an aqueous suspension of minced testes with acetic acid, the sperm heads are precipitated as nuclei together with some cytoplasmic material. A purer and more homogeneous preparation is obtained by starting with the nuclei after complete removal of any cytoplasm. This is done by placing the spermatozoa in distilled water. Within a few minutes, the cytoplasm in the head and tail begins to swell. The fibrils of the latter form loops, while the nucleus itself remains unaltered. Then the nuclei can be separated from the cytoplasm and tails by homogenizing the suspension briefly (5 min) and centrifuging. The fibrils and cytoplasm remain in the supernatant, which gives a reaction for tyrosine, tryptophan, cystine, and ribonucleic acid. The nuclei are in the sediment. They are washed in the centrifuge with distilled water until the above reactions are negative, indicating that the cytoplasm has been completely removed (Felix *et al.*, 1951b). The above amino acids, as well as ribonucleic acid, are missing from the nuclei of trout and herring sperm. This method of preparation does not impair the biological potency of the nuclei; it has been possible, in a few cases, to use them to fertilize the corresponding fish eggs and to promote the development of normal fish (Felix *et al.*, 1952b) (see Figs. 1-5).

When the nuclei are precipitated with acetic acid, they are gray rather than pure white and sometimes give a reaction for tyrosine (Steudel, 1911; Felix *et al.*, 1951b).






FIG. 1. Trout sperm in Ringer solution. Osmium fixation. Magnification:  $\times 11,000$ . Electron micrographed by Drs. W. Lippert and O. Hug of the Max Planck Institute for Biophysics, Frankfurt/Main.




FIG. 2. Trout sperm suspended in distilled water for 10 min. Osmium fixation. Magnification:  $\times 10,000$ . Electron micrograph by Drs. W. Lippert and O. Hug of the Max Planck Institute for Biophysics, Frankfurt/Main.






FIG. 3. Trout sperm suspended in distilled water for 30 min. Osmium fixation. Magnification:  $\times 8,500$ . Electron micrograph by Drs. W. Lippert and O. Hug of the Max Planck Institute for Biophysics, Frankfurt/Main.




FIG. 4. Trout sperm: isolated nuclei after repeated washings with distilled water. Osmium fixation. Magnification:  $\times 9,000$ . Electron micrograph by Drs. W. Lippert and O. Hug of the Max Planck Institute for Biophysics, Frankfurt/Main.

*b. Method.* The best starting material is freshly drawn milt, adequate amounts of which may be obtained from trout in fish hatcheries. Care must be taken to avoid contamination with blood or feces. Two hundred trout yield about 180 ml of milt. If it cannot be used immediately, the milt may be stored overnight in a cold room without loss of mobility of the sperm. The total amount of milt collected is added, with stirring, to about 4 liters




FIG. 5. Trout sperm: nuclei treated on the blend with 10% NaCl and subsequently washed with distilled water. Magnification:  $\times 6,000$ . Electron micrograph by Drs. W. Lippert and O. Hug of the Max Planck Institute for Biophysics, Frankfurt/Main.

of ice-cold distilled water and cytolysis is allowed to proceed for about 20 min. The supernatant opalescent liquid is somewhat yellowish, probably due to lactoflavine, while the very loosely packed sediment is pure white and consists almost exclusively of the nuclei. It is taken up in 500 ml of water and evenly dispersed, strained through several layers of muslin, and then homogenized in small portions for about 3 min. The combined homogenates are centrifuged at 3000 rpm. The pellet of loose nuclei is repeatedly washed with distilled water and centrifuged until the washings no longer give a reaction for tryptophan. This procedure yields very pure nuclei; however, the losses are fairly large, since the supernatant always

contains some suspended nuclei. When 0.005 % citric acid is used for the suspension, the supernatant remains clear but the nuclei stick together. If possible, the nuclei should be used immediately. To store them, they are dried with acetone and ether, which causes them to shrink a little.

## 2. Preparation of Nucleoprotamine

The nuclei of trout sperm consist almost exclusively of the salt of protamine with deoxyribonucleic acid (DNA), i.e., of nucleoprotamine.

Miescher (1874) observed that in a 10–15 % sodium chloride solution, sperm is transformed into a translucent gelatinous mass. Mirsky and Pollister (1942) were the first to use a 10 % sodium chloride solution for extraction of nuclear material from sperm.

TABLE I  
*Analyses of Sperm Nuclei and Fibers of Nuclei*

Source	N	P	N/P	Arginine	P/Arginine
<i>Analyses of sperm nuclei</i>					
<i>Salmo irideus</i>	19.67	5.87	3.35	30.86	1:0.94
<i>Salmo trutta</i>	19.80	5.72	3.46	30.42	1:0.95
<i>Salmo fontinalis</i>	19.78	5.88	3.36	30.44	1:0.92
<i>Clupea harengus</i>	19.73	5.76	3.42	30.42	1:0.94
<i>Analyses of fibers of nuclei</i>					
<i>Salmo irideus</i>	19.52	5.65	3.43	30.60	1:0.96
<i>Salmo trutta</i>	19.67	5.71	3.44	30.20	1:0.94
<i>Salmo fontinalis</i>	19.67	5.73	3.43	30.15	1:0.94
<i>Clupea harengus</i>	19.57	5.68	3.44	30.77	1:0.96

If the wet pellet of nuclear material obtained from 180 ml of trout milt is taken up in 2000 ml of 10 % sodium chloride solution and homogenized briefly, it will dissolve almost immediately without leaving any residue. (If acetone-dried nuclei are used, a small insoluble residue remains.) The solution is stirred in an ice bath for 12 hr. On addition of 9 volumes of water to this solution the entire nuclear preparation (that is, the nucleoprotamine) precipitates in long threads which can be wrapped around a glass rod and washed with water. They are washed further with alcohol and dried with ether.

As shown in Table I, the nuclei and fibers contain equal amounts of nitrogen, phosphorus, and arginine.

If desired, the nucleoprotamine may be redissolved in 10 % sodium chloride solution and reprecipitated with water before proceeding to the next step. It is not recommended that the nucleoprotamine be reprecipitated more often since it gradually loses its solubility.

### 3. Preparation of Protamine Hydrochloride

The dried and powdered nucleoprotamine is extracted several times with small portions of ice-cold 0.2 % hydrochloric acid. About 150 ml of acid are needed for 1.5 gm of nucleoprotamine. The extract is treated with 20 volumes of acetone. At first, the solution merely becomes cloudy, but on standing at 4°C for 12 hr, the protamine hydrochloride collects on the walls of the container. It is dissolved in a little water, frozen, and dried under high vacuum.

In general, the preparations are pure enough so as not to require further purification by way of the picrates. However, this should be done in case of doubt.

There is still another way in which protamine may be prepared from nucleoprotamine (Goppold-Krekels and Lehmann, 1958). The nucleoprotamine is again dissolved in a 10 % sodium chloride solution and then the solution is saturated with sodium chloride. On standing at 4°C, the protamine gradually settles out. This process may be speeded up by centrifugation. A clear supernatant is obtained, and also an oily sediment consisting of protamine and sodium chloride. This is dissolved in 5 % trichloroacetic acid and then the concentration of the trichloroacetic acid is increased to 10 %, causing the protamine to precipitate free of sodium chloride. For certain purposes (see Section VI, C), the oily sediment may be used directly after first dissolving it in a little water and freeze-drying it. This preparation is referred to as "clupeine oil."

### III. COMPOSITION OF THE PROTAMINES

The protamines contain very few kinds of amino acids. All of them have been shown to contain arginine, alanine, and serine; proline and valine occur in most of them, and in many glycine and isoleucine are present. Individual protamines contain histidine, lysine, threonine, and aspartic and glutamic acids. Tyrosine has been demonstrated by Kossel only in thynnine from tunny and in cyclopterine from *Cyclopterus lumpus* (Baltic Sea). None of the protamines analyzed thus far contain tryptophan, phenylalanine, or sulfur-containing amino acids. An exception is mugiline  $\beta$ , which contains methionine.

In all protamines, the basic amino acids predominate to the extent that there are about two basic amino acids to every nonbasic amino acid. The most important of the former is arginine, which, in most protamines, constitutes about two-thirds of all amino acids. Several protamines contain still another basic amino acid, either histidine or lysine, and in a few, all three of them occur. Based on this, Kossel divided the protamines into three groups: monoprotamines, containing only arginine; diprotamines, containing lysine or histidine as well as arginine; and triprotamines, con-